

THE GIFT OF Bur. of Bort. Sal.



BENEFAL LIBRARY, UNIV. OF MIGH. JUN 16 1905

No. 27.-APRIL, 1905

DEPARTMENT OF THE INTERIOR
BUREAU OF GOVERNMENT LABORATORIES

A REVIEW OF THE IDENTIFICATIONS OF THE SPECIES DESCRIBED IN BLANCO'S FLORA DE FILIPINAS

BY

ELMER D. MERRILL, BOTANIST

MANILA
BUREAU OF PUBLIC PRINTING
1905

PREVIOUS PUBLICATIONS OF THE BUREAU OF GOVERNMENT LABORATORIES.

- No. 1, 1902, Biological Laboratory.—Preliminary Report of the Appearance in the Philippine Islands of a Disease Clinically Resembling Glanders. By R. P. Strong, M. D.
- No. 2, 1902, Chemical Laboratory.—The Preparation of Benzoyl-Acetyl Peroxide and Its Use as an Intestinal Antiseptic in Cholera and Dysentery. (Preliminary notes.) By Paul C. Freer, M. D., Ph. D.
- No. 3, 1903, Biological Laboratory.—A Preliminary Report on Trypanosomiasis of Horses in the Philippine Islands. By W. E. Musgrave, M. D., Acting Director Biological Laboratory, and Norman E. Williamson, Assistant Bacteriologist Bureau of Government Laboratories
- No. 4, 1903, Serum Laboratory.—Preliminary Report on the Study of Rinderpest of Cattle and Carabaos in the Philippine Islands. By James W. Jobling, M. D., Director of the Serum Laboratory.
- No. 5, 1903, Biological Laboratory.—Trypanosoma and Trypanosomiasis, with Special Reference to Surra in the Philippine Islands. By W. E. Musgrave, M. D., Acting Director Biological Laboratory, and Moses T. Clegg, Assistant Bacteriologist Biological Laboratory.
- No. 6, 1903.—I, New or Noteworthy Philippine Plants. II. The American Element in the Philippine Flora. By Elmer D. Merrill, Botanist.
- No. 7, 1903, Chemical Laboratory.—The Gutta Percha and Rubber of the Philippine Islands. By Penoyer L. Sherman, jr., Ph. D., Chemist, Chemical Laboratory.
- No. 8, 1903.—A Dictionary of the Plant Names of the Philippine Islands. By Elmer D. Merrill, Botanist.
- No 9, 1905, Biological Laboratory.—A Report on Hemorrhagic Septicæmia in Animals in the Philippine Islands. By Paul G. Woolley, M. D., and J. W. Jobling, M. D.
- No. 10, 1903, Biological Laboratory.—Two Cases of a Peculiar Form of Hand Infection (Due to an Organism Resembling the Koch-Weeks Bacillus). By John R. McDill, M. D., and Wm. B. Wherry, M. D.
- No. 11, 1903, Biological Laboratory.—Entomological Division, Bulletin No. 1, Preliminary Bulletin on Insects of the Cacao. (Prepared especially for the benefit of farmers.) By Charles S. Banks, Entomologist Bureau of Government Laboratories.
- No. 12, 1903, Biological Laboratory.—Report on Some Pulmonary Lesions Produced by the Bacillus of Hemorrhagic Septicæmia of Carabaos. By Paul G. Woolley, M. D.
- No. 13, 1904, Biological Laboratory.—A Fatal Infection by a Hitherto Undescribed Chromogenic Bacterium: Bacillus aureus fœtidus. By Maximilian Herzog, M. D.
- No. 14, 1904, Serum Laboratory.—Texas Fever in the Philippine Islands and the Far East. By J. W. Jobling, M. D., and Paul G. Woolley, M. D. Biological Laboratory.—Entomological Division, Bulletin No. 2, the Australian Tick (Boophilus Australia Fuller) in the Philippine Islands. By Charles S. Banks, Entomologist.
- No. 15, 1904, Biological and Serum Laboratories.—Report on Bacillus Violaceus Manilæ: A Pathogenic Micro-Organism. By Paul G. Woolley, M. D.
- No. 16, 1904, Biological Laboratory.—Protective Inoculation Against Asiatic Cholera: An Experimental Study. By Richard P. Strong, M. D.
- No. 17, 1904.—New or Noteworthy Philippine Plants. II. By Elmer D. Merrill, Botanist.
- No. 18, 1904, Biological Laboratory.—Part I. Amebas: Their Cultivation and Etiologic Significance. By W. E. Müsgrave, M. D., and Moses T. Clegg. Part II. Treatment of Uncomplicated Intestinal Amebiasis (Amebic Dysentery) in the Tropics. By W. E. Musgrave, M. D.

(Continued on third page of cover.)

No. 27.—APRIL, 1905


DEPARTMENT OF THE INTERIOR BUREAU OF GOVERNMENT LABORATORIES

A Review of the Identifications of the Species Described in Blanco's Flora de Filipinas

BY

ELMER D. MERRILL,
BOTANIST

MANILA
BUREAU OF PUBLIC PRINTING
1905



LETTER OF TRANSMITTAL.


DEPARTMENT OF THE INTERIOR,
BUREAU OF GOVERNMENT LABORATORIES,
OFFICE OF THE SUPERINTENDENT OF LABORATORIES,
Manila, P. I., July 25 1904.

SIR: I have the honor to transmit herewith a paper entitled "A Review of the Identifications of the Species Described in Blanco's Flora de Filipinas," by Elmer D. Merrill, botanist.

I am, very respectfully,

Paul C. Freer, Superintendent of Government Laboratories.

Hon. DEAN C. WORCESTER, Secretary of the Interior.



A REVIEW OF THE IDENTIFICATIONS OF THE SPECIES DESCRIBED IN BLANCO'S FLORA DE FILIPÍNAS.

By Elmer D. Merrill, Botanist.

INTRODUCTION.

"Il est à regretter que ces révérends ecclésiastiques (Blanco and Loureiro) et même le Père Plumier, leur prédécesseur, ne se soient pas contentés d'écrire des homélies." Thus A. DeCandolle¹ expressed his opinion of Blanco's botanical work, an opinion in which all botanists who have occasion to study the Philippine flora will heartily concur.

While we must agree with DeCandolle's opinion regarding Blanco's work, and while undoubtedly the publication of Blanco's Flora de Filipinas has retarded rather than advanced our knowledge of the Philippine flora, still much credit is due to this author for prosecuting his investigations under most unfavorable conditions. The preparation, by the unaided efforts of one man, of a work of the magnitude of Blanco's Flora de Filipinas, with no correspondence with contemporary European botanists, with few reference books, and no authentically named botanical specimens, and where almost every fact noted is the result of the personal observation of the author, is in itself a monument to the zeal and ability of Blanco, and, after a consideration of the circumstances under which the work was accomplished, one can but wonder that the author did not commit more errors.

Blanco's generic and specific descriptions, judged from the standpoint of modern botany, are certainly very imperfect and incomplete, yet many of them are equal or even superior to many of those published by this contemporaries. However, while on the one hand the types of the species described by various European botanists are still perserved in various herbaria in Europe, and an examination of the type specimen will usually clear up any question existing as to the identity of any doubtful species, on the other the types of all of Blanco's species have been destroyed. It is certain that Blanco had at least a working herbarium, but unfortunately, after his death, the value of his collection was not recognized. Accordingly, to-day we are dependent entirely on his imperfect descriptions for the identification of his species.

In the two editions of the Flora de Filipinas, Blanco described no less than 1,127 species and varieties, of which about 623 were proposed by him as new to science, and 219 were correctly and 285 incorrectly identified with species of other authors. A large percentage of Blanco's new species are to-day unknown, but the greater proportion will eventually be reduced as synonyms of those described by other authors. At the present time but 90 are known to be valid. Considering the early date when Blanco did his work and the fact that the Philippine flora is especially rich in endemic species, the percentage of valid ones among those proposed by Blanco is exceedingly small, but this is due to several factors. Most of Blanco's botanical material was from the immediate vicinity of towns or from thickly settled regions, and accordingly a great proportion of his species were based on widely distributed and cosmopolitan forms. He had practically no material from the higher mountains, where the flora is especially rich and interesting; he made no attempt to make his work exhaustive for any one family or group—in fact, he almost entirely ignored such large and critical families as the Orchidacea, Graminea, Cyperacea, etc. He endeavored particularly to include all species of economic value, especially cultivated plants, and he erroneously identified with species of other authors very many which would have been valid had he given them distinctive specific names.

THE FIRST EDITION OF BLANCO'S FLORA DE FILIPINAS.

In the year 1837 there was published in Manila the first edition of Blanco's Flora de Filipinas, an octavo volume of 887 pages, with an introduction of 78 additional pages, the latter containing an explanation of the principles of botany, of the technical terms used in the text, of the Linnean system of classification, and an enumeration of the genera treated in the work, arranged in classes

according to the latter. The work is entirely in the Spanish language, and contains the descriptions of 903 species and varieties under Latin and 31 under native names only. This work is the only one of its kind treating of the flora of the Philippines, and on account of the many imperfect descriptions, and also due to the fact that the types of Blanco's species were never accessible to European botanists, the publication of this work has caused considerable confusion in the subsequent treatment of the flora of the Philippines and the adjacent Malayan region, although had the types of Blanco's species been preserved his work would have compared very favorably with that of many of his contemporaries. According to Blanco's statement he had few botanical books, chiefly Linnaeus' Systema Vegetabilium, Jussieu's Genera Plantarum, Aublet's Histoire des Plantes de la Guiane françois, and some of the earlier volumes of DeCandolle's Prodromus.

THE SECOND EDITION.

In 1845, shortly after Blanco's death, a second edition of his Flora de Filipinas was published, an octavo volume of 619 pages, very similar to the first one in its style and arrangement, but scarcely an improvement. In the second edition 1,131 species and varieties are described under Latin and 27 under native names only. Six described in the first are excluded from the second. while 198 are added in the second, and no less than 247 described in the first are included in the second under different generic or specific names. As the descriptions of the species in the second edition contain no references to those in the first, it was in many cases difficult to correlate the species of the two editions up to the time of publication of the Novissima Appendix. In some cases the specific or generic names were arbitrarily changed by Blanco, while in others he attempted to identify certain species which were treated of in the first edition with those which had previously been described by other authors. Very few of these changes are correct. In other cases in the first edition he described species under their proper names, which in the second he erroneously altered.

It is frequently rather difficult to determine definitely whether or not Blanco intended to describe a species as new or to identify the same with those of other authors. In some cases, but not always, he indicates these new species. Authorities for species are

¹Blanco died in Manila April 1, 1845.

not given, and only when he gives a citation at the end of a description, to Linnaeus, DeCandolle, or other authors, can we be sure that he intended these to represent species of other authors.

Blanco had little conception of geographical distribution, and on the one hand identified many endemic species with those of Asia, and even of tropical America, in the latter being influenced by Aublet's Histoire des Plantes de la Guiane françoise, of which work he had a copy, while on the other hand he described as new many species, native of tropical America, introduced as weeds or as cultivated plants. The second edition like the first follows the Linnean system of classification, with some changes in the sequence of families and genera, but is marred by many typographical errors, which are probably due to the fact that Blanco's death occurred some months before the book was printed, and the correction of the proof devolved on some person who had little or no botanical knowledge.

THE THIRD EDITION.

The third or Augustinian edition of Blanco's Flora de Filipinas was published in Manila between the years 1877 and 1883, under the auspices of the Order of Augustinians. It is a pretentious publication of four volumes of text and two volumes of plates, and is the work of two Augustinian priests, Andreas Naves and Celestino Fernandez-Villar. The first three volumes practically consist of only a reprint of the second edition with a Latin translation, although in rare instances names that are used for species in the first edition are substituted for those in the second. In the third no page references are given to species in the first and second, and accordingly the third edition is of little practical value, and shows no improvement over the second except that it is more available to botanists who have no knowledge of the Spanish language. authors of the third edition practically succeeded only in extending the data included in the small, convenient octavo volume of the second, through three bulky folio volumes, and in so doing added nothing to our knowledge of Blanco's species or to our knowledge of the Philippine flora. Volume 1 of the third edition was published in the year 1877 and contains 350 pages, including the species from Monandra, Monogynia to Hexandra, Hexagynia; volume 2 was published in the year 1878 and consists of 418 pages, including the species from Octandria, Monogynia, to Polyadelphia, Polyandra, while volume 3, published in 1879, consists of 271 pages and contains those from Syngenesia to Cryptogamia, and the uncertain species. The fourth volume and the two of plates have no direct connection with the first and second editions of Blanco's Flora de Filipinas, the Novissima Appendix, one of the papers in the fourth volume, being the only part of interest to us here.

THE NOVISSIMA APPENDIX.

The Novissima Appendix, the third and last article in the fourth volume of the third edition of Blanco's Flora de Filipinas, consists of 375 pages and was published in Manila between the years 1880 and 1883, the last part of the work being printed in June, 1883. This paper in the work of Celestino Fernandez-Villar and Andreas Naves, and upon it the present paper is largely based. object of the publication of the Novissima Appendix was not only to reduce the species of Blanco's Flora de Filipinas to known ones of older authors but also to enumerate all the species of plants then known from the Philippines. In both cases, but more especially in the former, the authors fell far short of the mark. The authors of the Novissima Appendix made the serious mistake of identifying Blanco's Philippine species with those of other authors from British India and the Malayan region, and, except in the cases of widely distributed and well-known species, nearly all of their identifications are erroneous; and in so identifying Blanco's species they only succeeded in further complicating Philippine botany by crediting to the Archipelago scores of species that do not extend to the Philippines. Of the 1,127 species and varieties described by Blanco, 5 only were considered by the authors of the Novissima Appendix to be unknown, 1,122 being for most part reduced without question and without discussion to species of other authors. In the present paper it is considered that no less than 289 of Blanco's species are unknown at the present time, notwithstanding the work that has been done on the Philippine flora since the publication of the Novissima Appendix, and in no less than 490 other cases it has been found necessary to change or doubt the identifications of Villar and Naves. It is evident that these authors accomplished their work in the library rather than in the field and in the herbarium, and that they reduced Blanco's species to those of other authors not so much by the study of botanical material as by the examination and comparison of descriptions. It is also very evident that, no matter how well they understood Blanco's species, they had little knowledge of the extra-Philippine flora, and very little conception of geographical distribution. While comparatively few of the specific identifications are correct, it is a remarkable fact that very few errors are to be found in their generic identifications.

In the preparation of the manuscript for the Novissima Appendix the treatment of the Dicotyledons is by Villar, and follows the arrangement of Bentham and Hooker's Genera Plantarum. The Monocotyledons were commenced by Naves and completed by him to page 307 (Fimbristylis nutans Vahl.), the remainder of the work being by Villar. The last part of the Genera Plantarum was not published in time to be utilized in the treatment of the Monocotyledons.

The Novissima Appendix serves an excellent purpose in that it for the first time correlates the species of the three editions of Blanco's Flora de Filipinas, page references being given to all those described by Blanco in the three editions. Previous to the publication of the Novissima Appendix it was very difficult to correlate the species of the first two editions of the Flora de Filipinas owing to the fact that Blanco changed the names of so many species in the second edition, giving no references to the names found in the first. The Novissima Appendix also gives us a clew to the proper families and genera of Blanco's species, for it is rare indeed that they are referred to wrong genera.

The identifications of Villar and Naves have been used as a basis of the present work, and these have been accepted in all cases where the facts seem to warrant. Identifications have been discarded in some cases by a careful comparison of descriptions, and in others by a comparison of descriptions in connection with the known geographical distribution of species.

GENERA.

In the first edition of the Flora de Filipinas Blanco described seventeen new genera, four of which were discarded by him in the second, while in the second he described six additional new ones, making a total of twenty-three. Of these twenty-three genera, three—Lunasia, Palaquium, and Malaisia—are recognized to-day as valid; two—Balingayum and Soala—are unknown or uncertain, while the remaining eighteen have been satisfactorily reduced as synonyms of previously established genera of other authors. Of

the two unknown, Soala belongs to the Anonacea, but can scarcely be Cyathocalya, to which it has been reduced by later authors. Balingayum has been reduced by various authors questionably to Erythropalum, a genus of the Olacacea, while Fernandez-Villar retains it as a distinct genus of the Onagracea. Specimens are necessary to determine just what these genera may be, and should be found in the localities cited by Blanco in his descriptions.

SPECIES.

Of the 1,127 species and varieties described by Blanco in the two editions of his Flora de Filipinas, so far as we have been able to determine them, 289 are to-day either unknown or at least very imperfectly known from Blanco's descriptions only. These species which Blanco erroneously identified with those of older authors are of comparatively little importance to us, whether or not such are ever identified they will have no effect on the nomenclature of Philippine botany. However, with respect to those which Blanco described as new to science, it is very important to have them properly identified at the earliest date possible. Of the 289 species at present unknown, 65 Blanco erroneously identified with those of other authors, while 224 were first described by him. Of the total of 1,127 species, 623 were described by Blanco as new, 219 were correctly and 285 incorrectly identified with those of other authors. Of the 623 species described by Blanco as new, so far as known to-day, 297 have been satisfactorily reduced as synonyms of those previously described, 236 are quite unknown, while of 90 species Blanco's descriptions are the first known, and his specific names are or should be retained. It is confidently expected that we shall in the future be able to identify a large percentage of these unknown species, many of which will prove to be valid, Blanco's names replacing names of those at present credited to other authors, while many will be satisfactorily reduced as synonyms of previously described species.

THE IDENTIFICATION OF BLANCO'S SPECIES.

In identifying Blanco's species numerous factors must be considered, but it is at once evident that the identification of those which are at present unknown will depend primarily on the field worker, and accordingly the present paper has been prepared with the view of aiding collectors in the field. Blanco frequently gives much more

space and attention to the description of gross characters and to the discussion of the economic uses of his species than to an enumeration of strictly technical characters, data and information which the ordinary dried botanical specimen does not show unless the collector's notes are very complete. Accordingly, European botanists, working only with dried material, have been unable to identify many of Blanco's species, but investigators working in the Philippines have been able to do this with little trouble because they have the advantage of knowing the various species as they appear in the field, and have to a greater or less extent the opportunity of collecting in the localities from which Blanco received his material, and of securing data regarding the economic uses of the different species, the native names, and other information which is often essential to correct identification. However, botanists working in the Philippines have been at a disadvantage in that they have not had access to the types of Malayan species in the various European herbaria, and accordingly have in many cases erroneously reduced those of Blanco to entirely different species, which in many cases do not extend to the Philippines.

Blanco's botanical material was, for most part, from the Island of Luzon, very few species being from the southern islands—Mindoro, Negros, Cebu, etc.—and not a single one from as far south as Mindanao. By far the greater part of his Luzon material came from provinces near Manila, especially from Rizal and Bulacan, while many specimens were from Pampanga, Bataan, Union, Ilocos Sur, and Camarines, few being from other provinces, such as Ilocos Norte, Cagayan, etc. In nearly all cases where Blanco does not cite the locality from which his material was received it is probable that it came from provinces near Manila, at least for the greater part. Very few were from the higher altitudes, and none from the higher mountains. Piwus taeda Blanco, non Linn., Llanosia toquian Blanco, and Blechnum colubrinum Blanco are examples of species from the mountains, but none of these are from the higher peaks.

The native names of plants are a very important factor in the identification of Blanco's species, but these are frequently very local and more or less subject to variation, both in pronunciation and in application. However, as a rule the native names recorded

by Blanco are to-day applied to the same species in the same locality, hence it becomes necessary, when collecting with the view to the rediscovery of those of Blanco, to secure, so far as possible, material in the very towns from which be received his specimens, for frequently a name in almost universal use in one locality to designate a certain species may be quite unknown to natives of a neighboring one, or if used by the latter may be applied to an entirely different Native names of species of economic value, those which yield valuable timbers, dves, and tans, food, fiber, medicines, etc., are frequently quite invariable and are universally used throughout the provinces inhabited by any one tribe of people to designate a certain definite species, and in many cases such plant names are found to be common to several dialects. In questioning natives in regard to the names and uses of plants one will frequently encounter people who have almost no knowledge even of the most common species, and unless the investigator has more or less experience in dealing with natives, and some knowledge of the subject of plant names, he is apt to receive a worthless list, made up for the occasion. However, in every town there will be found a certain number of persons who are thoroughly acquainted with the names and properties of the plants growing in the vicinity, and names secured from such persons are usually correct, at least for that locality, and can be depended upon. Certain names are used in a generic rather than in a specific sense to designate all species in a closely related group or species which have common characteristics, such as Balete, for most scandent species of Ficus; Malaruhat, for many species of Jambosa; Dapo, for most epiphytic plants, etc.

The present paper has been prepared in order to summarize in a convenient form our present knowledge of the species of plants described by Blanco and to call especial attention to such as are at present unknown, in order that collectors in the future may have some guide in collecting material which may serve to clear up the identity of some of his unknown species. In the preparation of this paper, wherever Blanco's names are found to be valid, and there is no doubt as to the identity of the species, they have been retained in accordance with the principles of priority. For the best interests of Philippine botany it is very essential that every effort be made to properly identify the large number of unknown species proposed by Blanco, as with so many in our flora a stable nomenclature can not be established until the majority are properly identified.

The arrangement of the present paper follows that of the Novissima Appendix. All the species described by Blanco are included with page references in the text under each to the first and second editions of the Flora de Filipinas, while in the general index the same are given to the third edition and to the Novissima Appendix. All the native names given by Blanco have been included. They are only in the following dialects: Tagalog, Pampangan, Visayan, Bicol, and Ilocano, represented respectively by the following letters: T., P., V., B., I. Such data as the date of flowering, locality, size of the plant, and habitat are given in the case of those species which to-day are doubtful or unknown, when such data are given by Blanco. Species of Blanco, of the correctness of the identification of which we are quite certain, are followed by an exclamation mark, while the unknown and uncertain ones are preceded by an asterisk.

ENUMERATION OF THE SPECIES.

RANUNCULACEÆ.

Atragene zeylanica Blanco, ed. 1, 461; ed. 2, 321, non Linn. Villar considers that Blanco includes two distinct species in his description and refers Blanco's species in part to Naravelia zeylanica DC., a species that does not extend to the Philippines, and in part to N. laurifolia Wall. Blanco's description seems to include the two species known as Naravelia laurifolia Wall., and Clematis gouriana Roxb., both these species being common and widely distributed in the Philippines. I., Banaibanai, Ovai na panglao.

DILLENIACEÆ.

- Dillenia indica Blanco, ed. 1, 472, non Linn.; Dillenia speciosa Blanco, ed. 2, 329, non Thunb. = Dillenia philippinensis Rolfe! T., Catmon.
- Delima aspera Blanco, ed. 1, 429; ed. 2, 299 = Tetracera sarmentosa (Linn.) Vahl.! (Delima sarmentosa Linn.) The pubescent form of this species. T., Malbas tagbalang.
- Tetracera monocarpa Blanco, ed. 1, 459; Tetracera sarmentosa (Linn.) Vahl., ed. 2, 320. The latter name is correct.! Erroneously referred by Villar to *Tetracera macrophylla* Wall., a species that does not extend to the Philippines. T., *Malacatmon*.

MAGNOLIACEÆ.

- Magnolia angatensis Blanco, ed. 1, 859; ed. 2, 328 = Talauma angatensis (Blanco) F. Vill.! T., Anobling.
- Michella champaca Linn., Blanco, ed. 1, 462; ed. 2, 322.! T., Sampac, Sampaca.
- * 'Kadsura blancoi Azaola in Blanco, ed. 2, 594. Reduced by Villar to Schizandra elongata Hook. f. et Th., a species that does not extend to the Philippines. Doubtless a true Kadsura and a good species, possibly represented by No. 23 Loher.

ANONACEÆ.

Unona setigera Blanco, ed. 1, 468; Uvaria setigera Blanco, ed. 2, 323. Reduced by Villar to *Uvaria purpurea* Blume, but is probably a synonym of *Uvaria rufa* Blume. Blanco's specimens were from Mandaloyon, near Manila.

¹Unknown species are preceded by an asterisk.

- * Unona camphorata Blanco, ed. 1, 486; ed. 2, 326. Reduced by Villar to *Uvaria dulcis* Dunal., a species at present unknown from the Philippines. Certainly, however, a species of *Uvaria.!* A tall scandent shrub in forests, the roots said to have the odor of camphor, bearing fruit in December. T., *Taghivalas*; V., *Dalaganum*, *Dalagao*.
- * Soala litoralis Blanco, ed. 1, 437; ed. 2, 304. Reduced by Villar to Cyathocalyx zeylanicus Champ., a species that does not extend to the Philippines. Certainly not a species of this genus.! Blanco's specimens were from Bauang, Province of Batangas; a weak shrub near the seashore, flowering in July. T., Soal.
- Uvaria sinensis Blanco, ed. 1, 465; Unona uncinata Blanco, ed. 2, 324, non DC. Reduced by Villar to Artabotrys odoratissimus R. Br., which is certainly correct.! T., Alangilang sonson.
- Unona corniculata Blanco, ed. 1, 469; ed. 2, 326. Reduced by Villar to Artabotrys suaveolens Blume, which is probably correct. Blanco's specimens were from Arayat, Province of Pampanga; the fruit said to be fragrant and edible. P., Susong damolag.
- Unona odoratissima Blanco, ed. 1, 467; ed. 2, 325 = Canangium odoratum (Lam.) Baill.! (Cananga odorata Hook. f. et Th.) T., Alangilang.
- Unona cabog Blanco, ed. 1, 466; Unona odorata? Blanco, ed. 2, 325.
 Reduced by Villar to Unona desmos Dunal., which is probably correct.
 T., Cabog.
- Unona ossea Blanco, ed. 1, 467; Uvaria ossea Blanco, ed. 2, 322. Reduced by Villar to *Unona discolor* Vahl., which is certainly correct.!
- * Macanea arborea Blanco, ed. 1, 431; Monodora myristica Blanco, ed. 2, 300. This species is transferred by Villar to the genus Monocarpia under the name Monocarpia blancoi F.-Vill. If this is the correct genus Blanco's first name should be taken up. A tree known in Cebu and Batangas, flowering in September. T., Palo calai, Calai, Caningag.
- Unona dehiscens Blanco, ed. 1, 466; ed. 2, 325 = Xylopia dehiscens (Blanco) Merrill, Forest. Bureau Bull. 1:20. 1903. (Xylopia blancoi Vidal.) This species was erroneously reduced by Villar to Anaxagorea luzonensis A. Gray.
- * Uvaria lanotan Blanco, ed. 1, 464; Unona latifolia? Blanco, ed. 2, 324. Reduced by Villar to Goniothalamus giganteus Hook. f. et Th., which can not be correct according to Blanco's description. Possibly a species of Mitrephora. A tree flowering in March. T., Lanotan.
- Uvaria tripetala Blanco, ed. 1, 465, non Roxb.; Unona tripetala Blanco, ed. 2, 324 = Phaeanthus cumingii Miq.! Erroneously reduced by Villar to Phaeanthus nutans Hook. f. et Th., a species that does not extend to the Philippines. A tree flowering in August. T., Lanotang itim.
- Anona squamosa Linn., Blanco, ed. 1, 469; ed. 2, 327.! T., Ates, Yates. Anona reticulata Linn., Blanco, ed. 1, 470; ed. 2, 327.! T., Anonas.

- Anona muricata Linn., Blanco, ed. 2, 326.! T., Guanabanos.
- *Uvaria amuyon Blanco, ed. 1, 463; Unona cauliflora Blanco, ed. 2, 323. Reduced by Villar to Melodorum fulgens Hook. f. et Th., but this is certainly an error, as Blanco's species is a tree, while Melodorum fulgens is a large woody climber. T., Amuyon, Grano del Parayso.

MENISPERMACEÆ.

- Menispermum rimosum Blanco, ed. 1, 810; ed. 2, 558 non Spreng. = Tinospora crispa Miers.! T., Macabuhay; V., Pangiavan, Panavan, Taganagtagua, Paliavan.
- Menispermum cocculus Linn., Blanco, ed. 1, 809; ed. 2, 557 = Anamirta cocculus (Linn.) W. et A.! T., B., P., Lactang, Lictang, Suma, Lanta, Lintang baguin, Tuba, Balasin, Bayati.
- Cissampelos pareira Linn., Blanco, ed. 1, 815; ed. 2, 563.! T., Sansao, Sansaosansauan, Calacalamayan, Batangbatang, Cuscusipa, Calaad.

NYMPHAEACEÆ.

- Nymphaea lotus Linn., Blanco, ed. 1, 456; ed. 2, 317.! T., B., P., Lavas, Talailo, Gavai gabai, Tunas.
- Nymphaea turbinatum Blanco, ed. 1, 458; Nelumbium speciosum Willd., Blanco, ed. 2, 318. The latter name is correct.! T., Baino; I., Sucao.

PAPAVERACEÆ.

Argemone mexicana Linn., Blanco, ed. 1, 454; ed. 2, 316.! I., Casubhang aso.

CRUCIFEREÆ.

- Cardamine glandulosa Blanco, ed. 1, 521; Cardamine impatiens?

 Blanco, ed. 2, 363. Referred by Villar to Nasturtium indicum DC.,
 which is undoubtedly correct. Blanco's specimens were from Pasig,
 Province of Rizal, flowering in January.
- Brassica orientalis Blanco, ed. 1, 519; ed. 2, 361 non Linn. Referred by Villar to *Brassica nigra* Koch., which is probably correct. Blanco's specimens were from Pasig, Province of Rizal, flowering in March.
- Sinapis brassicata Linn., Blanco, ed. 2, 362. Referred by Villar, and probably correctly so, to *Brassica campestris* Linn. Cultivated.
- Sinapis sinensis Blanco, ed. 1, 520; Sinapis juncea Linn., Blanco, ed. 2, 362. Referred by Villar to Brassica juncea Hook. f. et Th., which is probably correct. Sp.-Fil., Mostaza.

CAPPARIDACEÆ.

Cleome viscosa Linn., Blanco, ed. 1, 522; Polanisia viscosa DC., Blanco, ed. 2, 364. The former name is correct.! T., Balabalanoyan, Apoiapoyan.

21896 - - 2

- Cleome alliacea Blanco, ed. 1, 522; Cleome alliodora Blanco, ed. 2, 363 = Pedicellaria pentaphylla (Linn.) Schrank.! (Gynandropsis pentaphylla DC.) T., Balabalanoyan, Apoiapoyan.
- *Cleome pentaphylla Blanco, ed. 1, 523; Cleome gigantea Blanco, ed. 2, 364. Reduced by Villar to the American species, Cleome speciosa DC. Blanco states that his species was known in Manila under the name of Araña. Probably a cultivated species, but Blanco does not state this fact.
- Crataeva religiosa Forst., Blanco, ed. 1, 399; ed. 2, 279.! I., Balai namoc.
- *Capparis baducca Blanco, ed. 1, 438 non Linn.; Capparis mariana Jacq., Blanco, ed. 2, 305. The latter name is possibly correct, Blanco's specimens being from Parañaque, near Manila, said to have been introduced from the Marianne Islands. T., Alcaparras.
- Crataeva octandra Blanco, ed. 1, 400; d. 2, 380. Reduced by Villar to Capparis floribunda Wall., which may possibly be correct. Blanco's specimens were from Piddig, Province of Ilocos Norte.
- Capparis nemorosa Blanco, ed. 1, 439; C. micracantha Blanco, ed. 2, 305, non DC. Reduced by Villar to Capparis horrida Linn. f., which is probably correct. T., Dauag. I., Talactac.
- *Capparis linearis Blanco, ed. 1, 438; ed. 2, 305. Reduced by Villar to Capparis viminea Hook. f. et Th., which is certainly an error; a shrub said by Blanco to be common in forests.
- *Capparis odorata Blanco, ed. 1, 439; ed. 2, 305. A species known only from Blanco's description, and one that Villar makes no attempt to reduce. Blanco's specimens were from Angat, Province of Bulacan.

BIXACEÆ.

- Bixa orellana Linn., Blanco, ed. 1, 456; ed. 2, 317.! T., Achote.
- Banara racemosa Blanco, ed. 1, 425; Flacourtia corollata Blanco, ed. 2, 559 = Scolopia crenata Clos.! Erroneously reduced by Villar to Scolopia rhinanthera Clos., a species not known from the Philippines. T., Pilapil.
- * Banara brevifolia Blanco, ed. 1, 426; Flacourtia parviflora Blanco, ed. 2, 560. Reduced by Villar to Scolopia dasyanthera Benn., a species not known from the Philippines. Possibly identical with Flacourtia ramontchi L'Herit.
- Miroxylum decline Blanco, ed. 1, 813; Stigmarota edulis Blanco, ed. 2, 560 = Flacourtia sepiaria Clos.! T., Bitonyol.
- Hydnocarpus polyandra Blanco, ed. 2, 545 = Pangium edule Reinw.! V., Pangui.

PITTOSPORACEÆ.

Bursaria inermis Blanco, ed. 2, 124. Possibly identical with Pittosporum ferrugineum Ait.

Aquilaria pentandra Blanco, ed. 1, 373; Limonia laureola Blanco, ed. 2, 251, non DC. = PITTOSPORUM PENTANDRUM (Blanco) (Pittosporum fernandezii, Vidal, in F.-Vill. Nov. App. 13. 1880; Sinopsis, Atlas, t. 8. f. A. 1883; Pittosporum brachysepalum Turcz.! 1854.) Blanco's name is the earliest for this species which is represented by Nos. 500, 1308, 1438, 2804, 2951, and 3177 Merrill. T., Mamalis.

POLYGALACEÆ.

- Polygala monspeliaca Blanco, ed. 1, 557; ed. 2, 388, non Linn. Reduced by Villar to *Polygala telephioides* Willd., which is probably correct. Blanco's specimens were from Mandoloyon, near Manila.
- Amorpha pedalls Blanco, ed. 1, 553; ed. 2, 387. Reduced by Villar to Salomonia oblongifolia DC., which is probably correct. Blanco's specimens were from Malinta, near Manila, flowering in October.
- Securidaca volubilis Blanco, ed. 1, 556, non Linn.; S.? complicata Blanco, ed. 2, 388, non H. B. K. Reduced by Villar to Securidaca tavoyana Wall., which is certainly an error, as this species does not extend to the Philippines. Probably identical with Securidaca corymbosa Turcz. The bark is used as a substitute for soap. T., B., P., Gogong bisaya, Gogong bacay, Balago.

CARYOPHYLLACEÆ.

Polycarpon polyphyllum Blanco, ed. 1, 53; ed. 2, 36. Reduced by Villar to *Polycarpon loeflingiae* Benth. et Hook. f., which is probably correct. Blanco's specimens were from Pasig, Province of Rizal, flowering in February.

PORTULACACEÆ.

- Portulaca oleracea Linn., Blanco, ed. 1, 407; ed. 2, 284.! T., Olasiman, Colasiman; B., Ausiman.
- Portulaca meridiana Linn. f., Blanco, ed. 1, 408; ed. 2, 285 = Portulaca quadrifida Linn.! T., Sayican.

ELATINEACÆ.

- Tillaea rubella Blanco, ed. 1, 75; ed. 2, 56. Reduced by Villar to Bergia ammannioides Roxb., which is probably correct. Blanco's specimens were from Pasig, Province of Rizal.
- Bergia serrata Blanco, ed. 1, 387; Spergula serrata Blanco, ed. 2, 271. Reduced by Villar to Bergia verticillata Willd., which is probably correct.

HYPERICACEÆ.

Hypericum olympicum Blanco, ed. 1, 613; ed. 2, 429, non Linn. = Cratoxylon blancoi Blume.! Erroneously reduced by Villar to Cratoxylon polyanthum Korth. T., Guyong guyong.

Hypericum aegyptium Blanco, ed. 1, 615; ed. 2, 430, non Linn. Reduced by Villar to *Cratoxylon formosum* Dyer, which is probably correct. Blanco's specimens were from Angat, Province of Bulacan, a tree flowering in August.

GUTTIFEREÆ.

- Cambogia binucao Blanco, ed. 1, 434; ed. 2, 302 = Garcinia binucao (Blanco) Choisy.! Erroneously reduced by Villar to Garcinia cornea Desrouss. T., Binucao; V., Bagobago.
- Cambogia venulosa Blanco, ed. 1, 435; ed. 2, 302 = Garcinia venulosa (Blanco) Choisy.! Erroneously reduced by Villar to Garcinia cornea Linn., a species that does not extend to the Philippines. A tree flowering in August. T., Taclang anac.
- * Stalagmites? cambogloides Blanco, ed. 2, 301, non Murr. Certainly a species of *Garcinia.!* This species is not mentioned by Villar in the Novissima Appendix. Blanco's specimens were from Bauang, Province of Batangas, a tree flowering in August. T., Banago.
- * Cambogla crassifolia Blanco, ed. 2, 304. Reduced by Villar to Garoinia cowa Roxb., which is undoubtedly an error. Blanco's specimens were from Cebu. V., Sagudan.
- *TovomIta pentapetala Blanco, ed. 1, 432; ed. 2, 301. This species was transferred by Villar to Ochrocarpus, as a distinct species, Ochrocarpus pentapetalus (Blanco) F.-Vill. It is, however, known only from Blanco' description. A tree flowering in December, growing along the seashore in the Provinces of Ilocos Norte and Ilocos Sur. I., Darigayos, Pamitlain, Pamitlatin.
- *Willughbeia multilocularis Blanco, ed. 1, 131; ed. 2, 94. This species was first referred by Villar to the Apocinaceæ, but in the Addenda et Corrigenda he refers it to Garcinia ovalifolia var. spicata Hook. f. et Th. Possibly correct as to the genus, but certainly not this species. A tree flowering in August. T., Malabatoan.
- Calophyllum inophyllum Linn., Blanco, ed. 1, 612; ed. 2, 428.! T., Palo maria, Bitanhol, Tamauian, Dancalan, Dincalan; B., Dincalan, Bitaog; P., Bitaog, Palo maria; I., Bitaog.
- * Calophyllum apetalum Blanco, ed. 2, 429, non Willd. Reduced by Villar to Calophyllum pulcherrimum Wall., a species that does not extend to the Philippines. Possibly identical with Calophyllum spectabile Willd. Blanco's specimens were from Angat, Province of Bulacan. T., Malabochoc.
- Plinia paniculata Blanco, ed. 1, 423; ed. 2, 296 = Kayea paniculata (Blanco) Merrill, Govt. Lab. Publ. 17:29. 1904. Erroneously reduced by Villar to Kayea racemosa Pl. et Tr., a species that does not extend to the Philippines. T., Guisan.

TERNSTROEMIACEÆ.

- Guettarda vermicularis Blanco, ed. 1, 723; ed. 2, 500. Reduced by Villar to Anneslea fragrans Wall., a species that does not extend to the Philippines. Probably identical with Anneslea crassipes Hook. Blanco's specimens were from the seashore, Bauang, Province of Batangas; a shrub, flowering in October. T., Malatibig.
- Lianosia toquian Blanco, ed. 2, 319 = TAONABO TOQUIAN (Blanco) (Ternstroemia toquian F.-Vill.!) This species is certainly represented by Nos. 2288 and 3206 Merrill; 150, 137, and 208 Forestry Bureau, collected by Barnes, and No. 57 Ahern, Blanco's description in some respects being erroneous. Gilg, in Engler und Prantl, Natürlichen Pflanzenfamilien retains Llanosia as an imperfectly known genus of the Dilleniaceæ. T., Toquian.
- *Camellia sasanqua Blanco, ed. 2, 371 non Thunb. This species is reduced by Villar to Camellia drupifera Lour., but this identification is very doubtful. Blanco's specimens were from a plant cultivated in Manila.
- Salceda montana Blanco, ed. 2, 374 = Thea Montana (Blanco) (Camellia lanceolata F.-Vill., non Seem.). Nos. 304, 441 Whitford; Nos. 1905, 2778, 2402 Forestry Bureau. Blanco's specimens were from Angat, Province of Bulacan.
- * Helianthemum triflorum Blanco, ed. 2, 309. Reduced by Villar to Archytaea vahlii Choisy, which is undoubtedly an error. Blanco's specimens were from Angat, Province of Bulacan; flowering in February. T., Iring.

DIPTEROCARPACEÆ.

- Mocanera mayapis Blanco, ed. 1, 449; Dipterocarpus mayapis Blanco, ed. 2, 313 = Anisoptera vidaliana Brandis?; Dipterocarpus turbinatus F.-Vill., non Gaertn.! Blanco's species is probably identical with Anisoptera vidaliana Brandis, although Blanco describes the petioles as short. This is apparently one of the most common species of Dipterocarpacea in Luzon, and is universally known to the Tagalogs as Mayapis, the name cited by Blanco. Blanco's specimens were from the mountains near Angat, Province of Bulacan. T., Mayapis.
- Mocanera vernicifiuua Blanco, ed. 1, 540; Dipterocarpus vernicifiuus Blanco, ed. 2, 314. The latter name is the correct one, Dipterocarpus velutinus Vidal being certainly a synonym. Erroneously referred by Villar to Dipterocarpus hispidus Thwaites. T., Panao, Balao, Malapao.
- Mocanera grandifiora Blanco, ed. 1, 451; Dipterocarpus grandifiorus Blanco, ed. 2, 314.! The latter name is correct. Erroneously referred by Villar to Dipterocarpus griffithii Miq., a species that does not extend to the Philippines. T., Apiton, Hapiton.

- Mocanera thurifera Blanco, ed. 1, 446; Dipterocarpus thurifer Blanco, ed. 2, 310 = Anisoptera thurifera (Blanco) Blume.! T., Sandana, Lauan; B., Lauaan.
- * Vatica sinensis Blanco, ed. 1, 401; ed. 2, 280, non Gmel. Reduced by Villar to Vatica grandiflora Dyer, but this species does not extend to the Philippines. Said by Blanco to be closely related to Vatica mangachapoi, and probably only a form of that species. Blanco's specimens were from San Mateo, Province of Rizal.
- Vatica mangachapoi Blanco, ed. 1, 401; Vatica apteranthera Blanco, ed. 2, 281. Erroneously reduced by Villar to Vatica scaphula Dyer, a species that does not extend to the Philippines. The first name is the correct one. T., Mangachapoi.
- Mocanera polysperma Blanco, ed. 1, 448; Dipterocarpus polyspermus Blanco, ed. 2, 312 = Shorea polysperma (Blanco). Erroneously referred by Villar to Shorea talura Roxb., a species that does not extend to the Philippines. This distinct species is represented by Nos. 606, 734, 787, and 819 Forestry Bureau. T., Tangili.
- * Dipterocarpus palosapis Blanco, ed. 2, 312. Erroneously referred by Villar to Shorea floribunda Kurz, a species that does not extend to the Philippines. Probably a Shorea, but known only from Blanco's description. T., Palosapis.
- Mocanera malaanonan Blanco, ed. 1, 858; Dipterocarpus malaanonan Blanco, ed. 2, 312 = Shorea malaanonan (Blanco) Blume. Probably a distinct species, but not well-known. Blanco's specimens were from Angat, Province of Bulacan. T., Malaanonan.
- *Euphoria malaanonan Blanco, ed. 1, 289; Euphoria? nephelium? Blanco, ed. 2, 200. Reduced by Villar to Shorea robusta Gaertn. f., which is certainly an error, at least as to the species, possibly identical with the preceding species. Blanco's specimens were from Angat, Province of Bulacan, his material being very imperfect. T., Malaanonan.
- Mocanera guiso Blanco, ed. 1, 449; Dipterocarpus guiso Blanco, ed. 2, 313 = Shorea guiso (Blanco) Blume.! Erroneously reduced by Villar to Shorea robusta Gaertn. f., a species that does not extend to the Philippines. T., Guiso, Guijo.
- Mocanera plagata Blanco, ed. 1, 447; Dipterocarpus plagatus Blanco, ed. 2, 311 = Hopea plagata (Blanco) Vidal.! Erroneously referred by Villar to Shorea reticulata Thwaites, an extra-Philippine species. T., Yacal, Sapolongan.
- * Mocanera mangachapoi Blanco, ed. 1, 450; Dipterocarpus mangachapoi Blanco, ed. 2, 313 = Shorea mangachapoi (Blanco) Blume.! This species is known only from Blanco's description, and should not be confused with Vatica mangachapoi Blanco. Blanco's specimens were from Angat, Province of Bulacan, and from San Mateo, Province of Rizal. T., Guisong dilao, Mangachapoi.

MALVACEÆ.

- Althaea sinensis Blanco, ed. 1, 552, non Cav.; A. rosea Cav., Blanco, ed. 2, 386.! The latter name is correct.
- Malva coromandelina Blanco, ed. 2, 551, non Cav.; Malva luzonica Blanco, ed. 2, 385 = Malvastrum tricuspidatum A. Gray.! T., Quinalumpang.
- Sida lanceolata Roxb., Blanco, ed. 1, 548;
 Sida frutescens Blanco, ed. 2, 384, non Cav. = Sida carpinifolia Linn.!
 T., V., P., Ualualisan, Escobang haba, Pamalis, Higot balato, Mamalis.
 I., Taquing baca.
- Sida truncatula Blanco, ed. 1, 548; Sida philippica DC., Blanco, ed. 2, 283 = Sida rhombifolia Linn. var.! T., Escobang bilog.
- Sida semicrenata Link., Blanco., ed. 2, 384 = Sida rhombifolia Linn. var.!
- Sida indica Linn., Blanco. ed. 1, 547; ed. 2, 283 = Abutilon indicum (Linn.) G. Don.! T., Cuacuacohan, Guling gulingan, Tabing, Malis, Dalupang, Palis, Malvas de Castilla.
- Napea latifolia Blanco, ed. 2, 387 = Malachra capitata Linn.!
- Napea scabra Blanco, ed. 1, 553; ed. 2, 386, non Linn. This is Malachra lineariloba Turcz.!, erroneously reduced by Villar to Malachra bracteata Cav., an American species. T., Lapnis na bolohan, Anabo. V., Vocabul.
- Urena multifida Blanco, ed. 1, 540; ed. 2, 378, non Cav. = Urena lobata Linn.! T., V., P., Colotan, Colotcolotan, Molopolo, Dalupan.
- * Hibiscus batacensis Blanco, ed. 1, 544; ed. 2, 380. Reduced by Villar to Kosteletzyka, as a distinct species K. batacensis (Blanco) F.-Vill. A species known only from Blanco's description, whose specimens were from the town of Batac, Province of Ilocos Norte.
- Hibiscus bifurcatus Blanco, ed. 1, 545; ed. 2, 380 = Hibiscus surratensis Linn.!
- Hibiscus abelmoschus Linn., Blanco, ed. 1, 545; ed. 2, 380.! T., Castoli, Castiocastiogan, Dalupan. V., Maricum Ducum, Marucum, Maropoto, Sapinit. P., Castiocastiolan.
- Hibiscus tiliaceus Linn., Blanco, ed. 1, 541; ed. 2, 379.! T., P., Balibago. V., Malibago, Raguindi.
- Hibiscus rosa-sinensis Linn., Blanco, ed. 1, 543; ed. 2, 379.! T., V., P., Tacurangan, Arogangan, Antolangan, Cayanga, Tapolanga, Tarocanga, Gomamila. I., Cayanga rosa.
- Hibiscus mutabilis Linn., Blanco, ed. 1, 546; ed. 2, 381.!
- Thespesia sublobata Blanco, ed. 2, 382 = Thespesia lampas D. et G.!
- Hibiscus populneus Linn., Blanco, ed. 1, 544; Thespesia populnea (Linn.) Corr., Blanco, ed. 2, 381. The latter name is correct.! T., Boboi gubat.
- Thespesia banalo Blanco, ed. 2, 382 = Thespesia macrophylla Blume.! T., Banalo, P., Banaro, V., Banago.

- Gossypium herbaceum Linn., Blanco, ed. 1, 534; ed. 2, 374; G. religiosum L., Blanco, ed. 2, 374. T., Bulac.
- Gossypium paniculatum Blanco, ed. 1, 539; ed. 2, 378 = Gossypium barbadense Linn. var.! I., Capas.
- Gossypium perenne Blanco, ed. 1, 537; ed. 2, 376 = Gossypium arboreum Linn.! T., Bulac na monti, Bulac na totoo, Bulac cahoy.
- Melaleuca grandifiora Blanco, ed. 1, 615; Bombax ceiba Linn., Blanco, ed. 2, 372. The latter name is correct.! T., Taglinao, Malabulac.
- Bombax pentandrum Linn., Blanco, ed. 1, 531; ed. 2, 371 = Ceiba pentandra (Linn.) Gaertn.! (Eriodendron anfractuosum DC.) T., Boboi, V., Doldol, P., Bulac castila.

STERCULIACEÆ.

- Sterculia foetida Linn., Blanco, ed. 1, 763; ed. 2, 524.! T., Calumpang, I., Bangar.
- Sterculia cordifolia Blanco, ed. 1, 764; ed. 2, 525, non Cav. = Sterculia philippinensis Merrill, Govt. Lab. Publ. 17: 29. 1904. Erroneously reduced by Villar to Sterculia urens Roxb., a species that does not extend to the Philippines. T., Banilad, Banilar.
- Sterculia balanghas Blanco, ed. 1, 765; ed. 2, 525, non Linn. Reduced by Villar to Sterculia rubiginosa Vent., a species that does not extend to the Philippines. Apparently the same as Sterculia cuneata R. Br. T., Nato, Malabunot.
- Sterculia alata Blanco, ed. 2, 525, non Roxb. = Sterculia blancoi Rolfe.! Villar erroneously considered that Blanco's species was identical with that of Roxburgh.
- Heritiera tinctoria Blanco, ed. 1, 653; ed. 2, 456 = Pterocymbium tinctorium (Blanco) (Pterocymbium javanicum R. Br.; Sterculia campanulata Wall.) Blanco's name is the earliest one published for this species and should be retained. T., Taloto, V., Duha?, Busain.
- *Sterculia glandulosa Blanco, ed. 1, 764. This species is known only from Blanco's short and imperfect description, and is excluded from the second edition. No locality or native name is given.
- *Sterculia glomerata Blanco, ed. 1, 764; ed. 2, 525. This species is quite unknown, but is probably a species of *Myristica* as suggested by Villar. A tree having a red gum. T., *Hindurugu*.
- *Sterculia decandra Blanco, ed. 1, 766; ed. 2, 526. This species is probably also a *Myristica* as suggested by Villar. Blanco's specimens were from Balanga, Province of Bataan, a large tree, the wood having a foetid odor. T., *Banabanalo*.
- Helicteres apetala Blanco, ed. 1, 383; Sterculia cymbiformis Blanco, ed. 2, 526 = Heritiera littoralis Dry.! T., Dongon, I., Paronapin.
- Kleinhovia hospita Linn., Blanco, ed. 1, 652; ed. 2, 455.! T., B., Tanag, B., Hamitanag, P., Panampat, I., Bitnong.

- Dombeya decandra Blanco, ed. 1, 349; Dombeya biserrata Blanco, ed. 2, 244 = Helicteres spicata Coleb.! T., Dangling aso, Buntot usa, I., Baquinbaquit.
- Mimusops talosan Blanco, ed. 1, 284; ed. 2, 198. Reduced by Villar to a variety (lanigera) of the preceding and certainly only a form of the species.! Blanco's specimens were from Malinta, near Manila, a shrub flowering in September. T., Talosan.
- Pterospermum hastatum Blanco, ed. 1, 526; ed. 2, 367 = Pterospermum diversifolium Blume.! T., Bayog.
- Pterospermum obliquum Blanco, ed. 1, 529; Pterospermum semisagittatum Blanco, ed. 2, 368, non Buch.-Ham. A good species, the first name being the correct one.! Erroneously reduced by Villar to Pterospermum rubiginosum Hiern, a species that does not extend to the Philippines. T., Calocatingan.
- Pentapetes coccinea Blanco, ed. 1, 529; Pentapetes cebuana Blanco, ed. 2, 369 = Pentapetes phoenicea Linn.!
- Melochia supina Linn., Blanco, ed. 1, 524; ed. 2, 365 = Melochia corchorifolia Linn.!
- Geruma subtrilobata Blanco, ed. 1, 182; ed. 2, 130 = the preceding species!
- Melochia arborea Blanco, ed. 1, 524; ed. 2, 130 = Melochia indica (Houtt.) A. Gray.! (Melochia velutina Bedd.)
- Hypericum pentandrum Blanco, ed. 1, 616; ed. 2, 430 = the preceding species!
- Waltheria americana Linn., Blanco, ed. 1, 523; ed. 2, 364.! (Waltheria indica Linn.)
- Abroma communis Blanco, ed. 1, 606; ed. 2, 160 = Abroma augusta Linn.!
- Abroma alata Blanco, ed. 1, 605; ed. 2, 422 = the preceding species!

 This species is retained by Villar as distinct, but is certainly only a form of Abroma augusta Linn.! T., Anibong, Anibiong, V., Nabo, Labon, Anabon.
- Theobroma cacao Linn., Blanco, ed. 1, 601; ed. 2, 419!
- Commersonia echinata Blanco, ed. 2, 160 = Commersonia platyphylla Andr.!

TILIACEÆ.

*Triopteris poliandra Blanco, ed. 1, 380; ed. 2, 268. Reduced by Villar to Berrya ammonilla Roxb., which is certainly an error, as Blanco's description does not apply to that species. Blanco's specimens were from Angat, Province of Bulacan, and from Malinta, near Manila, a shrub, flowering in July. T., Pacpacbalang.

- *Grewia serrata Blanco, ed. 1, 444; Columbia serratifolia Blanco, ed. 2, 427, non DC. Reduced by Villar to Grewia columnaris Sm., a species not known from the Philippines. Possibly the same as Grewia multiflora Juss. Blanco's specimens were from the Tala Mountains, Province of Rizal or Bulacan.
- Mallococca crenata Blanco, ed. 1, 442; Grewia multiflora? Blanco, ed. 2, 309. Reduced by Villar to *Grewia umbellata* Roxb., which is certainly correct.
- Mallococca parva Blanco, ed. 1, 443; Grewia mallococca, Blanco, ed. 2, 310 = Grewia multiflora Juss.! T., Danglin.
- Columbia anilao Blanco, ed. 1, 554; ed. 2, 426 = Colona serratifolia Cav.! Erroneously retained by Villar as a distinct species. T., Anilao.
- Triumfetta semitriloba Linn., Blanco, ed. 1, 406; ed. 2, 283.! T., Colotan bilog.
- Corchorus capsularis Linn., Blanco., ed. 1, 442; ed. 2, 308.! T., Pasao na bilog.
- Corchorus catharticus Blanco, ed. 1, 442; Corchorus olitorius Linn., Blanco, ed. 2, 308. The latter name is correct.! T., Salayong.
- Corchorus aestuans Blanco, ed. 1, 441; Corchorus acutangulus Linn., Blanco, ed. 2, 308. The latter name is correct.! T., Pasao na haba.
- * Vallea calomala Blanco, ed. 1, 439; Ageratium calomala Blanco, ed. 2, 306. Reduced by Villar to Elaeocarpus lancaefolius Roxb., a species that does not extend to the Philippines, certainly, however, correct as to the genus. Blanco's specimens were from the Province of Batangas, flowering in August. T., Calomala.
- * Elaeocarpus sylvestris Blanco, ed. 2, 306. This species is reduced by Villar to *Elaeocarpus oblongus* Gaertn., but Blanco's description scarcely applies to this species. Blanco's specimens were from Cebu. V., Cabilte, Cabalte.

MALPHIGIACEÆ.

- Hiraea reclinata Blanco, ed. 1, 378, non Jacq.; Tristillateia malintana Blanco, ed. 2, 267 = Tristillateia australasiaca A. Rich.!
- Triopteris jamaicensis Blanco, ed. 1, 379; ed. 2, 207, non Linn. = Hiptage madablota Gaertn.! V., Comimpol.

ZYGOPHYLLACEÆ.

Tribulus lanuginosus Blanco, ed. 1, 350; ed. 2, 245, non Linn. = Tribulus cistoides Linn.!

GERANIACEÆ.

Malva moschata Blanco, ed. 1, 551; ed. 2, 385, non Linn. A cultivated species, reduced by Villar to *Pelargonium odoratissimum* Ait., which is probably correct. Sp. Fil., *Malva rosa*.

- Oxalis acetosella Blanco, ed. 1, 388; ed. 2, 272, non Linn. = Oxalis corniculata Linn.! T., Tangain daga, Susocoyoli. B., Daraisig. P., Malabalugbug daguis, Ayo, Cungi, Yayo.
- Oxalis sensitiva Linn., Blanco, ed. 1, 399; Biophytum sensitivum DC., Blanco, ed. 2, 272. The last name is correct. T., Macahiya, Damonghiya, I., Mahahiin.
- Averrhoa carambola Linn., Blanco, ed. 1, 391; ed. 2, 274.! T., Balimbin, Bilimbin.
- Averrhoa bilimbi Linn., Blanco, ed. 1, 392; ed. 2, 273.! T., Calamias, Camias, B., Quilingiva, I., Pias.
- *Averrhoa pentandra Blanco, ed. 1, 392; ed. 2, 274. Reduced by Villar to Connaropsis as a distinct species, Connaropsis philippica F.-Vill. If correctly referred, Blanco's specific name should be retained. Blanco's specimens were from Malinta, near Manila, flowering in October.
- Impatiens triflora Blanco, ed. 1, 636; ed. 2, 443, non Willd. = Impatiens balsamina Linn.! T., P., Camantigui. V., Suranga, Solanga.

RUTACEÆ.

- Cissus frutescens Blanco, ed. 1, 70; Cissus arborea Blanco, ed. 2, 51.

 Reduced by Villar to Evodia roxburghiana Benth., which is probably correct. Blanco's specimens were from the Province of Batangas, and from Malinta, near Manila. T., Matang diablo.
- Melicope tetrandra Blanco, ed. 1, 293, non Roxb.; Evodia triphylla DC., Blanco, ed. 2, 50. The last name is correct.
- Evodia bintoco Blanco, ed. 2, 50. Referred by Villar to Evodia latifolia DC., which is probably correct. Blanco's specimens were from Bohol and Samar, flowering in October. V., Bintoco, Tancapan.
- *Orixa ternata Blanco, ed. 1, 62; ed. 2, 45. Reduced by Villar to Evodia robusta Hook. f., a species that is unknown from the Philippines, probably, however, correct as to the genus. A shrub flowering in July.
- * Fagara octandra Blanco, ed. 1, 67; ed. 2, 47. Reduced by Villar to Melicope ternata Forst., a species that certainly does not extend to the Philippines. Blanco's specimens were from San Mateo, Province of Rizal, flowering in June. T., Matang arao.
- Fagara piperita Blanco, ed. 1, 64; ed. 2, 47, non Linn. Reduced by Villar to Zanthoxylum oxyphyllum Edgw., a species that certainly does not extend to the Philippines. Correct as to the genus and probably identical with Zanthoxylum piperatum DC. T., Cayutana.
- Fagara pterota Blanco, ed. 1, 66; ed. 2, 47, non Linn. Referred by Villar to Zanthowylum avicennæ DC., which is probably correct. P., T., Cayutana, V., Salay marbar, Baga tambal, P., Salay cangay.
- Lunasia amara Blanco, ed. 1, 783; Pilocarpus amara Blanco, ed. 2, 540. The first name is correct, the type of the genus *Lunasia*, one of the few valid genera established by Blanco. T., *Lunas*.

- * Bergera compressa Blanco, ed. 1, 360; ed. 2, 254. Reduced by Villar to Clausena willdenovii W. et A., a species that does not extend to the Philippines. Probably, however, correct as to the genus. A small tree flowering in August. T., Piris.
- * Bergera inodora Blanco, ed. 1, 361; Bergera koenigii Blanco, ed. 2, 254, non Linn. Reduced by Villar to *Clausena indica* Oliv., a species that does not extend to the Philippines, probably, however, correct as to the genus. Blanco's specimens were from Malinta, near Manila.
- Limonia linearis Blanco, ed. 1, 357; Limonia monophylla Linn., Blanco, ed. 2, 252 = Atalantia monophylla (Linn.) Correa. ? T., Dayap na monti.
- Limonia disticha Blanco, ed. 1, 356; Limonia corymbosa Blanco, ed. 2, 251 = Atalantia disticha (Blanco) (Atalantia nitida Oliv., Journ. Linn. Soc. 5, Suppl. 2:25. 1861.) Blanco's name is the oldest for this species and should be retained. Erroneously referred by Villar to Atalantia racemosa W. et A., a species that does not extend to the Philippines. T., Malarayap.
- Limonia trifoliata Linn., Blanco, ed. 1, 357; ed. 2, 252 = Triphasia trifoliata (Linn.) DC.! T., Limoncitos.
- Melicope conferta Blanco, ed. 2, 205. Reduced by Villar to Acronychia laurifolia Blume, which is probably correct.
- Murraya cerassiformis Blanco, ed. 1, 363; Murraya exotica Blanco, ed. 2, 255, non Linn. = Glycosmis pentaphylla Correa.! T., Guinguen.
- Murraya lobata Blanco, ed. 1, 363; ed. 2, 256. Reduced by Villar to a variety of *Glycosmis pentaphylla* Correa, which is certainly correct.
- *Bergera ternata Blanco, ed. 1, 360; ed. 2, 254. Reduced by Villar to Glycosmis bilocularis Thwaites, a species that does not extend to the Philippines. Probably correct as to the genus, and a form of Glycosmis pentaphylla Correa. Blanco's specimens were from Tala, Province of Rizal or Bulacan, a shrub with fragrant fruits. T., Malamolauin.
- Andromeda japonica Blanco, ed. 1, 371; ed. 2, 261, non Thunb. = Micromelum tephrocarpum Turcz.! Erroneously referred by Villar to Micromelum glabrescens Benth., a species that does not extend to the Philippines. T., Tulibas.
- Connarus foetens Blanco, ed. 1, 525;
 Connarus santaloides Blanco, ed.
 2, 366 = Murraya exotica Linn.! T., Camuning, Molavin. V., P., Camunin, Banati.
- Cookia anisum olens Blanco, ed. 1, 359; Cookia anisodora Blanco, ed.
 2, 253 = Clausena anisum olens (Blanco) Merrill, Govt. Lab. 17: 21.
 1904. T., Maisipaisi, Calomata, Camanguianis.
- Cookia wampi Blanco, ed. 1, 358; ed. 2, 253 = Clausena wampi (Blanco) Oliver.! T., Huampit.
- Citrus notissima Blanco, ed. 607; ed. 2, 424. Reduced by Villar to Citrus medica Linn., var. acida Hook. f., which may be correct. T., Dayap. Sp.-Fil., Limonoito.

- * Citrus mitis Blanco, ed. 1, 610; ed. 2, 426. Reduced by Villar to Citrus medica Linn., var. limetta Hook. f., but Blanco's description does not apply to this variety. A small spineless shrub. T., Aldonises, Calamondin.
- Citrus reticulata Blanco, ed. 1, 610; ed. 2, 424 = Citrus aurantium Linn.! T., Sintonis. Sp.-Fil., Narangitas.
- Citrus aurantium Blanco, ed. 1, 609; ed. 2, 425, non Linn. Reduced by Villar to Citrus aurantium Linn., var. bigrardia Hook. f., but Blanco's description does not agree with that of this variety. A form or variety of Citrus aurantium Linn.! T., Dalandan, Cahil.
- Citrus papillaris Blanco, ed. 1, 610; ed. 2, 425. Reduced by Villar to Citrus aurantium Linn., var. bergamina Hook. f., which is certainly an error. Certainly a form or variety of Citrus aurantium Linn.! T., Pisong.
- Citrus torosa Blanco, ed. 1, 609; ed. 2, 425. Reduced by Villar to Citrus hystrix DC., which is probably correct. T., Suya, Colobot.
- Citrus decumana Linn., Blanco, ed. 1, 606; ed. 2, 424.! T., Lucban, Suha.
- Murraya odorata Blanco, ed. 2, 256. Reduced by Villar to Feronia elephantum Correa, which is probably correct. Blanco's specimens were from Angat, Province of Bulacan, flowering in February.
- Limonia glutinosa Blanco, ed. 1, 358; Feronia ternata Blanco, ed. 2, $252 = Aegle\ glutinosa\ (Blanco)\ Merrill, Govt. Lab. 6: 12. 1904. (Aegle decandra Naves.) T., Malacabuyao. P., T., Tabog.$

SIMARUBACEÆ.

- Manungala pendula Blanco, ed. 1, 306; Niota tetrapetala Lam., Blanco, ed. 2, 213 = Samadera indica Gaertn.! T., V., P., B., Manungal V., Linatong anac, Linton gamai, Palagium, Palagarium, Ponoan, Mavindato, Daraput.
- Paliurus perforatus Blanco, ed. 1, 174; ed. 2, 122 = Harrisonia brownii A. Juss.! T., Asimao.
- Paliurus dubius Blanco, ed. 1, 175; ed. 2, 122. Reduced by Villar to *Harrisonia bennetii* Hook. f., var. multijuga F.-Vill. Probably identical with H. bennetii Hook. f. T., Laiya.
- Allanthus pongelion Blanco, ed. 1, 380; ed. 2, 268. Reduced by Villar to Allanthus malabarica DC., which is certainly correct.! T., Macaisa, Balocas.

OCHNACEÆ.

Ochna fascicularis Blanco, ed. 2, 246.! A good species. Erroneously reduced by Villar to Brackenridgea. T., Aniatan.

BURSERACEÆ.

- Guiacum abilo Blanco, ed. 1, 364; Icica abilo Blanco, ed. 2, 256 = Garuga floribunda Decne.! Villar reduces the species as described in the first edition to Garuga pinnata Roxb., and the species as described in the second edition to Garuga floribunda Decne., although Blanco's descriptions manifestly apply to the same species. T., Abilo.
- Canarium album Blanco, ed. 1, 793; ed. 2, 546, non Raeusch = Canarium luzonicum Miq.! (Canarium carapifolium Perk.) Erroneously reduced by Villar to Canarium commune Linn., a closely related species. The source of Manila Elemi.! T., Pisa, Pilani. I., Anten.
- Canarium commune Blanco, ed. 1, 791, non Linn.; Canarium pimela Blanco, ed. 2, 545, non Koen. = Canarium villosum (Miq.) F.-Vill. (Canariopsis villosa Miq.; Canarium cumingii Engler.!) P., Palsahinhin, Anagatli, Anagatti. I., Anten.
- Boswellia? obliqua Blanco, ed. 2, 243 (err. typ., oblicua) = Ganophyllum obliquum (Blanco) (Ganophyllum falcatum Blume!!) Blanco's name is the earliest one published for this species and should be retained. It is described in the first edition under the native name only, Pandapanda.
- * Boswellia? integra Blanco, ed. 2, 242. Referred by Villar to Protium javanicum Burm., but as no species of this genus is at present known from the Philippines Villar's identification must be considered doubtful. This species is described in the first edition under the native name Calamansalai. Blanco's specimens were from the mountains near San Mateo, Province of Rizal, a tree fruiting in June. T., Polyagan.

MELIACEÆ.

- Plagianthus humilis Blanco, ed. 1, 526; ed. 2, 366 = Turraea humilis (Blanco) (Turraea pumila Benn. Pl. Jav. Rar. 183. 1840.) Blanco's name is the earliest for this species and should be retained.
- Melia azedarach Blanco, ed. 1, 345, non Linn.; Melia composita Blanco, ed. 2, 241, non Willd. = Melia candollei Juss.! Erroneously reduced by Villar to Melia dubia Cav., a species that does not extend to the Philippines. T., Malongain.
- Turraea decandra Blanco, ed. 1, 347; Turraea virens Blanco, ed. 2, 243, non Linn. = Dysoxylon decandrum (Blanco) (Dysoxylon blancoi Vidal, Cat. Pl. Prov. Manila, 22; Sinopsis, Atlas, t. 39. f. A.; D. salutare F.-Vill.; D. amooroides Miq.) DeCandolle 1 retains Blanco's Turraea virens and T. decandra as distinct, uncertain species, but it is evident, however, that the two descriptions apply to the same species, as noted by Villar. This species is represented by Nos. 141, 2140, 2758, and 2776 Merrill, the two former numbers having been identified by Perkins 2 as Dysoxylon amooroides Miq. P., Iguiu, Aguiu, Taclitan, Malabangao. B., Ananangtang, Mananantang, Bacugan, Mocasisi, Bingliu, Palohambobocag, Bolong tambal. I., Pasiloag.

¹ Monog. Phan. 1: 749.

² Frag. Fl. Filip. 31. 1904.

- Trichilia pentandra Blanco, ed. 1., 355; ed. 2, 249. = Chisocheton Pentandrum (Blanco) (C. philippinum Harms; C. ceramicus F. Vill., non Miq.) T., Salaqui, Agapanga.
- *Turraea octandra Blanco, ed. 1, 349; ed. 2, 244. Reduced by Villar to Dysoxylon schizochitode (Turcz.) DC., which may possibly be correct, in which case Blanco's name should be retained. A tree flowering in August. T., Himamao.
- Sandoricum ternatum Blanco, ed. 1, 346; Sandoricum indicum Cav., Blanco, ed. 2, 242. The last name is correct.! T., Santol.
- *Trichilia tripetala Blanco, ed. 1, 354; ed. 2, 248. Reduced by Villar to Amoora rohituka W. et A., a species not known from the Philippines. Probably correct as to the genus although excluded from the Meliaceæ by DeCandolle.
- Lansium domesticum Jack, Blanco, ed. 2, 228.! This species is described in the first edition under the native name Lansones. T., Lansone. V., Bulahan, Boocan, Boboa.
- * Portesia rimosa Blanco, ed. 1, 297; Trichilia rimosa Blanco, ed. 2, 250. Reduced by Villar to Amoora canarana Hiern. Probably correct as to the genus, but the species does not extend to the Philippines. Blanco's specimens were from the Province of Batangas, a tree flowering in May. T., Busilac.
- *Trichilia? volubilis Blanco, ed. 2, 249. Reduced by Villar to Aglaia grandis Miq., a species not known from the Philippines. Excluded from the Meliaceæ by DeCandolle, which is probably correct. A vine flowering in January. T., Gogong toco, Cubatili, Gogong bisaya.
- * Argophyllum pinnatum Blanco, ed. 1, 186; ed. 2, 131. Reduced by Villar to Aglaia angustifolia Miq., a species that does not extend to the Philippines, probably, however, correct as to the genus. Blanco's specimens were from Angat, Province of Bulacan, a shrub flowering in March.
- Melia iloilo Blanco, ed. 2, 241 = Aglaia argentea Blume! Described in the first edition under the native name only. P., Iloilo.
- Xylocarpus granatum Koenig., Blanco, ed. 1, 298; ed. 2, 207.! T., Tabigui, Nigui, Calumpang sa lati. P., Migi.
- *Cedrela odorata Blanco, ed. 1, 184; ed. 2, 130, non Linn. Reduced by Villar to Cedrella toona Roxb., a species not known from the Philippines. Correct as to the genus. T., P., Calantas. B., Lanigpa, Lanigda.
- *Cedrela taratara Blanco, ed. 2, 131. A species known only from Blanco's description, whose specimens were from Bohol. T., Taratara. V., Sandana. B., Bolongcauit.

CHAILLETIACEÆ.

- * Elaeocarpus? integrifolius Blanco, ed. 2, 306. Reduced by Villar to Chailletia gelonioides Hook. f., which must be considered doubtful, as this species has not been found in the Philippines. Described in the first edition under the native name, Bitlag. A tree flowering in March.
- * Quilesia sericea Blanco, ed. 1, 177; ed. 2, 125. Reduced by Villar to Chailletia griffithii Hook. f., a species not known from the Philippines, and which must accordingly be considered doubtful. A shrub found in Ilocos Province, flowering in January. I., Decdec.
- *Riana tricapsularis Blanco, ed. 1, 850; ed. 2, 126. Reduced by Villar to Chailletia helferiana Kurz, a species that certainly does not extend to the Philippines. A small shrub flowering in July. Blanco's specimens were from Angat, Province of Bulacan, no native name given.

OLACACEÆ.

Fissilia psittacorum Blanco, ed. 1, 28; ed. 2, 20, non Lam. = Olaw imbricata Roxb.! T., Malabaguio, I., Tadiag.

CELASTRACEÆ.

- Diosma serrata Blanco, ed. 1, 168; ed. 2, 119 = Celastrus paniculata Willd.! T., Bilogo.
- Cupania spinosa Blanco, ed. 1, 184; ed. 2, 204 = Gymnosporia montana Roxb.! T., Malarayap.
- * Hippocratea volubilis Blanco, ed. 1, 27; ed. 2, 20. Reduced by Villar to *Hippocratea obtusifolia* Roxb., a species as yet not collected in the Philippines. Probably *Hippocratea indica* Willd. T., *Tatlong*.
- Salacia sinensis Blanco, ed. 1, 26; ed. 2, 19. Reduced by Villar to Salacia prinoides DC., which is probably correct. T., Matang olang.
- *Comocladia serrata Blanco, ed. 1, 30. Reduced by Villar to Salacia oblonga Wall., a species that certainly does not extend to the Philippines. Blanco does not include this species in the second edition. Blanco's specimens were from the seashore at Batangas, flowering in February, no native name given.

RHAMNACEÆ.

- Enrilla dichotoma Blanco, ed. 1, 709; Ventilago monoica Blanco, ed. 2, 124 = VENTILAGO DICHOTOMA (Blanco) (Ventilago luzonensis Vidal, Rev. Pl. Vasc. Filip. 90. 1886.) Villar erroneously reduces this species to Ventilago maderaspatana Gaertn. T., Salapao, Silipao.
- Rhamnus jujuba Linn., Blanco, ed. 1, 172; Zizyphus jujuba (Linn.) Lam., Blanco, ed. 2, 120. The last name is correct. Sp.-Fil., Manzanitas.

- Rhamnus zizyphus Blanco, ed. 1, 171, non Linn.; Zizyphus lotus Blanco, ed. 2, 121, non Lam. = Zizyphus trinervis Poir.! Erroneously reduced by Villar to Zizyphus oenoplia Mill. T., Duclap, Talanai. I., Labba labba, Lanoti.
- * Rhamnus talanai Blanco, ed. 1, 170; Zizyphus latifolia Blanco, ed. 2, 121, non Poir. Reduced by Villar to Zizyphus œnoplia Mill., which is certainly an error. Possibly a form of Zizyphus trinervis Poir. Blanco's specimens were from San Mateo, Province of Rizal. T., Talinas.
- Rhamnus zonulatus Blanco, ed. 1, 172 = Zizyphus zonulatus Blanco, ed. 2, 120.! (Zizyphus arborea Merrill, Govt. Lab. 1:11. 1904.!)
 Blanco's second name is the correct one for this species, which was erroneously referred by Villar to Zizyphus xylopyrus Willd. T., Balacat.
- *Rhamnus dalanta Blanco, ed. 1, 170 = Zizyphus dalanta Blanco, ed. 2, 121. Probably a distinct species. Blanco's specimens were from Bonbonon, Negros. V., Dalanta.
- Rhamnus carolinianus Blanco, ed. 1, 169; ed. 2, 119, non Walt. = Colubrina asiatica L. C. Rich.! Erroneously reduced by Villar to Rhamnus wightii Wight et Arn., a species that certainly does not extend to the Philippines. T., Cabatiti.
- Gouania domingensis Blanco, ed. 1, 196; ed. 2, 138, non Linn. = Gouania leptostachya DC.!

AMPELIDACEÆ.

- Cissus quadrangularis Linn., Blanco, ed. 1, 72; ed. 2, 52.!
- * Cissus vesicatoria Blanco, ed. 12, 50. Reduced by Villar to Cissus (Vitis) repens W. et A., which may possibly be correct. T., Pirapit angin.
- Cissus latifolia Blanco, ed. 2, 52, non Lam.; Cissus simplex Blanco, ed. 1, 72. Reduced by Villar to Cissus (Vitis) adnata, which is probably correct.
- Cissus acida Blanco, ed. 1, 69; ed. 2, 50, non Linn. = Cissus carnosa Lam.! T., Calitcalit, Cavilan, Pacopol. B., Lupo, Langingi, Lagini, Raguindi, P., Culutpamo.
- Cissus rubescens Blanco, ed. 1, 71; Cissus alata? Blanco, ed. 2, 51, non Jacq. Reduced by Villar to Cissus geniculata Roxb., which may be correct. T., Sampang. P., Loba.
- *Cissus pedata Blanco, ed. 1, 71; ed. 2, 52, non Lam. Reduced by Villar to *Vitis capriolata* Don., which is probably an error, as this species does not extend to the Philippines. T., Ayo.
- Aquilicia sambucina Blanco, ed. 1, 178, non Linn.; Leea sambucina Blanco, ed. 2, 126, non Blume = Leea rubra Blume.! T., Caliantang, Abangabang.

Ticoria aculeata Blanco, ed. 1, 85; Leea aculeata Blume, Blanco, ed 2, 127.

The latter name is correct.! P., Malimali.

SAPINDACEÆ.

- Cardiospermum halicacabum Blanco, ed. 1, 312; ed. 2, 218 = Cardiospermum halicacabum Linn., var. microcarpum Blume.! T., Bangcolon.
- *Sapindus koelreuteria Blanco, ed. 1, 289; Koelreuteria arborea Blanco, ed. 2, 202. Reduced by Villar to *Hemigyrosa perrottetii* Blume = *Guioa perrottetii* Radlk., which is certainly an error. Blanco gives no native name, and no locality.
- Molinaea arborea Blanco, ed. 1, 292; ed. 2, 204 = Guioa perrottetii (Blume) Radlk.! Erroneously reduced by Villar to Hemigyrosa canescens Thw., a species that does not extend to the Philippines. Blanco's specimens were from Batangas, and also from Parañaque, near Manila, flowering in January. T., Salab.
- Quassia tricarpa Blanco, ed. 1, 351; ed. 2, 246. Reduced by Villar to Sapindus saponaria Linn., and is certainly the form described by Radikofer as Sapindus saponaria Linn., forma microcarpus, also probably identical with the form described by Vidal as Sapindus turczaninowii. T., Tiquis tiquis, Catiquis, Sausanli, I., Cusiben, P., Malapalitpit.
- Sapindus saponaria Blanco, ed. 1, 292, non Linn.; Sapindus guisian Blanco, ed. 2, 201 = Sapindus rarak DC. (Dittelasma rarak Hook. f.) A tree flowering in April, no herbarium specimens known from the Philippines at the present time. T., Guisan, V., Duca.
- Sapindus edulis Blume, Blanco, ed. 2, 201 = Erioglossum rubiginosum Blume!
- * Schmidelia conferta Blanco, ed. 2, 217. Reduced by Villar to Allophylus zeylanicus Linn., a species that does not extend to the Philippines. Blanco's specimens were from Angat, Province of Bulacan, possibly not a species of Sapindaceæ.
- Aporetica ternata Blanco, ed. 1, 290, non Forst. Aporetica gemella Blanco, ed. 2, 203, non DC. = Allophylus grossedentata (Turcz.)
 Radlk.! Erroneously reduced by Villar to Allophylus cobbe Blume.
- * Aporetica penicellata Blanco, ed. 1, 291; ed. 2, 203. Reduced by Villar to Allophylus cobbe Blume, var. villosa, which is certainly an error, although certainly correct as to the genus.
- * Quassia simaruba Blanco, ed. 2, 247, non Linn. Reduced by Villar to Cupania glabrata Kurz., a species that does not extend to the Philippines. Blanco's specimens were from Angat, Province of Bulacan, fruiting in February.
- * Euphoria annularis Blanco, ed. 1, 285; ed. 2, 199. Reduced by Villar to Ratonia litoralis Benth. et Hook. f., which is certainly an error. It may be identical with Lepidopetalum perrottetii Blume. A tree flowering in August. T., Guisian, Guisihan.

- Euphoria cubili Blanco, ed. 1, 287; ed. 2, 200 = Cubilia blancoi Blume.! Reduced by Villar to Cubilia rumphii Blume, which is a synonym of C. blancoi.! See Icon. Bogor. 1, part 4, 51, tt. 92, 93. 1901. This species is represented by No. 445 Forestry Bureau, and No. 33 Decades Philippine Forest Flora. T., Cubili, Lubilubi.
- * Sapindus baccatus Blanco, ed. 1, 290; Koelreuteria edulis Blanco, ed. 2, 202. Blume transferred this species to Otophora, under the name Otophora blancoi. The species is known only from Blanco's description, his material being from the Provinces of Pampanga and Pangasinan, a tree flowering in December. P., Talinouno. I., Dirig.
- Capura pinnata Blanco, ed. 1, 264; Capura purpurata Blanco, ed. 2, 184 = Otophora fruticosa Blume.! (O. nigrescens Vill.; O. pinnata Merrill.) Erroneously reduced by Villar to Otophora paucijuga Hiern, a species that does not extend to the Philippines.
- *Ornitrophe triandra Blanco, ed. 1, 291; Schmidelia triandra Blanco, ed. 2, 218. Reduced by Villar to Schleichera trijuga Willd., which is very doubtful. Blanco gives no native name, and no locality.
- * Euphoria didyma Blanco, ed. 1, 288; ed. 2, 201. Referred by Villar to Nephelium glabrum Noronh., which is possibly correct. T., Guisian, Guisian, Tinaingui.
- Euphoria litchi Blanco, ed. 1, 285; ed. 2, 199, non Juss. = Euphoria cinerea Radlk.! T., Alpai, Lechias, Alipai, Alupai. B., Boboa.
- Ptelia arborea Blanco, ed. 1, 63; Seringia lanceolata Blanco, ed. 2, 45=

 Harpullia arborea (Blanco) Radlk.! Erroneously reduced by Villar
 to Harpullia cupanioides Roxb. T., Poas. I., Oas.
- Melicocca triptera Blanco, ed. 2, 203 = Tristira triptera (Blanco) Radlk.!

 Dodonæa angustifolia Linn. f., Blanco, ed. 1, 312; ed. 2, 217 = Dodonæa viscosa Linn. var. vulgaris Benth.! T., Calapinai.

ANACARDIACEÆ.

- Mangifera indica Linn., Blanco, ed. 1, 179; ed. 2, 127.! T., Manga.
- Mangifera rostrata Blanco, ed. 2, 129. Reduced by Villar to Mangifera silvatica Roxb., which is probably an error, as Blanco's species appears to be only a form of the common Mangifera indica Linn. Sp.-Fil., Manga-pico.
- Mangifera altissima Blanco, ed. 1, 181; ed. 2, 129.! Reduced by Villar to Mangifera longipes, a species not as yet discovered in the Philippines. A distinct species related to Mangifera quadrifida Jack. See Merrill, Govt. Lab. Publ. 17: 27. 1904. T., Pajo, Pao.
- * Mangifera anisodora Blanco, ed. 2, 129. Reduced by Villar to Mangifera macrocarpa Blume (M. fragrans Maingay.). This may be a distinct species, or a form of Mangifera indica Linn. Blanco saw no specimens, but established his species on hearsay. His information was from the pueblo of Sinait, Province of Ilocos Norte. Sp.-Fil, Manga de anis.

- * Mangifera pinnata Blanco, ed. 1, 182, non Lam. This species is not included in the second edition, and is certainly not a Mangifera. It may be a species of some other genus of Anacardiaceæ, or may be a species of the Meliaceæ. T., Tagapi.
- Cassuvium reniforme Blanco, ed. 1, 322; ed. 2, 227 = Anacardium occidentale Linn.! T., Casoy. I., Bollugo.
- Fagara decandra Blanco, ed. 1, 66; ed. 2, 48 = Buchanania florida Schauer. var. arborea Engl.! T., Balinhasa.
- Helicteres pinnata Blanco, ed. 1, 384; Crytocarpa quinquestila Blanco, ed. 2, 269 = Koordersiodendron pinnatum (Blanco) Merrill, Forestry Bureau, Bull. 1: 33. 1903. (K. celebicum Engl.) T., Amoguis.
- Semecarpus cuneiformis Blanco, ed. 1, 220; ed. 2, 155. Apparently a form or variety of Semecarpus perrottetii March.! T., Ligas.
- Semecarpus anacardium Blanco, ed. 1, 217; ed. 2, 152, non Linn. f. = Semecarpus perrottetii March.! Erroneously reduced by Villar to Semecarpus pubescens Thw. T., Ligas. I., Camiring. V., Langas.
- Spondias dulcis Blanco, ed. 1, 390; ed. 2, 273, non Forst. = Spondias lutea Linn.! Erroneously reduced by Villar to Spondias purpurea Linn. Sp.-Fil., Sirihuelas.
- Poupartia pinnata Blanco, ed. 1, 392; ed. 2, 275 = Spondias mangifera Willd.! T., B., Dao.
- Paliurus edulis Blanco, ed. 1, 173; Paliurus Iamio, Blanco, ed. 2, 122 = Dracontomelum cumingianum Baill.! Although Blanco's name is the earlier, Baillon's should be retained, as Blanco in establishing the species describes the flowers (and leaves?) of some species of Canarium, while the rest of his description applies to Dracontomelum. Blanco undoubtedly intended to describe the species known to-day as Dracontomelum cumingianum, as that species is universally known by the Tagalogs by the name of Lamio. T., Lamio, Malaiyo. V., Batoan.
- Paliurus dao Blanco, ed. 1, 174; ed. 2, 122 = Dracontemelum mangiferum Blume.! T., Dao.

MORINGACEÆ.

Moringa oleifera Lam., Blanco, ed. 1, 341; ed. 2, 238.! T., Malungay, Camalungai, Calungai. V., P., Malungai, Calungai, Dool, Malungit.

CONNARACEÆ.

- Cnestis volubilis Blanco, ed. 1, 385; Cnestis trifolia Blanco, ed. 2, 270, non Lam. = ROUREA VOLUBILIS (Blanco) (Rourea heterophylla Planch.) Blanco's name is the earlier, and should be retained. T., Camagsa taquilis, Palo Santo, Guicos guicos. V., P., Hanambabao, Ungalina mapula, Magtabig, Mavindato.
- Cnestis diffusa Blanco, ed. 1; Cnestis polyphylla Blanco, ed. 2, 270 = Cnestis ramiflora Griff.! Erroneously reduced by Villar to Rourea rugosa Planch. T., Ibaiban.

- * Cnestis erecta Blanco, ed. 1, 387; Omphalobium pictum Blanco, ed. 2, 271. Reduced by Villar to Connarus monocarpus Linn., a species which certainly does not extend to the Philippines. Possibly a distinct species. A small shrub, flowering in October, no native name given.
- *Cnestis corniculata Blanco, ed. 1, 386; ed. 2, 270, non Lam. Reduced by Villar to Connarus ferrugineus Jack., which is certainly an error. Blanco's specimens were from the pueblo of Tagudin, Province of Ilocos Sur. I., Sal-laday.
- Cnestis glabra Blanco, ed. 1, 387; ed. 2, 271, non Lam. Reduced by Villar to Connarus paniculatus Roxb., a species that certainly does not extend to the Philippines. Certainly a synonym of Rourea multiflora Planch.! No locality or native name given.

LEGUMINOSÆ.

- Crotalaria linifolia Linn. f., Blanco, ed. 1, 570; Quirosia secunda Blanco, ed. 2, 398. The former name is correct.
- Crotalaria pallida Blanco, ed. 1, 570, non Ait.; Crotalaria pumila Blanco, ed. 2, 397, non Schrank = Crotalaria sessiliflora Linn.!
- Phaseolus bulai Blanco, ed. 1, 572; Quirosia anceps Blanco, ed. 2, 398 = Crotalaria verrucosa Linn.! T., Bulai lava.
- Crotalaria quinquefolia Linn., Blanco, ed. 1, 569; ed. 2, 397.! T., Catanda, Susoi, Susosusoyan, Balatong aso.
- * Liparia badocana Blanco, ed. 1, 597; Psoralea badocana Blanco, ed. 2, 416. Retained by Villar as a distinct species under the second name, which may be correct, citing *Meladenia densiftora* Turcz., as a synonym. Blanco's specimens were from the pueblo of Badoc, Ilocos Province. No native name given.
- Amorpha glandulosa Blanco, ed. 1, 555; Dalea alopecuróides Blanco, ed. 2, 389, non Willd. = Dalea glandulosa (Blanco) (Dalea nigra Mart. et Gal.). Although this plant is a native of America, yet the earliest description is that of Blanco, and his name should be retained. T., Durang parang, Camangi.
- Indigofera angustifolia Blanco, ed. 1, 596; ed. 2, 415, non Linn. = Indigofera hirsuta Linn.! I., Tayomtayom.
- Indigofera argentea Blanco, ed. 2, 415, non Linn. = Indigofera tinctoria Linn.!
- Indigofera tinctoria Blanco ed. 1, 591; ed. 2, 413, non Linn. = Indigofera anil Linn.! T., Tayom. P., Tayung. V., Tagung.
- Indigofera hirsuta Blanco, ed. 1, 591; non Linn.; Indigofera senegalensis Blanco, ed. 2, 412, non Lam. = Tephrosia purpurea Pers.! T., Tayomtayoman. V., Dagangdang. P., Maasic.
- Cylista piscatoria Blanco, ed. 1, 589; Galactia ? terminiflora Blanco, ed. 2, 411 = MILLETTIA PISCATOBIA (Blanco). Erroneously reduced by Villar to *Milletia splendens* W. et A., a species that certainly does not extend to the Philippines. T., *Tubli*.

- Galedupa pungam Blanco, ed. 1, 558; ed. 2, 390, non Gmel. = Gliricidia sepium (Jacq.) Steud.! (G. maculata H. B. K.) Sp.-Fil., Madre cacao.
- Coronilia emerus Blanco, ed. 1, 582, non Linn.; Sesbania cannabina Blanco, ed. 2, 418. Reduced by Villar to Sesbania aculeata, var. paludosa Baker, which is probably correct. T., Malacaguois.
- Sesbania grandiflora Pers., Blanco, ed. 1, 599; ed. 2, 418.! T., Caturai.
- Lupinus angustifolius Blanco, ed. 1, 566, non Linn.; Smithia bigeminata Blanco, ed. 2, 395. Reduced by Villar to Zornia diphylla Pers., which is probably correct.
- Arachia hypogaea Linn., Blanco, ed. 1, 567; ed. 2, 396.! T., Mani.
- Hedysarum vespertilionis Linn., Blanco, ed. 1, 581; ed. 2, 407 = Lourea vespertilionis (Linn.) Desv.
- Tetragonolobus simplicifolius Blanco, ed. 397. This species was reduced by Villar to Alyscicarpus tetragonolobus Edgw., a species that certainly does not extend to the Philippines. It is undoubtedly identical with Alyscicarpus bupleurifolius DC. Blanco's specimens were from Para-fiaque, near Manila.
- Aeschynomene arborea Blanco, ed. 1, 581; ed. 2, 406 = Desmodium umbellatum (Linn.) DC.!
- Cytisus quinquepetalus Blanco, ed. 1, 598; Cajanus quinquepetalus Blanco, ed. 2, 417. Reduced by Villar to Desmodium cephalotes Wall., which is probably correct.
- Hedysarum puicheilum Linn., Blanco, ed. 1, 581; Dicerma puicheilum DC., Blanco, ed. 2, 407 = Desmodium pulcheilum (Linn.) Benth.! T., Payangpayang. V., Calaicai.
- Hippocrepis rhomboidea Blanco, ed. 1, 585; Desmodium spirale DC., Blanco, ed. 2, 408. The latter name is correct.
- Hippocrepis comosa Blanco, ed. 1, 584, non Linn.; Desmodium diversifoilum Blanco, ed. 2, 408, non DC. = Desmodium gangeticum DC.! T., Manquit.
- Hippocrepis multisiliquosa Blanco, ed. 1, 584, non Linn.; Desmodium gangeticum Blanco, ed. 2, 408, non DC. = Desmodium latifolium DC.! T., Manquit.
- Hippocrepis humilis Blanco, ed. 1, 585; Desmodium parvifolium Blanco, ed. 2, 408, non DC. = Desmodium triflorum DC.! T., Pacpac langao.
- Abrus precatorius Linn., Blanco, ed. 1, 565; ed. 2, 394.! T., Saga, Sagamamin, Bangati. B., Bangati, Gicos gicos, Agaiyangyiang, Mangadolong, Caloo, Matangpune, Aroyangyang. P., Cansasaga. I., Bugayon.
- Negretia urens Blanco, ed. 1, 586; ed. 2, 409 = Mucuna imbricata DC.!
 (No. 3783 Merrill.) Erroneously reduced by Villar to Mucuna monosperma DC., a species that certainly does not extend to the Philippines.
 A vine, flowering in November. T., Buquiquit, Lipai.
- Negretia pruriens Blanco, ed. 2, 411. Reduced by Villar to Mucuna atropurpurea DC., which may be correct. V., Nipai, Lipai.

- Negretia mitis Blanco, ed. 1, 588; ed. 2, 410, non Beauv. Reduced by Villar to Mucuna nivea DC., but probably only a form of Mucuna pruriens DC. T., Garbanzos, Habas.
- Erythrina carnea Blanco, ed. 1, 564; ed. 2, 393, non Ait. = Erythrina indica Lam.! T., B., P., Dapdap. T., Casindac, B., Cabrab. P., Sulbang.
- Erythrina picta Blanco, ed. 1, 565, non Linn.; Erythrina caffra Blanco, ed. 2, 394, non Thunb. = Erythrina ovalifolia Roxb.! T., Anii.
- Dolichos ensiformis Linn., Blanco, ed. 1, 577; Canavalia gladiata Blanco, ed. 2, 403 = Canavalia ensiformis (Linn.) DC.! T., Habas. V., Magtambocao.
- Dolichos acinaciformis Blanco, ed. 1, 578, non Jacq.; Canavalia ensiformis Blanco, ed. 2, 404, non DC. = Canavalia obtusifolia Cav.! T., Pataning dagat.
- Dioscorea bolojonica Blanco, ed. 1, 800; ed. 2, 551; Pachyrhizus teres Blanco, ed. 1, 580; Pachyrhizus montanus Blanco, ed. 2, 406 = Pueraria phaseoloides (Roxb.) Benth.! Villar probably correctly reduces Dioscorea bolojonica Blanco to Pueraria, Blanco having described that species from leaf specimens only. The native name given by Blanco for his Dioscorea is Bahai., for Pachyrhizus, Sincamas aso.
- Phaseolus inamoenus Blanco, ed. 1, 571; ed. 2, 399, non? Linn. = Phaseolus vulgaris Linn.! This is reduced by Villar to a variety of Phaseolus lunatus Linn. T., Patani.
- * Phaseolus ilocanus Blanco, ed. 1, 572; Phaseolus tunkinensis Blanco, ed. 2, 399, non? Lour. Reduced by Villar to a variety of *Phaseolus lunatus* Linn., probably only a cultivated form of this species. Sp.-Fil., *Frijoles del Abra*.
- * Phaseolus vexillatus Blanco, ed. 1, 574, non Linn.; Phaseolus vulgaris
 Blanco, ed. 2, 401, non Linn. Reduced by Villar to a variety of
 Phaseolus lunatus Linn., which may be correct. Much cultivated in
 the Province of Batangas. T., Buttingi, Biringi.
- * Phaseolus caracalla Linn., Blanco, ed. 1, 575; ed. 2, 401. Retained under this name by Villar; doubtful. T., Sitao.
- Phaseolus mungo Linn., Blanco, ed. 1, 573; ed. 2, 400.! T., Balatong, Mongos.
- * Dolichos trilobus Blanco, ed. 2, 403, non Linn. Reduced by Villar to *Phaseolus calcaratus* Roxb., which may be correct. Blanco's specimens were from Guadalupe, near Manila, flowering in January.
- Phaseolus lunatus Linn., Blanco, ed. 1, 573; ed. 2, 400.! T., Zabache.
- * Dolichos repens Blanco, ed. 1, 577; ed. 2, 402, non Linn. Reduced by Villar to *Vigna repens* Baker, which is certainly an error, as the latter species does not extend to the Philippines. Blanco's specimens were from Batangas, flowering in August. No native name given.

- Dollchos sesquipedalis Blanco, ed. 2, 402, non Linn.; Dollchos echinulatus Blanco, ed. 2, 401 = Vigna catjang Endl.! T., Quibal, Sitao.
- Pachyrhizus jicamas Blanco, ed. 1, 579; Pachyrhizus angulatus Rich., Blanco, ed. 2, 405. The latter name is correct. T., Sincamas, Hicamas.
- Clitorea ternatea Linn., Blanco, ed. 1, 590; ed. 2, 412.! T., Colocanting.
- Glycine lucida Blanco, ed. 1, 578, non Forst.; Lablab cultratus DC., Blanco, ed. 2, 405 = Dolichos lablab Linn.! T., Bulai, Batao.
- Dolichos tetragonolobus Linn., Blanco, ed. 1, 576; ed. 2, 402 = Psophocarpus tetragonolobus (Linn.) DC.! T., Calamismis. I., Pal-lam.
- Cytisus volubilis Blanco, ed. 1, 599; Cajanus volubilis Blanco, ed. 2, 417. Reduced by Villar to Atylosia mollis Benth., which is probably correct.
- Cytisus cajan Linn., Blanco, ed. 1, 597; Cajanus bicolor DC., Blanco, ed. 2, 416 = Cajanus indicus Spreng.! T., Caguois.
- *Amerimnum mimosella Blanco, ed. 1, 563; ed. 2, 393. Reduced by Villar to Dalbergia lanceolaria Linn., a species that certainly does not extend to the Philippines. Prain retains this as an imperfectly known species, Dalbergia mimosella (Blanco) Prain. Blanco's specimens were from Tala, Province of Rizal or Bulacan, flowering in June. T., Macapil.
- Pterocarpus pallidus Blanco, ed. 1, 560; ed. 2, 391 = Pterocarpus indicus Willd.! T., Asana. C., V., Naga. P., Daitanag.
- Pterocarpus santalinus Blanco, ed. 1, 561; ed. 2, 392, non Linn. = Pterocarpus blancoi Merrill, Govt. Lab. Publ. 6:7. 1904. Pterocarpus santalinus Linn., does not extend to the Philippines. T., Narra, Naga. P., Apalit, Daitanag. V., Sanque?.
- Galedupa maculata Blanco, ed. 1, 559; ed. 2, 390 = Pongamia glabra Vent.! T., Balicbalic.
- Galedupa frutescens Blanco, ed. 1, 559; ed. 2, 391. Reduced by Villar to *Derris scandens* Benth., which is probably correct. T., *Malasaga*.
- Pterocarpus frutescens Blanco, ed. 1, 562; ed. 2, 392. Reduced by Villar to *Derris uliginosa* Benth., which is probably correct. T., Silasila.
- * Pterocarpus diadelphus Blanco, ed. 1, 563; ed. 2, 393. Reduced by Villar to Derris thyrsiflora Benth., a species not known from the Philippines. Flowering in June. T., Silasila.
- Sophora tomentosa Linn., Blanco, ed. 1, 328; ed. 2, 229.! T., B., Tambalisa, Cabaicabai, Yabag, Mantala, etc.
- Sophora heptaphylla Blanco, ed. 1, 329; ed. 2, 229, non DC. Villar considers that Blanco describes DeCandolle's species, but Sophora heptaphylla certainly does not extend to the Philippines. It is apparently but a form of the preceding. T., Tambalisa, Cabaicabai.

¹ Ann. Bot. Gard., Calcutta, 10: 42. 1904.

- Ormosia calavensis Blanco, ed. 2, 230.!
- Guilandina bonducella Linn., Blanco, ed. 1, 343; ed. 2, 239 = Caesalpinia bonducella (Linn.) Fleming.! T., Bayag cambing, Calambibit. V., Dalugdug.
- Guilandina nuga Linn., Blanco, ed. 1, 344; ed. 2, 240 = Caesalpinia nuga (Linn.) Ait.! T., Camit cabag.
- Caesalpinia sappan Linn., Blanco, ed. 1, 335; ed. 2, 234.! T., Sapang, Sibucao.
- Poinciana pulcherrima Linn., Blanco, ed. 1, 333; ed. 2, 232 = Caesalpinia pulcherrima (Linn.) Swartz.! Sp.-Fil., Flores, Rosas, Caballero.
- Caesalpinia torquata Blanco, ed. 1, 336; Mezoneurum procumbens
 Blanco, ed. 2, 235 = Mezoneurum glabrum Desf.! T., Sagnit, Sapnit,
 Capitcabag. P., Sapnit. V., Tugabang, Ugabang.
- Caesalpinia ignota Blanco, ed. 1, 336; ed. 2, 235 = Mezoneurum pubescens Desf.!
- Mimosa membranulacea Blanco, ed. 1, 739; Reichardia pentapetala Blanco, ed. 2, 233 = Pterolobium indicum A. Rich.! I., Palo santo.
- Cassia fistula Linn., Blanco, ed. 1, 339; ed. 2, 237.! Blanco includes in this species Cassia javanica Linn. T., P., Canafistula. V., Lombayong, Ibabao, Balayong.
- Cassia occidentalis Linn., Blanco, ed. 1, 338; ed. 2, 236.! T., Tighiman.
- Cassia tora Linn., Blanco, ed. 1, 337; ed. 2, 235.! T., Manimanihan, Mongomongohan, Catandang aso.
- Cassia longisiliqua Blanco, ed. 1, 338, non Linn.; Cassia sulcata Blanco, ed. 2, 236, non DC. = Cassia hirsuta Linn.! T., Tighiman.
- Cassia alata Linn., Blanco, ed. 1, 339; ed. 2, 237.! T., Sonting, Acapulco, Gamot sa buni, Catanda. V., Sunting, Casitas. P., Pacayomcom Castila.
- Cassia mimosoides Linn., Blanco, ed. 1, 340; ed. 2, 237.!
- Schotia speciosa Blanco, ed. 1, 356; ed. 2, 251, non Linn. Reduced by Villar to Cynometra ramiflora Linn., var. mimosoides Baker, which is probably an error. Doubtless the same as Cynometra inequalifolia A. Gray. T., Balitbitan.
- Crudia spicata Blanco, ed. 2, 261, non Willd. = Crudia blancoi Rolfe.! T., Malatumbaga, Hintotoor.
- Tamarindus indica Linn., Blanco, ed. 1, 29; ed. 2, 20.! T., P., C., V., Sampaloc, Maca sampaloc. V., Sambac, Sumalagui, Camalagui, Sampalagui. I., Salomagui.
- Eperua decandra Blanco, ed. 1, 368; ed. 2, 259 = Intsia bijuga O. Kuntze.! (Afzelia bijuga A. Gray.) T., Ipil, Taal.
- Eperua falcata Blanco, ed. 1, 369, non Aubl.; Eperua rhomboidea Blanco,
 ed. 2, 260 = Pahudia rhomboidea (Blanco) Prain. (Afzelia rhomboidea Vidal.) T., Tindalo, Balayon.

- Bauhinia tomentosa Blanco, ed. 1, 330; ed. 2, 230, non Linn. Reduced by Villar to Bauhinia malabarica Roxb., which is probably correct.
 T., Alibanban. B., Livas, Alibanban, Balibanban, Marulinao, Diis, Ahihiro, Alambihor, Alibihil. P., Alibanban.
- Bauhinia binata Blanco, ed. 1, 331; ed. 2, 231. Reduced by Villar to Bauhinia blancoi Baker, but if this is correct Blanco's name is the earlier. T., Alibanban.
- Bauhinia scandens Blanco, ed. 1, 332; ed. 2, 232, non Linn. = Bauhinia cumingiana Benth.! Erroneously referred by Villar to Bauhinia vahlii W. et A., a species that does not extend to the Philippines. T., Banot.
- * Bauhinia castrata Blanco, ed. 1, 331; Bauhinia purpurea Blanco, ed. 2, 231, non Linn. Considered by Villar to be identical with Bauhinia purpurea Linn., but this is certainly an error. Blanco's specimens were from Pasig, flowering in October. Very near Bauhinia malabarica Roxb. T., Alibanban.
- * Bauhinia grandiflora Blanco, ed. 1, 332; ed. 2, 231, non Juss. Reduced by Villar to Bauhinia variegata Linn., a species not known from the Philippines. Blanco's specimens were from Tayabas, a tree flowering in August. T., Bongalon.
- Mimosa acle Blanco, ed. 1, 738; ed. 2, 509 = Pithecolobium acle (Blanco) Vidal.! Erroneously reduced by Villar to Xylia dolabriformis Benth. T., Acle.
- Adenanthera gogo Blanco, ed. 1, 353; Entada pursaetha DC., Blanco, ed. 2, 247 = Entada scandens (Linn.) Benth.! T., Gogo, Bayogo. B., P., Balogo, Gogong bacay.
- Mimosa virgata Blanco, ed. 1, 737; Mimosa punctata Blanco, ed. 2, 508 = Adenanthera pavonina Linn.! T., Quinasaicasai.
- Mimosa peregrina Blanco, ed. 1, 737; ed. 2, 509, non Linn. = Parkia roxburghii G. Don.! T., Copang.
- Mimosa asperata Blanco, ed. 1, 732; ed. 2, 505, non Willd. = Mimosa pudica Linn.! T., Macahiya.
- * Mimosa quadrivalvis Blanco, ed. 1, 732; ed. 2, 506. Considered by Villar to be the same as Schranckia aculeata Willd., an American species, and one not definitely known from the Philippines. Blanco's specimens were from Bauang, Province of Batangas, flowering in September. T., Sapinit.
- Mimosa farnesiana Linn., Blanco, ed. 1, 729; ed. 2, 504 = Acacia farnesiana (Linn.) Willd.! T., Aroma.
- Mimosa tenuifolia Blanco, ed. 1, 739; ed. 2, 510, non Linn. = Acacia concinna DC.! T., Sibog.
- Mimosa lebbek Blanco, ed. 1, 733; ed. 2, 506, non Linn. Reduced by Villar to Albizzia retusa Benth., which is probably correct. T., Langil.
- Mimosa coriaria Blanco, ed. 1, 734; ed. 2, 506 = Albizzia procera Benth.! T., Ayangao, Dariangao, Ananaplas. P., Anitap. I., Adaan.

- Mimosa carisquis Blanco, ed. 1, 731; ed. 2, 507. Reduced by Villar to Albizzia julibrissin Durazz., which is probably correct. T., Carisquis.
- Mimosa unguis-cati Blanco, ed. 1, 731, non Linn.; Inga lanceolata Blanco, ed. 2, 370, non HBK. = Pithecolobium dulce Benth.! T., Camochiles, Camonsiles.
- Mimosa scutifera Blanco, ed. 1, 735; ed. 2, 507. Reduced by Villar to Pithecolobium lobatum Benth., which is probably correct, although held by Bentham to be a distinct species, Pithecolobium scutiferum. T., Anagap.

ROSACEÆ.

- * Prunus armeniaca Blanco, ed. 2, 290, non Linn. Reduced by Villar to Prunus triflora Roxb., which is evidently an error. Blanco's specimens were from cultivated plants in Manila and in the Province of Cagayan. Sp.-Fil., Albaricoques.
- Rubus moluccanus Linn., Blanco, ed. 1, 428; ed. 2, 298. Considered by Villar to have been correctly identified by Blanco. T., Dagamit.
- Rubus jamaicensis Blanco, ed. 1, 427, non Linn.; Rubus comintanus Blanco, ed. 2, 298. Reduced by Villar to Rubus rosæfolius Smith, which is probably correct. T., Sapinit, Sagmit.

SAXAFRAGACE AL

* Malesherbia globosa Blanco, ed. 2, 454. Reduced by Villar to Hydrangea oblongifolia Blume, which is probably an error, possibly the same as Hydrangea lobbii Maxim. Blanco's specimens were from Bolhoon, Cebu. V., Mampol.

CRASSULACEÆ.

- Cotyledon paniculata Blanco, ed. 1, 381, non Linn. f.; Bryophyllum germinans Blanco, ed. 2, 220 = Bryophyllum calycinum Salisb.! T., Angelico, Catacataca.
- Cotyledon serrata Blanco, ed. 1, 382, non Linn.; Bryophyllum serratum Blanco, ed. 2, 220 = Kalanchæ laciniata DC.!
- Cotyledon lanceolata Blanco, ed. 1, 382, non Forsk; Bryophyllum triangulare Blanco, ed. 2, 221 = Kalanchæ spathulata DC.!

DROSERACEÆ.

Drosera hexagynia Blanco, ed. 1, 226; ed. 2, 159 = Drosera indica Linn.! T., Hintipalo.

RHIZOPHORACEÆ.

- Rhizophora mangle Blanco, ed. 1, 397; ed. 2, 278, non Linn. = Rhizophora mucronata Lam.! T., Bacao, Bacauan.
- Rhizophora longissima Blanco, ed. 1, 398; ed. 2, 278 = Rhizophora conjugata Linn.! T., Tangal. V., Tongog.

- Rhizophora candel Blanco, ed. 1, 396; ed. 2, 277, non Linn. = Ceriops candolleana Arn.! T., Ligasin, Tigasan. V., Pototan.
- Rhizophora tinctoria Blanco, ed. 1, 394; Rhizophora gymnorrhiza Linn., Blanco, ed. 2, 276 = Bruguiera gymnorrhiza Lam.! T., Bacao, Bacauan. I., Oongon.
- Rhizophora polyandra Blanco, ed. 1, 396; ed. 2, 277 = Bruguiera eriopetala W. et A.! T., Bacao, Bacauan.
- Rhizophora plicata Blanco, ed. 1, 398; ed. 2, 279. Reduced by Villar to Bruguiera malabarica Arn., a species not known from the Philippines, probably only a form of Bruguiera eriopetala W. et A. T., Bacao, Bacauan.
- Brugulera nemorosa Blanco, ed. 2, 275. Reduced by Villar to Carallia integerrina DC., which is probably correct.

COMBRETACEÆ.

- Terminalia latifolia Blanco, ed. 1, 376, non Swartz; Terminalia mauritiana Blanco, ed. 2, 264, non Lam. = Terminalia catappa Linn.! T., P., Talisai. P., Banilac, Nato, Hitam, Dalasa, Calisai. I., Lugo, Pandan.
- Terminalia angustifolia Blanco, ed. 1, 377, non Jacq.; Terminalia edulis Blanco, ed. 2, 265. Erroneously reduced by Villar to Terminalia belevica Roxb., a species that does not extend to the Philippines. A very characteristic species, the second name being the correct one. T., Calumpit.
- *Bucida comintana Blanco, ed. 1, 856; ed. 2, 265. Erroneously reduced by Villar to *Terminalia chebula* Retz., a species not known from the Philippines. Certainly not a species of *Terminalia*, but judging from Blanco's description, a species of *Calycopteris*. Blanco's specimens were from the town of San Jose, Province of Batangas, a tree flowering in May. T., *Comintan*, *Dinglas*.?
- Gimbernatia calamansanay Blanco, ed. 2, 266 = Terminalia calamansanay (Blanco) Rolfe.! Erroneously referred by Villar to Terminalia bialata Kurz, a species that does not extend to the Philippines. T., Calamansanai.
- Petaloma coccinea Blanco, ed. 1, 345; ed. 2, 240 = Lumitzera purpurea Presl.! (Lumnitzera coccinea W. et A.) T., Culasi.
- Petaloma alba Blanco, ed. 1, 344; ed. 2, 240 = Lumnitzera racemosa Willd.! T., Culasi.
- Combretum distillatorium Blanco, ed. 1, 295; Combretum laxum Blanco, ed. 2, 206, non Roxb. = Combretum squamosum Roxb.! Erroneously reduced by Villar to Combretum ovalifolium Roxb., a species that certainly does not extend to the Philippines. I., Malacadog. T., Pamulaclaquin.

- *Gnidia oppositifolia Blanco, ed. 1, 299; ed. 2, 208. Reduced by Villar to Combretum wallichii DC., which is certainly an error, as that species does not extend to the Philippines. Possibly a species of Combretaceæ. Blanco's specimens were from Angat, Province of Bulacan, a small tree, flowering in December.
- Quisqualis indica Linn., Blanco, ed. 1, 361; Quisqualis spinosa Blanco, ed. 2, 254. This species is reduced by Villar to Quisqualis malabarica Bedd., which is certainly an error. Blanco's species is certainly identical with Quisqualis indica Linn.! T., Niogniogan, Tagarao. V., Tangolon. P., Bebevabe. I., Tartarao.
- Gronovia ternata Blanco, ed. 1, 187; ed. 2, 132 = Illigera luzonensis (Presl.) Merrill, Govt. Lab. Publ. 17: 18. 1904. (Illigera meyeniana Kunth.)
- Halesia ternata Blanco, ed. 1, 399; ed. 2, 279. Reduced by Villar to *Illigera dubia* Span., which is certainly an error. Certainly identical with the preceding.!
- Gyrocarpus lobatus Blanco, ed. 2, 54 = Gyrocarpus jacquini Roxb.! This species is described by Blanco, in the first edition under the native name only, Lapolapo.

MYRTACEÆ.

- * Metrosideros pictapetala Blanco, ed. 2, 295. Reduced by Villar to Metrosideros vera Rumph., which is evidently an error. Blanco's specimens were from Cebu, a tree flowering in September. V., Barit.
- Psidium aromaticum Blanco, ed. 1, 417, non Aubl.; Psidium pyriferum Blanco, ed. 2, 292, non Linn. = Psidium guayava Linn. var.! T., Bayabas, Guyabas, Calimbahin.
- Myrtus communis Blanco, ed. 1, 422; ed. 2, 295, non Linn. = Decaspermum blancoi Vidal.! Erroneously referred by Villar to Decaspermum rubrum Blume. T., Malatumbaga.
- Eugenia jambos Blanco, ed. 1, 416; ed. 2, 290, non Linn. = Jambosa malaccensis (Linn.) DC.! T., Macupa, Yambo.
- Eugenia malaccensis Blanco, ed. 1, 415; ed. 2, 290, non Linn. = Jambosa vulgaris DC.! T., Tampoi, Calobcob. V., Yampoi. P., Balobar.
- * Eugenia montana Blanco, ed. 1, 418; ed. 2, 291, non DC. Reduced by Villar to Eugenia macrocarpa Roxb., a species not known from the Philippines. Certainly a species of Jambosa. Blanco's specimens were in part from Balanga, Province of Bataan, flowering in August. T., Copcop, Malaracopcop.
- Myrtus mananquil Blanco, ed. 1, 421; Eugenia mananquil Blanco, ed. 2, 290. Reduced by Villar to Eugenia javanica Lam., probably the same as Jambosa aequa (Roxb.) DC. A tree flowering in April. T., Mananquil.

- * Eugenia bauanguica Blanco, ed. 1, 418; ed. 2, 290. Reduced by Villar to Eugenia laeta Ham., which is certainly an error. Related to if not identical with Jambosa malaccensis (Linn.) DC. Blanco's specimens were from Bauang, Province of Batangas, a shrub flowering in March.
- * Myrtus tripinnata Blanco, ed. 1, 421; Myrtus subrubens Blanco, ed. 2, 294. Reduced by Villar to Eugenia cymosa Lam., a species not known from the Philippines. Certainly a species of Jambosa. A tree flowering in February. T., Malaruhat na pula.
- * Calyptranthes makal Blanco, ed. 1, 419, non Rauesch.; Calyptranthes zuzygium Blanco, ed. 2, 293, non Swartz. Reduced by Villar to Eugenia operculata Roxb., which is certainly an error. A species of Syzygium.? A tree flowering in January. T., Malaruhat.
- Calyptranthes jambolana Willd., Blanco, ed. 1, 418; Syzygium jambolana DC., Blanco, ed. 2, 293. The last name is correct. T., P., Lumboi, Duhat, Duat.
- * Calyptranthes ramiflora Blanco, ed. 1, 420; Syzygium latifolium Blanco, ed. 2, 249, non DC. This species is reduced by Villar to Eugenia bracteata Roxb., a species that certainly does not extend to the Philippines. Blanco's specimens were from San Jose, Province of Batangas, a tree flowering in February. T., Dinglas.
- * Eugenia glandulosa Blanco, ed. 1, 417; ed. 2, 291. A shrub flowering in July. Blanco's specimens were from Malinta, near Manila. Villar makes no attempt to reduce this species.
- * Eugenia lobas Blanco, ed. 1, 857; Eugenia cauliflora Blanco, ed. 2, 291. A species quite unknown. Blanco's specimens were from trees cultivated in Manila, the fruit very acid. T., Lobas.
- Barringtonia speciosa Forst., Blanco, ed. 1, 533; ed. 2, 373.! T., Botong, Botongbotong. V., Bitoon, Bitung, Botong.
- Barringtonia stradivium Blanco, ed. 1, 533; Barringtonia racemosa Blume, Blanco, ed. 2, 373. The latter name is correct. T., Potat.
- *Sulipa globosa Blanco, ed. 2, 348. Reduced by Villar to *Planchonia littoralis* Blume, which may be correct. Blanco's specimens were from Maragondon, Province of Cavite, the ripe fruits used for poisoning fish.

MELASTOMATACEÆ.

- Osbeckla multiflora Blanco, ed. 1, 293, non Smith; Osbeckia sinensis Blanco, ed. 2, 205 = Osbeckia chinensis Linn.!
- * Melastoma aspera Blanco, ed. 1, 368; Melastoma obvoluta Blanco, ed. 2, 259. Reduced by Villar to Melastoma malabathricum Linn., but this is probably an error, as the latter species is not known from the Philippines. Blanco's specimens were from Angat. Probably the same as Melastoma obvolutum Jack.

- Melastoma tamonea Blanco, ed. 1, 367, non Sw.; Melastoma dodecandra Blanco, ed. 2, 258 = Melastoma polyantha Blume.! Reduced by Villar to Melastoma imbricatum Wall., a species that does not extend to the Philippines. Blanco's specimens were from Panay, a shrub flowering in March.
- Melastoma malabathrica Blanco, ed. 1, 367; ed. 2, 258, non Linn. = Melastoma molle Wall.! V., Buyong.
- Memecylon parviflorum Blanco, ed. 1, 300; Memecylon tinctorium Blanco, ed. 2, 208 = Memecylon edule Roxb.! Villar reduces Blanco's species to the variety ovata Clarke. T., Colis. I., Candon. V., Saguinsin.
- Memecylon lanceolatum Blanco, ed. 1, 301; ed. 2, 209. Reduced by Villar to Memecylon cumingianum Presl., but it is probably only a form or variety of the preceding. T., Colis. I., Candong.

LYTHRACEÆ.

- Ammania monoflora Blanco, ed. 1, 64; Ammania ramosior Blanco, ed. 2, 46, non? Linn. This species is reduced by Villar to Ammania peploides Spreng. (Rotalia indica (Willd.) Koehne), which is apparently correct, although Koehne considers Blanco's species to be identical with Rotalia ramosior (Linn.) Koehne, an American species. Blanco states that the plant is very common, flowering in October, but so far as known no collector has secured specimens of Rotalia indica (Linn.) Koehne, in the Philippines.
- Celosia nana Blanco, ed. 1, 192; Ammania debilis Blanco, ed. 2, 46, non Ait. = Ammania baccifera Linn.! T., Bias pogo.
- Pemphis setosa Blanco, ed. 1, 410; ed. 2, 285 = Pemphis acidula Forst.! Lawsonia inermis Linn., Blanco, ed. 1, 294; ed. 2, 206.! Sp.-Fil., Cinamomo.
- Quilamum luteum Blanco, ed. 1, 851; ed. 2, 136 = Crypteronia paniculata Blume.! Erroneously reduced by Villar to Crypteronia pubescens Blume, var. hookeri Clarke, a species that does not extend to the Philippines. T., Quilamo.
- Lagerstroemia indica Linn., Blanco, ed. 1, 454; ed. 2, 316.! Sp.-Fil., Melindres.
- Munchausia speciosa Linn., Blanco, ed. 1, 611; ed. 2, 427 = Lagerstroemia speciosa (Linn.) Pers.! T., Banaba.
- Sonneratia pagatpat Blanco, ed. 1, 424; ed. 2, 296.! This species was erroneously reduced by Villar to Sonneratia acida Linn. f. T., Pagatpat, Palatpat, Palapad.
- Punica granatum Linn., Blanco, ed. 1, 422; ed. 2, 295.! Sp.-Fil., Granada.

ONAGRACEÆ.

- Jussiæa Inclinata Blanco, ed. 1, 366, non Linn.; Jussiæa fluviatilis Blanco, ed. 2, 257, non Blume = Jussiæa repens Linn.! T., Cangeong dapo.
- Jussizea erecta Blanco, ed. 1, 365; ed. 2, 257, non Linn. = Jussizea suffrutiosa Linn.! T., Malapaco, Balachac.?
- * Balingayum decumbens Blanco, ed. 1, 187; ed. 2, 132. This species and the genus is unknown. Villar retains it as a distinct genus of the Onagraceæ, but Bentham and Hooker, and Engler and Prantl refer it doubtfully to the genus Erythropalum in the Olacaceæ. Blanco's specimens were from Malinta, near Manila, a small prostrate plant, growing in wet places, rare and not well known to the natives. T., Balingayo.

SAMYDACEÆ.

- *Laurus serrata Blanco, ed. 1, 319; ed. 2, 224. Reduced by Villar to Casearia glomerata Roxb., a species that certainly does not extend to the Philippines. Possibly a form of Casearia grewiæfolia Vent. A shrub or small tree flowering in August. T., Inignin.
- Samyda serrulata Blanco, ed. 1, 374, non Linn.; Samyda pubescens Blanco, ed. 2, 263, non Linn. = Casearia cinerea Turcz.! Erroneously reduced by Villar to Casearia tomentosa Roxb. T. Tulibas.
- Anavinga fuliginosa Blanco, ed. 1, 372; Casearia fuliginosa Blanco, ed. 2, 262. Casearia leucolepis Turcz., is certainly a synonym of this species, and Blanco's name being the earlier should be retained. Erroneously reduced by Villar to Casearia grewiæfolia Vent.
- *Samyda trivalvis Blanco, ed. 1, 374; ed. 2, 263. Reduced by Villar to Casearia fragilis Vent., which is certainly an error, although probably correct as to the genus. Blanco's specimens were from Angat, Province of Bulacan.
- * Gordonia polysperma Blanco, ed. 1, 549; ed. 2, 384. Reduced by Villar to *Homalium foetidum* Benth., a species not known from the Philippines. Undoubtedly a species of *Homalium*. Blanco gives no locality and no native name.

PASSIFLORACEÆ.

- Passiflora minima Blanco, ed. 1, 647, non Jacq.; Passiflora serrulata Jacq., Blanco, ed. 2, 452. The second name is undoubtedly correct. A cultivated plant flowering in August.
- * Passiflora parviflora Blanco, ed. 1, 469, non Sw.; Modecca? parviflora Blanco, ed. 2, 453. Reduced by Villar to Modecca cardiophylla Masters, a species that certainly does not extend to the Philippines. Blanco's specimens were from the mountains near San Mateo, Province of Rizal, fruit edible. T., Salapong.

- * Passifiora coccinea Blanco, ed. 1, 650, non Banks; Modecca? coccinea Blanco, ed. 2, 453. Reduced by Villar to Modecca heterophylla Blume, a species as yet not known from the Philippines. Blanco's specimens were from the same locality as the preceding species. Fruit 2 inches long, 2 inches in diameter, edible, flowering in June. T., Melong ouac.
- * Passiflora saponaria Blanco, ed. 1, 650; Modecca? saponaria Blanco, ed. 2, 453. Reduced by Villar to Modecca trilobata Roxb., a species that is unknown from the Philippines. Blanco's specimens were from Tagudin, Province of Ilocos Sur. Il., Libas.
- * Passiflora zucca Blanco, ed. 1, 648; Modecca trilobata Blanco, ed. 2, 452, non Roxb. Reduced by Villar to Modecca palmata Lam., a species that certainly does not extend to the Philippines, Blanco gives no locality for his specimens, which he states flower in June. Fruit red. T., Binoyoc boyoc.
- Carica hermaphrodita Blanco, ed. 1, 803; ed. 2, 554. Carica papaya Linn., Blanco, ed. 1, 803; ed. 2, 554. The latter name is correct. Carica hermaphrodita Blanco, is an abnormal form. Sp.-Fil., Papaya.

CUCURBITACEÆ.

- Trichosanthes amara Blanco, ed. 1, 774; ed. 2, 533, non Linn. Reduced by Villar to Trichosanthes palmata Roxb., but certainly identical with T. quinqangulata A. Gray! Not Luffa acutangula Roxb., as stated by Cogniaux. T., Salagsalag, Pacupis, Salimpocot, Hothot, Halahala, Buyocbuyoc. B., P., Tabobog, Cucubitan, Pocotpocot, Curagda. Sp.-Fil., Pepinillo de San Gregorio.
- Cucurbita lagenaria-oblonga Blanco, ed. 1, 772; ed. 2, 531 = Lagenaria vulgaris Seringe, var.! T., Tabayag.
- Cucurbita lagenaria-villosa Blanco, ed. 1, 772; ed. 2, 532 = Lagenaria vulgaris Seringe, var.! T., Opo.
- Mormordica operculata Blanco, ed. 1, 770; ed. 2, 530, non Linn. = Luffa cylindrica Roem.! T., Salag salag.
- Cucumis acutangulus Linn., Blanco, ed. 1, 776; ed. 2, 534 = Luffa acutangula (Linn.) Roxb.! T., Patola.
- Cucurbita pepo-aspera Blanco, ed. 1, 773; ed. 2, 532 = Benincasa cerifera Savi.! T., Condol.
- Momordica cylindrica Blanco, ed. 1, 769; ed. 2, 530, non Linn. = Momordica charantia Linn.! Native names the same as for the next species.
- Momordica balsamina Linn., Blanco, ed. 1, 768; ed. 2, 529.! T., Ampalaya, Apalaya, Amargoso. B., Palla, Margoso. P., Apalia. I., Paria.
- Momordica sphæroidea Blanco, ed. 1, 771; ed. 2, 531 = Momordica cochinchinensis Spreng.! T., Boyocboyoc.

- Cucumis melo Linn., Blanco, ed. 1, 775; ed. 2, 534. Reduced by Villar to Cucumis trigonus Roxb., but it is evident that Blanco correctly interpreted the species, and that the plant he describes is only a form of Cucumis melo Linn. T., Tabogo.
- Cucurbita sulcata Blanco, ed. 1, 773; ed. 2, 532 = Cucurbita maxima Duchesne.! T., Calabazang bilog.
- Cucumis luzonicus Blanco, ed. 1, 861; ed. 2, 534. Reduced by Villar to Melothria indica Lour., which is probably correct. T., Melon daga.

BEGONIACEÆ.

Begonia capensis Blanco, ed. 1, 724; ed. 2, 501, non Linn. Reduced by Villar to Begonia rhombicarpa A. DC., which is probably correct. T., Pingol bato.

CACTACEÆ.

- Cactus pitajaya Blanco, ed. 2, 289, non Linn. Reduced by Villar to Cereus triangularis Haw., which is probably correct. Cultivated only.
- Cactus opuntia Blanco, ed. 1, 414; ed. 2, 288, non Linn. Reduced by Villar to Opuntia cochinillifera Mill., which is probably correct. T., Sandoc sandoc, Dilang baca.

FICOIDEÆ.

- Sesuvium portulacastrum Linn., Blanco, ed. 1, 426; ed. 2, 297.! T., Dampalit, Tarumpalit. V., Bilang bilang, Dampalit. P., Dampalit, Carampalit.
- Portulaca toston Blanco, ed. 1, 408; Portulaca axilliflora Blanco, ed. 2, 285 = Trianthema monogyna Linn.! T., Toston.
- Glinus lotoides Linn., Blanco, ed. 1, 413; ed. 2, 288.! T., Lobio.
- Mollugo subserrata Blanco, ed. 1, 51; ed. 2, 34. The same as the following species. T., Malagoso, Molugoso.
- Mollugo stricta Linn., Blanco, ed. 1, 52; ed. 2, 35.! T., Malagoso, Mollugoso.

UMBELLIFERÆ.

- Hydrocotyle asiatica Linn., Blanco, ed. 1, 212; ed. 2, 149.! T., Taquip suso, Taquip cohol.
- Ammi glaucifolium Blanco, ed. 1, 213, non Linn.; Daucus anisodorus Blanco, ed. 2, 150. Reduced by Villar to Carum copticum Benth., which is probably correct. T., Lamudio, Damoro. B., Lamudio. P., Damoro.
- Anethum foeniculum Linn., Blanco, ed. 1, 214; ed. 2, 150 = Foeniculum vulgare Mill.! T., Haras.

ARALIACEÆ.

- * Aralia bipinnata Blanco, ed. 1, 222; ed. 2, 157. Reduced by Villar to Aralia javanica Thunb., a species not known from the Philippines. A shrub, flowering in February. T., Potat.
- Aralia tripinnata Blanco, ed. 1, 223; Panax fruticosa Blanco, ed. 2, 156 = Panax fruticosum Linn.! T., P., Papua. B., Macan, Papua.
- Aralia pendula Blanco, ed. 1, 223; ed. 2, 157 = Polyscias nodosa Seem.! T., B., Bongliu, Bingliu, Biasbias, Malapapaya.
- * Polyscias digitata Blanco, ed. 1, 224; Aralia? octophylla Blanco, ed. 2, 158, non Lour. Reduced by Villar to Heptapleurum rigidum Seem., a species not known from the Philippines. Blanco's specimens were from Cebu, flowering in September. B., Tagima.
- Polyscias odorata Blanco, ed. 1, 225; Paratropia crassa Blanco, ed. 2, 158 = Heptapleurum venulosum Seem.! T., Limalima, Galamai amo. B., Cayangcan, Taguima, Calangcang, Carangcang, Limalima, Taguilima, Tuglima. P., Limalima.
- Polyscias obtusa Blanco, ed. 1, 226; Paratropia obtusa Blanco, ed. 2, 226. Reduced by Villar to the preceding species, which is certainly correct.!
- * Nauclea digitata Blanco, ed. 2, 102. This species is reduced by Villar to Heptapleurum cephalotes Clarke, a species not known from the Philippines. Very doubtful.! Blanco's specimens were from Angat, Province of Bulacan, native name not given.
- * Polyscias disperma Blanco, ed. 1, 226. Reduced by Villar to Arthrophyllum diversifolium Blume, which is certainly an error. Blanco's specimens were from the seashore, Batangas. This species is excluded from the second edition.
- * Polyozus bipinnatus Blanco, ed. 2, 43. Reduced by Villar to Arthrophyllum pinnatum Clarke, a species not known from the Philippines. Blanco's specimens were from Angat, Province of Bulacan, flowering in June.

CORNACEÆ.

- Alangium octopetalum Blanco, ed. 2, 310 = Alangium lamarckii Thwaites.!
- Guettarda jazminifiora Blanco, ed. 1, 722; Guettarda speciosa Blanco, ed. 2, 499, non Linn. Reduced by Villar to Marlea begoniæfolia Roxb., which is probably correct. T., Calumpangin, Bagaolan.

CAPRIFOLIACEÆ.

Sambucus javanica Blume, Blanco, ed. 2, 151.!

RUBIACEÆ.

- Nauclea lutea Blanco, ed. 1, 141; Nauclea glaberrima Blanco, ed. 2, 100, non Bartl. = Sarcocephalus cordatus Miq.! Erroneously reduced by Villar to Sarcocephalus subditus Miq. T., Bangcal. V., Bancal, Cabag, Hanbabalos. P., Bancal. I., Bulala.
- Nauclea glandulosa Blanco, ed. 1, 143; Nauclea glabra Blanco, ed. 2, 101 = Nauclea blancoi Vidal.! Erroneously reduced by Villar to Anthocephalus cadamba Miq. T., Bagarilat.
- Nauclea adina Blanco, ed. 2, 102, non Smith = Stephegyne diversifolia Hook. f.! This species is described in the first edition under the native name only, Mambog. Erroneously reduced by Villar to Stephegyne parvifolia Korth., a species that does not extend to the Philippines. T., Mambog.
- * Mamboga capitata Blanco, ed. 1, 140; Nauciea Iuzoniensis Blanco, ed. 2, 102. Reduced by Villar to Stephegyne speciosa Korth., which is probably an error. Probably a Nauciea or a Sarcocephalus. A very large tree. T., Mambog.
- * Nauclea latifolia Blanco, ed. 1, 144; Nauclea obtusa Blanco, ed. 2, 101, non? Blume. Villar considers that Blanco was correct in reducing his species to that of Blume, but Nauclea obtusa Blume, has not as yet been found in the Philippines. A tree flowering in August. T., Balod.
- * Tapogomea rubra Blanco, ed. 1, 145; Cephaelis expaleacea Blanco, ed. 2, 103. Reduced by Villar to *Uncaria acida* Roxb., a species not known from the Philippines. Probably, however, a species of *Uncaria*. Blanco's specimens were from Cebu, flowering in April. B., *Mampol*.
- * Nauclea lanceolata Blanco, ed. 1, 144; Nauclea calycina Blanco, ed. 2, 101, non? Bartl. Reduced by Villar to Nauclea purpurea Roxb., a species not known from the Philippines. T., Bagarilao na itim.
- Exostemma philippicum Blanco, ed. 2, 113, non R. et S. = Hymenodictyon excelsum Wall.! T., Huliganga.
- Rondeletia asiatica Blanco, ed. 1, 146, non Linn.; Wendlandia exserta Blanco, ed. 2, 104, non DC. = Wendlandia luzonensis DC. Erroneously reduced by Villar to Wendlandia paniculata DC.
- Dentella repens Forst., Blanco, ed. 1, 146; ed. 2, 103.! T., Dilang butiqui.
- Oldenlandia capensis Blanco, ed. 1, 62; ed. 2, 45, non Thunb. = Oldenlandia diffusa Roxb.!
- Oldenlandia paniculata Linn., Blanco, ed. 1, 61; ed. 2, 44.!
- Oldenlandia affinis Blanco, ed. 2, 44, non R. et S. Reduced by Villar to Oldenlandia nudicaulis Roth., which is certainly an error, although the latter species has been found in the Philippines. Probably a form of Oldenlandia corymbosa Linn.

- Ophiorrhiza oblongifolia DC., Blanco, ed. 2, 64.!
- Mussaenda frondosa Blanco, ed. 1, 167; ed. 2, 118, non Linn. = Mussaenda grandiflora (Meyen.) Rolfe.! T., Tingatinga. I., Balailamoc.
- * Stigmanthus cymosus Blanco, ed. 2, 117. Reduced by Villar to Webera odorata Roxb., which is certainly an error, as that species does not extend to the Philippines. If a species of Webera, probably the same as W. luzonensis Vidal. Blanco's specimens were from Calauan, Province of Laguna, native name not given.
- * Randia aculeata Blanco, ed. 1; 141; ed. 2, 99. Reduced by Villar to Gardenia dumetorum Lam., which is probably an error, as the latter species is unknown from the Philippines. Blanco's specimens were from Bauang, Province of Batangas, a shrub growing on the seashore, flowering in May. T., Sinampaga.
- * Remijia odorata Blanco, ed. 2, 115. Reduced by Villar to Randia densiftora Benth., which may be correct, although the latter species is not known from the Philippines at the present time. Blanco's specimens were from Bauang, Province of Batangas, native name not given.
- * Serissa myrtifolia Blanco, ed. 1, 164; Remijia angatensis Blanco, ed. 2, 115. Transferred to Randia by Villar, as a distinct species Randia angatensis (Blanco) F. Vill. A shrub, known only from Blanco's description, flowering in August, Blanco's specimens being from Angat, Province of Bulacan. T., Caragli.
- Serissa pinnata Blanco, ed. 1, 163; Remijia obscura Blanco, ed. 2, 116 = Gardenia pinnata (Blanco) (Gardenia obscura Vidal; Randia obscura F.-Vill.). T., Caragli. P., Pagbut.
- Sulipa pseudopsidium Blanco, ed. 1, 497; ed. 2, 347 = Gardenia pseudopsidium (Blanco) F.-Vill.! T., Calapi, Sulipa, Malabayabas.
- Ixora manila Blanco, ed. 1, 60; ed. 2, 42 = Scyphiphora hydrophyllacea
 Gaertn.! T., Nilad, Nilar.
- * Ixora glandulosa Blanco, ed. 1, 61; ed. 2, 42. Reduced by Villar to Canthium confertum Korth, which is probably an error, as that species is unknown from the Philippines. T., Malapatopat.
- Ronabea bipinnata Blanco, ed. 1, 162; Ronabea arborea Blanco, ed. 2, 114 = Canthium bipinnatum (Blanco) (Canthium arboreum Vidal.). This species was erroneously referred to Canthium mite Bartl., by Villar. T., Tadiang anunang. I., Pamalatanguen.
- Canthium pauciflorum Blanco, ed. 1, 165; Canthium horridum Blume, Blanco, ed. 2, 116. The latter name is correct. T., Suliac daga.
- Canthium monoflorum Blanco, ed. 1, 166; Canthium pedunculare Cav., Blanco, ed. 2, 116. The latter name is correct.
- * Vangueria stellata Blanco, ed. 1, 167; ed. 2, 117. Reduced by Villar to Vangueria spinosa Roxb., which is probably an error, as Blanco's species does not appear to belong to this genus. A tree flowering in August. T., Malaas-is, Madondon.

- Ixora coccinea Linn., Blanco, ed. 1, 59; ed. 2, 41.! Villar reduces this species to Ixora stricta Roxb., but Blanco describes the common cultivated form, which is apparently not a native of the Philippines. T., Santan, Santa Ana.
- * ixora arborea Blanco, ed. 1, 61; ed. 2, 42. Reduced by Villar to Ixora stricta Roxb., which is certainly an error. A tree flowering in August. T., As-as.
- * Taligalea umbellata Blanco, ed. 2, 337. Reduced by Villar to Ixora macrohylla Bartl., which must be considered doubtful. Blanco's specimens were from Cebu, flowering in March. V., Bagobago.
- * Pavetta membranacea Blanco, ed. 1, 59; Pavetta sambucina Blanco, ed. 2, 41, non DC. Reduced by Villar to Pavetta angustifolia R. et S., which is probably an error. Blanco's species may be a form of Pavetta indica Linn., or a distinct species. His specimens were from Angat, flowering in September.
- Coffea arabica Linn., Blanco, ed. 1, 156; ed. 2, 110.! Sp.-Fil., Café.
- Morinda litoralis Blanco, ed. 2, 109 = Morinda citrifolia Linn.!
- Morinda citrifolia Blanco, ed. 1, 149; Morinda ligulata Blanco, ed. 2,
 105 = Morinda bracteata Roxb.! T., V., Bancudo, Pancudo, Bangcoro,
 Nino, Culit, Tumbong aso, Lino, Mambog, Tacpus. P., Taliantar.
 I., Apatot.
- Morinda royoc Blanco, ed. 1, 148; ed. 2, 105, non Linn. = Morinda tinctoria Roxb.! T., Tumbong asong hapay.
- Morinda umbellata Blanco, ed. 2, 110. Reduced by Villar to Morinda microcephala Bartl., but it is evident that Blanco correctly interpreted the Linnean species.
- * Coffea volubilis Blanco, ed. 1, 157; ed. 2, 111. Reduced by Villar to *Morinda tinctoria* Roxb., which may be correct. A scandent shrub, flowering in August. T., *Halon*.
- Pæderia tacpo Blanco, ed. 1, 160; ed. 2, 113 = Psychotria tacpo (Blanco) Rolfe.! Erroneously reduced by Villar to Psychotria malayana Jack. T., Tacpo.
 - Pæderia fœtida Blanco, ed. 1, 159; ed. 2, 112, non Linn. = Paederia tomentosa Blume.! T., Cantotai. B., Lilitan. P., Cantotai, Dicuta maboloc, Mabatang dicut.
 - Spermacoce mutilata Blanco, ed. 2, 43 = Spermacoce hispida Linn.!
 - Spermacoce muriculata Blanco, ed. 2, 44. This species is reduced by Villar to Spermacoce scaberrima Blume, probably a form of the preceding species. Blanco's specimens were from Guadalupe, near Manila.

COMPOSITÆ.

- Serratula multiflora Blanco, ed. 1, 617; ed. 2, 431, non Linn. = Vernonia chinensis Less.! T., Bayaquiboc.
- Elephantopus scaber Linn., Blanco, ed. 1, 634; ed. 2, 441.!

- Elephantopus serratus Blanco, ed. 1, 635; ed. 2, 442. Reduced by Villar to *Elephantopus mollis* HBK., which is probably correct.
- Ageratum quadriflorum Blanco, ed. 1, 624; Elephantopus? dubius Blanco, ed. 2, 442 = Elephantopus spicatus B. Juss.! T., Dilang usa, Tabatabacohan, Sigang dagat. P., Sumag. B., Habul, Ardotag. I., Cabcabron.
- Eupatorium ayapana Vent., Blanco., ed. 1, 619; ed. 2, 432.! Sp.-Fil., Ayapana.
- Knautia sagittata Blanco, ed. 1, 54; ed. 2, 36 = Mikania scandens Willd.!
 Perdicum tomentosum Blanco, ed. 1, 630; ed. 2, 439 = Grangea maderaspatana Poir.!
- * Baccharia ivæfolia Blanco, ed. 1, 627; ed. 2, 437, non Linn. Reduced by Villar to Conyza viscidula Wall., which may be correct. B., Tagabili.
- * Conyza gouani Blanco, ed. 1, 629, non Wild.; Conyza erosa Blanco, ed. 2, 439. Reduced by Villar to Blumea manillensis DC., which may be correct.
- Conyza balsamifera Linn., Blanco, ed. 1, 628; ed. 2, 438 = Blumea balsamifera DC.! T., Sambon. V., Lacdanbulan, Hamlibon, Lalacdan, Lacad bulan, Guintenguinten, Gabuen, Ayoban, Alibun. P., Sambon. I., Sobsob.
- Baccharis indica Linn., Blanco, ed. 1, 627; ed. 2, 438 = Pluchea indica (Linn.) Less.!
- Sphæranthus alatus Blanco, ed. 1, 635.; Sphæranthus indicus Blanco, ed. 2, 442, non Linn. = Sphæranthus africanus L.! T., Sambong gala.
- Sphæranthus elongatus Blanco, ed. 1, 636; ed. 2, 443 = Pterocaulon cylindrostachyum Clarke.! T., Sambong gala.
- Xeranthemum stæhelina Blanco, ed. 1, 629; non Don.; Gnaphalium dichotomum Blanco, ed. 2, 439, non Wild = Gnaphalium luteo-album Linn.! Reduced by Villar to Gnaphalium indicum Linn. Blanco's specimens were from Agoo, Province of Union. I., Badoc.
- * Conyza dentata Blanco, ed. 1, 629, non Linn.; Conyza cappa Blanco, ed. 2, 438, non Ham. Reduced by Villar to *Inula cappa* DC., which is certainly an error. Possibly a species of *Blumea*.
- Anthemis cotula Blanco, ed. 1, 633, non Linn.; Artemesia viridis Blanco, ed. 2, 436, non Linn. = Eclipta alba Hassk.! T., Higuis manoc. I., Tintatinta.
- Spilanthes acmella Blanco, ed. 1, 420; ed. 2, 433, non DC. = Wedelia biflora DC.! Villar considers that Blanco correctly interpreted DeCandolle's species, but his description certainly applies to Wedelia biflora DC. T., Hagonoi. V., Agonoi. P., Palunai.
- Spilanthes lobata Blanco, ed. 1, 622; ed. 2, 434 = Spilanthes acmella Linn.! T., Hagonoi.
- Spilanthes peregrina Blanco, ed. 1, 622; ed. 2, 434. Certainly only a form of Wedelia biflora DC.! T., Hagonoi.

- Coreopsis gracilis Blanco, ed. 2, 591 = Cosmos caudatus HBK.!
- Bidens bipinnata Blanco, ed. 1, 623; ed. 2, 435 = Bidens pilosa Linn.!
- Tagetes patula Linn., Blanco, ed. 1, 632; ed. 2, 440.! Cultivated only. Sp.-Fil., Amarillo.
- Matricaria chamomilla Blanco, ed. 1, 631; ed. 2, 440, non Linn. = Chrysanthemum indicum Linn.! Sp.-Fil., Manzanilla.
- Cotula quinqueloba Blanco, ed. 1, 626; ed. 2, 436 = Centripeda orbicularis Lour.! T., V., Harangan. V., Pissic.
- Artemesia vulgaris Linn., Blanco, ed. 1, 625; ed. 2, 435. The identification of this species seems to be correct. T., Ca Maria, Santa Maria, Tinisas.
- *Cacalia sarrasenica Blanco, ed. 1, 618; non Linn.; Senecio cacaliaster Blanco, ed. 2, 441, non Lam. Reduced by Villar to Gynura angulosa DC., a species that is not known from the Philippines. Flowering in August. T., Sampaga del monte.
- Cacalia sonchifolia Linn., Blanco, ed. 1, 618; Emilia sonchifolia (Linn.) DC. Blanco, ed. 2, 432. The second name is correct. T., Tagolinao. P., Tagulinao. B., Libun.
- Carthamus tinctorius Linn., Blanco, ed. 1, 616; ed. 2, 431.! T., Biri, Casubha, Lago. V., Casabha. P., Casubha, Cachumba.

GOODENIACEÆ.

Scaevola lobelia Blanco, ed. 1, 147; ed. 2, 104, non Murr. = Scaevola koenigii Vahl.! T., V., Boto, Bosboron, Mosboron, Bocaboc, Panabolong, Pangangtolong. Z., Linog.

CAMPANULACEÆ.

- Pongatium spongiosum Blanco, ed. 1, 86; Sphenoclea zeylanica Gaertn., Blanco, ed. 2, 62. The latter name is correct.! T., Silisilihan.
- Reichelia palustris Blanco, ed. 1, 220; ed. 2, 155. The same as the preceding.!

ERICACEÆ.

Clethra alnifolia Blanco, ed. 2, 259, non Linn. = Clethra lancifolia Turcz.! T., Malaclac.

PLUMBAGINACEÆ.

Plumbago viscosa Blanco, ed. 1, 78; ed. 2, 58 = Plumbago zeylanica Linn.! T., Sangdigquit. I., Bangbang, Talancao.

MYRSINACEÆ.

Bassovia sylvatica Blanco, ed. 2, 95. Reduced by Villar to Maesa indica A. DC., a species that does not extend to the Philippines. Probably the same as the species described by Mez as Maesa lawa. Blanco's specimens were from Angat, a small shrub, flowering in February. T., Solinao.

- Maesa membranacea Blanco, ed. 2, 590, non A. DC. = Maesa cumingiana Mez.!
- Bladhia japonica Blanco, ed. 1, 126; ed. 2, 90, non Thunb. Reduced by Villar to Ardisia pyramidalis Pers. = Ardisia serrata (Cav.) Pers. Questionably referred here by Mez, but Villar is probably correct. Flowering in August. T., Tacpong dalaga.
- Willughbeia drupacea Blanco, ed. 1, 132; ed. 2, 94. This species is reduced by Villar to *Ardisia obovata* Blume, which is a synonym of *Ardisia humilis* Vahl., certainly correctly referred.!
- Aegiceras corniculatum (Linn.) Blanco, ed. 1, 79; ed. 2, 59. This name is correct, Aegiceras majus Gaertn., is a synonym. T., Tingang baguis, Pipisic.

SAPOTACEÆ.

- Achras lucuma Blanco, ed. 1, 237; ed. 2, 166, non Ruiz et Pav. = Lucuma mammosa Gaertn.! T., Mamei.
- Sideroxylon duclitan Blanco, ed. 1, 129; ed. 2, 92. A distinct species, flowering in September. T., Duclitan, Dulitan.
- *Sideroxylon balitbitan Blanco, ed. 1, 130; ed. 2, 92. Probably a distinct species, but certainly closely related to the preceding. T., Balitbitan.
- Achras sapota Linn., Blanco, ed. 1, 236; ed. 2, 165.! Sp.-Fil., Chicos.
- Palaquium lanceolatum Blanco, ed. 1, 403; ed. 2, 282.! A distinct species. T., Bagalangit.
- Palaquium latifolium Blanco, ed. 1, 404; ed. 2, 282.! A distinct species. T., Palac palac.
- Palaquium oleiferum Blanco, ed. 1, 405; ed. 2, 283.! A species closely related to the preceding. I., Daracan.
- Azaola betis Blanco, ed. 1, 402; ed. 2, 281 = Illipe betis (Blanco) Merrill (Payena betis F.-Vill.). T., P., Betis.

EBENACEÆ.

- Diospyros pilosanthera Blanco, ed. 1, 304; ed. 2, 211. A good species. T., Bolong eta. B., Dalondong, Amaga.
- Diospyros multiflora Blanco, ed. 1, 303, non Wall.; Diospyros lotus Blanco, ed. 2, 210, non Linn. = Diospyros canomoi A. DC.! A distinct species. T., Canomoi, Canomai.
- * Diospyros bifiora Blanco, ed. 1, 303; ed. 2, 210. Possibly a good species, at present known only from Blanco's description. A tree flowering in June. T., Talang.
- * Diospyros kaki Blanco, ed. 2, 211, non? L. f. Possibly this may be the same as the species described under this name by Linnaeus, f., and Villar so considers it. Blanco's specimens were from Mahaihai, Province of Laguna, known there as Pagatpat. Blanco had only fruits. Diospyros kaki of the seond edition is entirely different from D. kaki of the first edition.

- Sapota nigra Blanco, ed. 1, 409; Diospyros nigra Blanco, ed. 2, 211 = Diospyros ebenaster Retz.! Sp.-Fil., Zapote negro.
- Diospyros kaki Blanco, ed. 1, 302, non Linn.; Diospyros embryopteris Blanco, ed. 2, 209, non Pers. = Diospyros philippensis (Desr.) Gurke! (D. discolor Willd.). T., Talang, Mabolo. V., Amaga. P., Talang.

STYRACEÆ.

Guettarda polyandra Blanco, ed. 2, 500 = Symplocos polyandra (Blanco?)

A. Brand. Villar erroneously refers this species to Symplocos racemosa Roxb.

OLEACEÆ.

- Nyctantes sambac Linn., Blanco, ed. 1, 9; ed. 2, 6 = Jasminum sambac (Linn.) Ait. T., Sampaga. V., Capopot visaya, Manul. P., Campopot, Sampagang pougso, Culatai.
- * Mogorium aculeatum Blanco, ed. 1, 9; ed. 2, 7. Reduced by Villar to Jasminum marianum DC., but this species is not known from the Philippines. Possibly a distinct species of Jasminum. Flowering in May.

SALVADORACEÆ.

Azima nova Blanco, ed. 1, 68; ed. 2, 49. Considered by Vidal as a distinct species, although reduced by Villar to Azima tetracantha Lam. It may be the same as Azima sarmentosa Blume.

APOCYNACEÆ:

- Allamanda cathartica Linn., Blanco, ed. 2, 64.!
- Brabejum? lucidum Blanco, ed. 2, 40 = GYNOPOGON MONILIFERA (Vid.).
 (Alyxia monilifera Vid.). Erroneously reduced by Villar to Alyxia stellatum R. et S. Blanco's specimens were from the pueblo of Bolohan, Cebu. V., Layo.
- * Brabejum? concatenatum Blanco, ed. 2, 40. Reduced by Villar to Alyxia laurina Gaud., a species not known from the Philippines. Blanco's specimens were from Cebu. Probably a distinct species of Gynopogon.
- * Brabejum ? pinnatum Blanco, ed. 2, 40. Reduced by Villar to Alyxia odorata Wall., which is probably an error. Blanco's specimens were from Cebu. Probably a distinct species of Gynopogon. V., Layo.
- Gerbera thevetla Linn., Blanco, ed. 1, 125; ed. 2, 89 = Thevetia neriifolia Juss.! Introduced from America. Sp.-Fil., Campanelo.
- Cerbera manghas Blume, Blanco, ed. 1, 125; ed. 2, 89 = Cerbera odollam Gaertn.! T., Toctoc calo.
- * Elcana seminuda Blanco, ed. 2, 584. Reduced by Villar to Cerbera lactaria Ham., a species that does not extend to the Philippines. Possibly a species of Cerbera. Blanco's specimens were from Angat, Province of Bulacan.
- Vinca rosea Linn., Blanco, ed. 1, 116; ed. 2, 84.! T., Cantotan.

- Plumeria alba Blanco, ed. 1, 111; ed. 2, 80, non Linn. = Plumeria acutifolia Poir. T., Carachucha, Calachuchi, Calasasi.
- Echites scholaris Linn., Blanco, ed. 1, 106; ed. 2, 77 = Alstonia scholaris (Linn.) R. Br.! T., Dita. I., Dallopaven.
- Echites trifida Blanco, ed. 1, 109; ed. 2, 79, non Jacq. Reduced by Villar to Alstonia spectabilis Miq., but it is certainly Alstonia macrophylla Wall.! T., Cayaocayao, Batino.
- Alstonia batino Blanco, ed. 2, 589 = Alstonia macrophylla Wall.! T., Batino.
- Tabernaemontana globosa Blanco, ed. 1, 116; ed. 2, 83 = Voacanga cumingiana Rolfe (V. foetida F.-Vill., non Blume). A small tree flowering in August. T., Bayag usa.
- Tabernaemontana laurifolia Blanco, ed. 1, 114; ed. 2, 82, non Linn. =

 Tabernaemontana pandacaqui Poir.! T., Campopot, Pandacaqui.

 B., Alibutbut, Toar, Pandaya. P., Pandacaqui. I., Curribuetbuet.
- *Tabernaemontana polygama Blanco, ed. 2, 82. Considered by Villar to be a distinct species. Blanco's specimens were from Parañaque and Mandoloyon, near Manila, flowering in May.
- Echites repens Blanco, ed. 1, 109, non Jaeq.; Echites procumbens Blanco, ed. 2, 78 = HOLARRHENA PROCUMBENS (Blanco). Erroneously reduced by Villar to Holarrhena macrocarpa (Hassk.) F.-Vill. See Nos. 1844 and 3299 Merrill, and No. 298 Whitford. T., Malayantoc, Hinguiu.
- Echites spiralis Blanco, ed. 1, 110; ed. 2, $79 = Parsonsia \ rheedii \ F. Vill.!$ This species is figured in the third edition of the Flora de Filipinas, t. 310, also by Vidal, Sinopsis, Atlas, t. 66. f. E., while a full description is given by Vidal, Revision, 184.
- Anasser laniti Blanco, ed. 1, 112; ed. 2, 81 = WRIGHTIA LANITI (Blanco) (Wrightia ovata A. DC.!). Blanco's name is the earliest for this species and should be retained. T., Laniti.
- Nerium oleander Blanco, ed. 1, 104; ed. 2, 75, non Linn. = Nerium odorum Soland.! Sp. Fil., Adelfa. T., Baladre.
- Echites caudata Blanco, ed. 1, 106; ed. 2, 77, non Linn. = Ichnocarpus navesii Rolfe.! Erroneously reduced by Villar to Ichnocarpus frutescens R. Br. T., Hinguio.
- Tabernaemontana elliptica Blanco, ed. 1, 116; ed. 2, 83, non Thunb. = Choneomorpha macrophylla Wall.! Blanco's specimens were from Tala, Province of Rizal or Bulacan, a scandent shrub, flowering in July.

ASCLEPIADACEÆ.

*Tabernaemontana cirrhosa Blanco, ed. 1, 115; ed. 2, 83. Reduced by Villar to Finlaysonia obovata Wall., which is certainly an error. Blanco's specimens were from Bauang, Province of Batangas, a scandent shrub growing along the seashore, in situations subject to inundation.

- * Apocynum mucronatum Blanco, ed. 1, 852; ed. 2, 143. Reduced by Villar to *Toxocarpos gracilis* Decne., which is very doubtful. A vine flowering in May.
- Asclepias gigantea Willd., Blanco, ed. 1, 207; ed. 2, 146 = Calotropis gigantea R. Br.! T., Capalcapal.
- Asclepias syriaca Blanco, ed. 1, 204; ed. 2, 144, non Linn. = Asclepias curassavica Linn.! T., Bulac castila.
- *Cynanchum viminale Blanco, ed. 1, 203; ed. 2, 143, non? Linn. Villar considers that Blanco describes the Linnean species, but this is probably an error, unless Blanco described an introduced plant. Blanco's species may be referable to Sarcostemma brevistigma W. et A. A vine, flowering in February, known in Manila as Mil leguas.
- * Asclepias peregrina Blanco, ed. 1, 207; ed. 2, 146. Reduced by Villar to Sarcolobus carinatus Wall., which is certainly an error. Blanco's specimens were from Bauang, Province of Batangas, growing in situations subject to inundation, along the seashore.
- Asclepias dæmia Blanco, ed. 1, 208; ed. 2, 146, non Forsk. Reduced by Villar to Gymnema tingens W. et A., which is probably correct.
- Cynanchium tenellum Blanco, ed. 1, 204; ed. 2, 143, non Linn. Reduced by Villar to Tylophora perrottetiana Decne., which is probably correct.
- Marsdenia akkar Blanco, ed. 1, 118; ed. 2, 85. Reduced by Villar to Marsdenia tinctoria R. Br., which is probably correct. B., Payanguit, Aringit.
- * Marsdenia tagudina Blanco, ed. 1, 121; ed. 2, 86. Considered by Villar to be a distinct species. Blanco's specimens were from the pueblo of Taguding, Province of Ilocos Sur. I., Tayomtayom.
- * Pergularia glandulosa Blanco, ed. 1, 202; ed. 2, 141. Reduced by Villar to *Pergularia minor* Andr., which is probably an error. Blanco's specimens were from Bauang, Province of Batangas, flowering in July.
- * Pergularia procumbens Blanco, ed. 1, 204; Pergularia glabra Blanco, ed. 2, 141. Considered by Villar to be a distinct species, possibly however a synonym of *Pergularia odoratissima* Sm. Blanco's specimens were from Manila. T., *Mil leguas*.
- Stapelia quadrangula Blanco, ed. 1, 202; ed. 2, 142, non Forsk. = Heterostemma cuspidatum Decne.! T., Biniguasan.
- Cynanchium hirtum Blanco, ed. 1, 203; ed. 2, 143, non Linn. Reduced by Villar to *Dregea viridiflora* (DC.) F.-Vill., which is probably correct. Blanco's specimens were from Malinta, near Manila.
- * Stapelia meliflua Blanco, ed. 1, 202; ed. 2, 142. Reduced by Villar to Hoya diversifolia Blume, a species not known from the Philippines. Certainly a species of Hoya, but not that species. A vine, flowering in June. T., Baliobalic.
- * Hoya carnosa Blanco, non? R. Br., ed. 2, 142. Identification possibly correct. Blanco's specimens were from cultivated plants.

- * Marsdenia parasita Blanco, ed. 1, 120; ed. 2, 86. Reduced by Villar to Hoya parasitica Wall., which is certainly an error. Certainly, however, a species of Hoya.! Blanco's specimens were epiphytes growing on Erythrina and on Mangifera, flowering in May, fruiting in June.
- Asclepias carnosa Blanco, ed. 1, 208; ed. 2, 147 = Hoya multiflora Blume.!

LOGANIACEÆ.

- Buddleia virgata Blanco, ed. 1, 57; ed. 2, 38, non Linn. Reduced by Villar to Buddleia neemda Hamilt., but certainly is Buddleia asiatica Lour.! T., Talicnono. I., Tugnang. B., Lagundi salasa.
- Fagraea scholaris Blanco, ed. 2, 93. Reduced by Villar to Fagraea cordifolia Blume, a species not known from the Philippines. Probably the same as Fagraea morindaefolia Blume. T., Cabal.
- * Tayotum nigrescens Blanco, ed. 1, 105; ed. 2, 76. Reduced by Villar to Norrisia malaccensis Gardn., which is undoubtedly an error. Blanco's specimens were from Angat, Province of Bulacan, a shrub flowering in June. T., Saguing saguing, Tayoto.
- Ignatia amara Linn. f., Blanco, ed. 1, 82; Strychnos philippensis Blanco, ed. 2, 61 = Strychnos ignatii Berg.! Retained under the second name by Villar. For full synonomy see Hooker's Icones, Vol. 25: pl. 2212, 2213. T., Pepita, Fruta. P., Pepita sa catbalogan. V., Pangaguason, Aguason, Canlara, Mana-naog, Dancagi, Catalonga, Igasud.

GENTIANACEÆ.

- Exacum albens Blanco, ed. 1, 58; ed. 2, 39, non Linn. f. Reduced by Villar to Exacum chironioides Griseb., probably, however, the same as Exacum tetragonum Roxb.
- Cobamba dichotoma Blanco, ed. 1, 510; ed. 2, 355. Reduced by Villar to Canscoria diffusa R. Br., which is certainly correct.! T., Codamba.
- * Codamba blancoi Azaola, in Blanco, ed. 2, 591. Reduced by Villar to Canscoria decussata R. et S., which is certainly an error. The material on which this species was based was from Gumai, Province of Pampanga. A form of the preceding species.?
- Menyanthes indica Bory, Blanco, ed. 1, 87; ed. 2, 63 = Limnanthemum cristatum Griseb.! T., Lauas. I., Locloquisin.

HYDROPHYLLACEÆ.

Hydrolea arayatensis Blanco, ed. 1, 211; Hydrolea zeylanica Vahl., Blanco, ed. 2, 148. The second name is correct.

BORAGINACEÆ.

Cordia banalo Blanco, ed. 1, 124; Cordia ? ignota Blanco, ed. 2, 88 = Cordia subcordata Lam.! T., Banalo.

- Cordia sebestena Blanco, ed. 1, 121; ed. 2, 87, non Linn. = Cordia dichotoma Blanco, ed. 1, 123; ed. 2, 88, non Forst. = Cordia blancoi Vidal.! Erroneously referred to Cordia myxa by Villar. T., Anonang. V. Anonang, Bilibi.
- Ehretia beurreria Blanco, ed. 1, 127; ed. 2, 91, non Linn. = Ehretia philippinensis A. DC.!
- Ehretia virgata Blanco, ed. 1, 127; ed. 2, 165, non Swartz. = Ehretia onava A. DC.! T., Onava.
- Menais mollis Blanco, ed. 1, 139; ed. 2, 99 = Ehretia navesii Vidal (Ehretia mollis Merrill). This species was erroneously referred by Villar to the preceding species. T., Taglocot.
- Carmonia heterophylla Cav., Blanco, ed. 1, 209; ed. 2, 147 = Ehretia buxifolia Roxb.! T., B., Alanguit, Alangitngit.
- Coldenia procumbens Linn., Blanco ed. 1, 74; ed. 2, 56.! T., Tapiasin, I., Tabtabocol.
- Tournefourtla arborea Blanco, ed. 1, 129; ed. 2, 91 = Tournefourtia argentea Linn. f.!
- Tournefourtia hirsutissima Blanco, ed. 1, 128; ed. 2, 91, non Linn = Tournefourtia sarmentosa Linn.! V., Patai ood. I., Sallacapo.
- Heliotropum parviflorum Blanco, ed. 1, 80; ed. 2, 59 = Heliotropum indicum Linn.! T., Hinlaylayon, Cotingcotingan.
- Borago Indica Linn., Blanco, ed. 2, 60. Reduced by Villar to *Trichodesma* indicum R. Br., which is probably correct. Blanco's specimens were from Paranaque, near Manila.
- Borago indica Blanco, ed, 1, 81, non Linn.; Borago africana? Blanco, ed. 2, 60 = Trichodesma zeylanicum R. Br.! T., Sigang dagat, Dilan usa.

CONVOLVULACEÆ.

- Convolvulus muricatus Blanco, ed. 1, 92; ed. 2, 68, non Linn. = Ipomæa bona-now Linn.! T., Camo camotihan.
- Ipomoea quamoclit Linn., Blanco, ed. 1, 97; ed. 2, 72 = Quamoclit vulgaris Choisy.! T., Agoho.
- Convolvulus nil Linn., Blanco, ed. 1, 92; ed. 2, 68 = Ipomæa nil (Linn.), Roth.! Erroneously reduced by Villar to Ipomæa hederacea Jacq.
- Convolvulus repens Blanco, ed. 1, 92; ed. 2, 68 = Ipomoea reptans (Linn.) Poir.! T., P., Cangeong. V., Tancong.
- Convolvulus pes-caprae Linn., Blanco, ed. 1, 88; ed. 2, 65 = Ipomæa pes-caprae (Linn.) Roth.! T., B., Catang catang, Lagayrai, Lampayong Bagasua, Camigang, Daripay, Tagaray, Arodayday, Lambayong. I., Lambayong.
- Convolvulus batatas Linn., Blanco, ed. 1, 93; ed. 2, 68 = Ipomæa batatas (Linn.) Lam.! T., Camoti.

- Chironia capsularis Blanco, ed. 1, 102; Chironia lanosanthera Blanco, ed. 2, 71 = Ipomæa nymphæfolia (Blume) Hallier.! Erroneously reduced by Villar to Ipomæa peltata Choisy. T., Bulacan.
- *Convolvulus distillatorius Blanco, ed. 1, 95; ed. 2, 70. This species was reduced by Villar to *Ipomæa campanulata* Linn., which is certainly an error. Probably a species of *Erycibe*. V., *Bulacan*, *Buracan*.
- Convolvulus maximus Blanco, ed. 1, 91; ed. 2, 67, non Linn. = Operculina turpethum S. Manso.!
- Convolvulus catharticus Blanco, ed. 1, 94; Convolvulus longifiorus Spreng., Blanco, ed. 2, 69 = Ipomæa glaberrima Boj.! Reduced by Villar to Ipomæa longiflora R. Br. T., Bulacan.
- Convolvulus colubrinus Blanco, ed. 2, 66. Reduced by Villar to *Ipomæa* muricata Jacq., which is probably correct. Introduced and cultivated.
- Ipomoea pes-tigridis Linn., Blanco, ed. 1, 97; ed. 2, 71.!
- Ipomoea hepaticifolia Blanco, ed. 2, 72, non Linn. Reduced by Villar to Ipomoea angustifolia Jacq., but this is certainly an error. Probably a form of Merremia umbellata var. orientalis Hallier f. Blanco's specimens were from Parañaque, near Manila, flowering in January.
- Convolvulus paniculatus Linn., Blanco, ed. 1, 96; ed. 2, 70 = Ipomæa paniculata (Linn.) R. Br.! T., Puntas puntas.
- Convolvulus dentatus Blanco, ed. 1, 89; ed. 2, 66, non Vahl. = Ipomæa blancoi Choisy.!
- Polemonium obscurum Blanco, ed. 1, 103; ed. 2, 75. Reduced by Villar to Lepistemon flavescens Blume, which is probably correct. Blanco's specimens were from Bauang, Tayabas, and from Santiago, Ilocos Sur, flowering in January.
- Convolvulus reniformis Roxb., Blanco, ed. 1, 91; ed. 2, 67 = Merremia emarginata Hallier.! Reduced by Villar to Lepistemon reniformis Hassk. T., Bato bato. I., Cupit cupit.
- Convolvulus hederaceus Blanco, ed. 1, 90; ed. 2, 66, non Linn. = Hewittia bicolor Wight.!
- Evolvulus linifolius Blanco, ed. 1, 221; ed. 2, 156. Reduced by Villar to Evolvulus alsinoides Linn., which is probably correct.
- * Porania volubilis Blanco, ed. 1, 88; ed. 2, 64. Considered by Villar to be identical with *Porania volubilis* Burm., which may be correct. Blanco's specimens were from Cebu. V., *Bulacan*.
- * Convolvulus valerianoides Blanco, ed. 1, 90; Convolvulus boerhaavoides ed. 2, 67. Reduced by Villar to Breweria, as a distinct species.

 B. valerianoides (Blanco) F.-Vill. This species is known only from Blanco's description, his specimens being from Santiago Point, Province of Batangas.

SOLANACEÆ.

Solanum lycopersicum Linn., Blanco, ed. 1, 134; ed. 2, 96 = Lycopersicum esculentum Mill.! T., Camatis, Tomates.

- Solanum nigrum Linn., Blanco, ed. 1, 134; ed. 2, 96.! T., Cunti, Onti, Gamagamatisan. V., Lagpacum, Bolagtob, Lubilubi. C., Cuti, Lubilubi.
- Solanum mauritianum Blanco, ed. 1, 134; ed. 2, 96, non Scop. = Solanum verbascifolium Linn.! V., Noog noog.
- Solanum zeylanicum Blanco, ed. 1, 136; ed. 2, 97, non Scop. = Solanum ferox Linn.! T., Talong gubat, Tarambolo.
- Solanum melongena Linn., Blanco, ed. 1, 135; ed. 2, 96.! T., Talong.
- Solanum coagulans Blanco, ed. 1, 135; ed. 2, 97, non Jacq. Reduced by Villar to Solanum sanctum Linn., but is probably only a variety of Solanum melongena Linn. T., Tarumbulo, Tangantangan.
- Solanum sinense Blanco, ed. 1, 137; Solanum tuberosum Linn., Blanco, ed. 2, 97. The latter name is correct.!
- Physalis peruviana Linn., Blanco, ed. 1, 138; ed. 2, 98.! T., Potocan.
- Physalis angulata Linn., Blanco, ed. 1, 138; ed. 2, 99.! I., Tuttullacac.
- Physalis pubescens Linn., Blanco, ed. 1, 138; ed. 2, 98.!
- * Solanum serratum Blanco, ed. 1, 136; ed. 2, 97. Reduced by Villar to *Physalis indica* Linn., which is probably correct. Blanco's specimens were from Pasig, Province of Rizal, flowering in January.
- Capsicum minimum Roxb., Blanco, ed. 1, 133; ed. 2, 95.! T., Pasitis.
- Datura metel Blanco, ed. 1, 98; ed. 2, 72, non Linn., = Datura fastuosa Linn., var.! Villar considers that Blanco's species is the same as Datura alba Nees. P., T., Talamponai. V., Tavbibung.
- Datura fastuosa Linn., Blanco, ed. 1, 100; ed. 2, 73.! T., Talamponai na itim.
- Nicotiana tabacum Linn., Blanco, ed. 1, 101; ed. 2, 74.! Sp.-Fil., Tabaco.
- * Nicotiana fruticosa Linn., ? Blanco, ed. 2, 74. V. frutescens Blanco, ed. 1, 101. This species may be the same as the form described by Linnaeus. Blanco's specimens were from the pueblo of San Jose, Batangas.
- * Nicotiana pusilla Blanco, ed. 1, 100; ed. 2, 74. Reduced by Villar to Nicotiana rustica Linn., but this reference is very doubtful. Blanco's specimens were from a garden in Manila.

SCROPHULARIACEÆ.

- Stemodia ruderalis Blanco, ed. 1, 498; ed. 2, 348, non Linn. = Lindenbergia philippensis (Cham.) Benth.!
- Tala odorata Blanco ed. 1, 485; ed. 2, 338. Reduced by Villar to Limnophila menthastrum Benth., but certainly the same as Limnophila roxburghii G. Don.! T., Tala, Taramhampan. P., Talatala. I., Taratara.
- * Diceros stoloniferus Blanco, ed. 2, 349. Reduced by Villar to Limnophila repens Benth., which is certainly an error. May be Limnophila conferta Benth. Sp.-Fil., Oregano.

- Calytriplex obovata Ruiz. et Pav., Blanco, ed. 2, 361. Thunbergia stolonifera Blanco, ed. 1, 517 = Herpestes monniera H. B. K.! T., Olasamang aso.
- Legazpia triptera Blanco, ed. 2, 339 = Torenia polygonoides Benth.!
- Mimulus violaceus Blanco, ed. 2, 357 = Torenia peduncularis Benth.!
- * Vandellia multiflora Blanco, ed. 1, 505, non Don. Reduced by Villar to *Torenia asiatica* Linn., which may be correct. A small plant, flowering in October. This species is not included in the second edition.
- *Torenia paniculata Blanco, ed. 1, 486; ed. 2, 339. This species is retained by Villar as distinct, which must be considered doubtful, as Blanco states that the plant is very common, although at present it is known only from Blanco's description.
- Vandelia soriana Blanco, ed. 1, 506; Torenia soriana Blanco, ed. 2, 340. Retained by Villar as a distinct species, but is apparently the same as *Vandelia peduncularis* Benth. Blanco's specimens were from Malinta and Guadalupe, near Manila, flowering in October.
- Kyrtandra personata Blanco, ed. 1, 18; Cyrtandra personata Blanco, ed. 2, 13. Reduced by Villar to *Bonnaya brachiata* Link et Otto, which is probably correct.
- Gratiola hyssopioides Blanco, ed. 1, 11; ed. 2, 8, non Linn. Reduced by Villar to Bonnaya reptans Spreng., which is probably correct. Flowering in January.
- Kyrtandra serrata Blanco, ed. 1, 18. Reduced by Villar to *Bonnaya* veronicaefolia Spreng., which may be correct. This species is not included in the second edition.
- Kyrtandra capsularis Blanco, ed. 1, 17; Cyrtandra glaberrima Blanco, ed. 2, 12. Certainly the same as the preceding.!
- Scoparia dulcis Linn., Blanco, ed. 1, 55; ed. 2, 37.! T., Chachachachan.

OROBANCHACEÆ.

Æginetia indica Roxb., Blanco, ed. 2, 342.! T., Dagatan.

GESNERIACEÆ.

- * Hydrocotyle monopetala Blanco, ed. 1, 213; Ophiorhiza triandra Blanco, ed. 2, 65. This species was reduced by Villar to Epithema, as a distinct species, Epithema triandrum (Blanco) F.-Vill. If a species of Epithema, possibly the same as Epithema benthami Clarke. A small plant, flowering in December.
- * Kyrtandra aristata Blanco, ed. 1, 18; Cyrtandra aristata Blanco, ed. 2, 13. Referred by Villar to *Didymocarpus*, as a distinct species *D. aristata* (Blanco) F. Vill., which reference must be considered doubtful. Blanco's specimens were from Malinta, near Manila, flowering in October. Probably not a species of this family.

BIGNONIACEÆ.

- Bignonia quadripinnata Blanco, ed. 1, 499; ed. 2, 349 = Oroxylum indicum (Linn.) Vent.! T., Pincapincahan, Pincapinca, Taghilao, Abangabang.
- Bignonia spathacea Linn., Blanco, ed. 1, 499; Spathodea luzonica Blanco, ed. 2, 350 = Dolichandrone spathacea (Linn.) K. Sch.! T., Tue.
- Millingtonia quadripinnata Blanco, ed. 1, 499; ed. 2, 351 = Stereospermum quadripinnatum (Blanco) F.-Vill.! T., Baticulin. V., Ansohan.
- Millingtonia pinnata Blanco, ed. 1, 501; ed. 2, 351 = Stereospermum pinnatum (Blanco) F.-Vill.! T., Banaibanai, Botong manoc.
- Crescentia trifolia Blanco, ed. 1, 489; ed. 2, 343 = Crescentia alata H. B. K.! Sp.-Fil., Hoya cruz.

PEDALIACEÆ.

Sesamum indicum Linn., Blanco, ed. 1, 507; ed. 2, 353.! T., Linga. V., Longa. P., Langis.

ACANTHACEÆ.

- Thunbergia subsagittata Blanco ed. 1, 518; Thunbergia fragrans Blanco, ed. 2, 360. Reduced by Villar to *Thunbergia javanica* Gaertn., but is probably *Thunbergia fragrans* Roxb.
- Elytraria amara Blanco, ed. 2, 8 = Elytraria crenata Vahl.!
- * Antirrhinum molle Blanco, ed. 1, 503; ed. 2, 353. Reduced by Villar to *Hygrophila undulata* Nees, which is certainly an error.! Blanco's specimens were from Malinta, near Manila, a common plant flowering in January.
- Antirrhinum comintanum Blanco, ed. 1, 502; ed. 2, 352 = Hygrophila salicifolia Nees.! T., Mamitic.
- Ruellia uliginosa Blanco, ed. 1, 494; ed. 2, 346, non Linn. = Blechum brownei Juss.! T., Sapinsapin, Dayang. P., Torrecilla, Calaboa.
- * Ruellia repens Blanco, ed. 1, 493; ed. 2, 345, non Linn. Reduced by Villar to *Hemigraphis*, as a distinct species, *Hemigraphis repens* (Blanco) F.-Vill. Known only from Blanco's description, whose specimens were from Cebu. V., *Palad palad*.
- Acanthus doloariu Blanco, ed. 1, 487; Acanthus ilicifolius Linn., Blanco, ed. 2, 341. The latter name is correct.! T., Dilivario, Dolariu, Tingloi, Laguiolaguio. V., Titio. P., Dulauari.
- Barreliera cristata Blanco, ed. 1, 492; ed. 2, 344 = Barleria cristata Linn.! Sp.-Fil., Flor de campana.
- Barreliera prionitis Blanco, ed. 1, 491; ed. 2, 343 = Barleria prionitis Linn.! T., Cocong manoc.

- Justicia gandarussa Blanco, ed. 1, 14; ed. 2, 10 = Pseuderanthemum bicolor (Schrk.) Radlk.! T., Cinco llagas, Silisilihan, Limang sugat.
 V., Mopio, Maladosos, Panaptun. P., Alyopyop, Mandalusa.
- * Ruellia secunda Blanco, ed. 1, 495; ed. 2, 346 = Lepidagathis secunda (Blanco) Nees. A species known only from Blanco's description, but which is probably identical with Lepidagathis luzona Nees.
- * Dianthera americana Blanco, ed. 1, 16, non Linn.; Dianthera ciliata Blanco, ed. 2, 12. Reduced by Villar to Justicia mollissima Nees, which is a synonym of Justicia simplex Don., and a species as yet not reported from the Philippines. Blanco's species is, possibly, a form of Justicia procumbens Linn. A plant a foot high, common in Malinta.
- Justicia dalaora Blanco, ed. 1, 14; ed. 2, 10. Reduced by Villar to Dianthera (Justicia) dichotoma (Blume) Clarke, which is probably correct. V., Dalaora.
- Justicia nasuta Linn., Blanco, ed. 1, 14; ed. 2, 10. Considered by Villar to be the Linnean plant = Rhinacanthus communis Nees. T., Tagactagac.
- Justicia picta Linn., Blanco, ed. 1. 12; ed. 2, 9 = Graptophyllum pictum (Linn.) Griff.! (Graptophyllum hortense Nees.) T., Moradong maputi, Balasbas. V., Balasbas, Antolang, San Francisco.
- Justicia ecbolium Blanco, ed. 1, 13; ed. 2, 10, non Linn. = the preceding! The variety with purplish leaves. T., Morado, Ternate, Balasbas. V., Balasbas. P., V., Hojas moradas, Balasbas malomay, Yovas, Ataiatai.
- * Ruellia contorta Blanco, ed. 1, 496; ed. 2, 347. Reduced by Villar to Peristrophe as a distinct species, P. contorta (Blanco) F. Vill., a species known only from Blanco's description. Blanco's specimens were from Bauang, Province of Batangas, flowering in February. T., Sapinsapin.
- Justicia viridis Blanco, ed. 1, 15; ed. 2, 11, non Forsk. = Hypoestes malaccanus Wight.!

VERBENACEÆ:

- Lantana viburnoides Blanco, ed. 2, 345, non Vahl. = Lantana camara Linn!
- Verbena capitata Forsk., Blanco, ed. 1, 19; ed. 2, 14 = Lippia nodiflora (Linn.) Rich.! T., Chachachachahan.
- *Callicarpa americana Blanco, ed. 1, 517; ed. 2, 360, non Linn. = Callicarpa blancoi Rolfe.! Reduced by Villar to Callicarpa bicolor Juss., and certainly closely related to that species. T., Palis, Tubang dalag.
- Tectona grandis Linn. f., Blanco, ed. 1, 130; ed. 2, 92.! Sp.-Fil., Teca, Ticla. V., Dalondon, Yate, Calayate.

- * Diospyros tectona Blanco, ed. 2, 609. This name is published in the index to native names, under *Dalandon*, without description. Villar reduces it to *Tectona hamiltoniana* Wall. The name should be discarded.
- * Premna serratifolia Blanco, ed. 2, 342, non Linn. Reduced by Villar to Premna foetida Reinw., which is undoubtedly an error.
- Premna odorata Blanco, ed. 1, 488; ed. 2, 341. Erroneously reduced by Villar to *Premna pubescens* Blume, a species that does not extend to the Philippines. Blanco's species is certainly identical with *Premna vestita* Schauer, which is a much later name. T., *Alagao*. V., *Adgao*, *Pamuhat*. P., *Tanglay maloto*. I., *Anobran*.
- * Premna cordata Blanco, ed. 1, 489; Premna tomentosa Blanco, ed. 2, 342, non Wall. Considered by Villar to be identical with *Premna tomentosa* Wall., but this is certainly an error. T., *Malaapi*.
- Premna nauseosa Blanco, ed. 1, 489; Premna integrifolia Blanco, ed. 2, 342. A good species, the first name being the correct one. Erroneously reduced by Villar to Premna mucronata Roxb.
- Gmelina asiatica Blanco, ed. 1, 492; ed. 2, 344, non Linn. = Gmelina villosa Roxb.! T., V., Bagaboboi, Balabalayan.
- Gmelina inermis Blanco, ed. 1, 493; ed. 2, 345 = Gmelina asiatica Linn. ?, or a form of the preceding.
- Vitex repens Blanco, ed. 1, 513; ed. 2, 358 = Vitex ovata Thunb. (Vitex trifolia Linn. f. var. obovata Benth.) T., Lagunding gapang.
- Vitex trifolia Linn. f., Blanco, ed. 1, 513; ed. 2, 358.! T., Lagundi. V., Gapasgapas. I., Danglay.
- Vitex leucoxylon Blanco, ed. 1, 516; ed. 2, 359, non Linn. Reduced by Villar to Vitex negundo Linn., which is probably correct. T., Lagundi, Molavin.
- Vitex altissima Blanco, ed. 1, 516; ed. 2, 359, non? Linn. = Vitex littoralis Decne.! T., Molavin, Lagundi.
- Vitex latifolia Blanco, ed. 1, 514; ed. 2, 358 = Vitex pubescens Vahl.! T., Molauin, Hamaraon, Bulaon.
- * Vitex geniculata Blanco, ed. 1, 514; ed. 2, 299. A species closely related to Vitex littoralis Decne., but perhaps distinct. T., Molavin.
- Clerodendron capsulare Blanco, ed. 1, 509; ed. 2, 355 = Clerodendron inerme R. Br.!
- Volkameria grandiflora Blanco, ed. 1, 512; ed. 2, 357 = Clerodendron macrostegium Schauer. T., Malapotocan, Bagauac. I., Agboligan, Nacboligan.
- Clerodendron fortunatum Blanco, ed. 1, 508; ed. 2, 354, non Linn. = Clerodendron blancoi Naves.! Reduced by Villar to Clerodendron infortunatum Gaertn. T., Casopanguil gubat.

- Volkameria inermis Blanco, ed. 1, 511, non Linn; Bolkameria casopanguil Blanco, ed. 2, 356 = Clerodendron intermedium Cham.! T., Casopanguil, Laroan, Anito, Macalalauang, Iginga. V., Asuangai, Pacapis, Colocolog, Alocasoc.
- Ligustrum quadriloculare Blanco, ed. 1, 10; ed. 2, 7. This species is the same as Clerodendron navesianum Vidal (Clerodendron blancoanum F.-Vill.) but the species is very near if not identical with Clerodendron longiforum Decne. T., Balictarin.
- Litsea luzonica Blanco, ed. 2, $284 = Symphorema\ luzonicum\$ (Blanco) F.-Vill.! (Symphorema luzoniense Vidal.) This species is described in the first edition under the native name only. P., Balaibai.
- Avicennia nitida Blanco, ed. 1, 504, non Jacq.; Avicennia tomentosa R. Br., Blanco, ed. 2, 353 = Avicennia officinalis Linn.!

LABIATÆ.

- Ocimum americanum Blanco, ed. 1, 480; ed. 2, 335, non Linn. = Ocimum basilicum Linn.! T., P., Solasi. V., Bonac.
- Ocimum citriodorum Blanco, ed. 2, 591 = the preceding species.!
- Ocimum album Blanco, ed. 1, 479, non Linn.; Ocimum virgatum Blanco, ed. 2, 334, non Linn. = Ocimum gratissimum Linn.! T., Locoloco. V., Coloncogon. P., Locoloco.
- Ocimum sanctum Linn., Blanco, ed. 1, 480; ed. 2, 334.! T., Balanoi.
- Ocimum flexuosum Blanco, ed. 1, 480; ed. 2, 334, non Linn. = the preceding,! the purple-stemmed form. T., Balanoi.
- Ocimum tenuiflorum Blanco, ed. 1, 481; ed. 2, 335 = Moschosma polystachyum Benth.!
- Coleus suganda Blanco, ed. 1, 483; ed. 2, 337. Reduced by Villar to Coleus aromaticus Benth., which is probably correct. Sp. Fil., Orega. o. T., Suganda.
- * Coleus pumilus Blanco, ed. 1, 482; ed. 2, 336. Reduced by Villar to Coleus acuminatus Benth., which may be correct. Blanco's specimens were from Pasig, Province of Rizal, flowering in October. T., Malamayana.
- Coleus grandifolius Blanco, ed. 1, 482; ed. 2, 336. Reduced by Villar to Coleus atropurpureus Benth., which is probably correct. T., V., P., Mayana, Maliana, Badiara, Marayapa.
- Pycnanthemum elongatum Blanco, ed. 2, 333 = Hyptis spicigera Lam.!
- Thymus virginicus Blanco, ed. 1, 478, non Linn.; Pycnanthemum decurrens Blanco, ed. 2, 333 = Hyptis capitata Jacq.! T., Combarcombaran, Lingalingahan.
- Thymus biserratus Blanco, ed. 1, 478; Pycnanthemum subulatum Blanco, ed. 2, 333 = Hyptis brevipes Poir.! T., Combarcombaran.
- Marrubium indicum Blanco, ed. 1, 477; ed. 2, 332, non Burm. = Hyptis suaveblens Poir.! T., Soob cabayao.

- Mentha cablin Blanco, ed. 1, 472; Mentha auricularia Blanco, ed. 2, 329, non Linn. = Pogostemon cablin (Blanco) Benth.! T., Cablin, Carlin. V., Cadlom, Catluen, Cadlin. P., Cablin.
- Mentha crispa Blanco, ed. 1, 474; ed. 2, 330, non Linn. = Mentha arvensis Linn.! Sp.-Fil., Yerba buena.
- Salvia violacea Blanco, ed. 2, 14, non Retz. et. Pav. = Salvia plebeia R. Br.!
- Rosmarinus officinalis Linn., Blanco, ed. 1, 20; ed. 2, 15.! Sp.-Fil., Romero.
- Phlomis alba Blanco, ed. 1, 474; ed. 2, 330, non Forsk. = Anisomelis ovata R. Br.! T., Talingharap.
- Stachys artemesia Lour., Blanco, ed. 1, 476; ed. 2, 331 = Leonurus sibiricus Linn.! T., Camariang songsong.
- Phlomis zeylanica Blanco, ed. 1, 475; ed. 2, 331, non Linn. Reduced by Villar to Leucas aspera Spreng., but certainly is Leucas linifolia Spreng.! T., Pansipansi, Solasolasihan, Carucansoli. V., Pansipansi, Paipaisi.

PLANTAGINACEÆ.

Plantago crenata Blanco, ed. 1, 56; Plantago media Blanco, ed. 2, 38, non Linn. Reduced by Villar to Plantago erosa Wall., which is a synonym of Plantago major Linn. Probably correctly referred. Flowering in February. T., Lantin.

NYCTAGINACEÆ.

- Mirabilis longiflora Blanco, ed. 1, 77; ed. 2, 57, non Linn. = Mirabilis jalapa Linn.! Sp.-Fil., Oracion, Diego de noche, Bella de noche. T., Guilalas, Suspiros.
- Boerhaavia diffusa Linn., Blanco, ed. 1, 8; ed. 2, 6 = Boerhaavia repens Linn.! T., Paanbalivis. I., Tabtabocol ti Nuang.
- Pisonia aculeata Linn., Blanco, ed. 1, 195; ed. 2, 137.! T., Digquit digquit. I., Puriquet. T., Panacla.
- Cordia olitoria Blanco, ed. 1, 123; ed. 2, 88. Reduced by Villar to *Pisonia inermis* Forst., which is cited as a synonym of *Pisonia alba* Spanoghe. A species rather commonly cultivated in Manila, but rarely producing flowers. T., *Maluco*, *Coles maluco*.
- * Buginvillea racemosa Blanco, ed. 1, 307; ed. 2, 214. Reduced by Villar to *Pisonia excelsa* Blume, which is certainly an error. Blanco's specimens were from Angat, Province of Bulacan, no native name given.
- *Cedrota guiarrensis Blanco, ed. 2, 213, not Ræusch. Reduced by Villar to *Pisonia umbellifera* Seem., which is the same as *Pisonia excelsa* Blume. Blanco's specimens were from Cebu. V., *Tac-an*.

AMARANTACEÆ.

- Celosia baccata Retz., Blanco, ed. 1, 193; Deeringia celosioides R. Br., Blanco, ed. 2, 135. The latter name is correct. T., Hagorilis. V., Hantilimocon.
- Celosia cristata Linn., Blanco, ed. 1, 191; Celosia coccinea Linn., Blanco, ed. 2, 134. The former name is correct. T., Palongpalongan.
- Celosia argentea Linn., Blanco, ed. 1, 192; ed. 2, 135.! T., Cadayohan, Quindayohan.
- * Celosia bicolor Blanco, ed. 1, 191; Celosia glauca Blanco, ed. 2, 135, non Rottl. Reduced by Villar to Celosia philippica (Weinm.) Steud.
- Amarantus spinosus Linn., Blanco, ed. 1, 710; ed. 2, 491.! T., Quilites, Orayi. P., Ayantoto. V., Calites, Tililes, Harum, Orayi, Bayangbayang. I., Cuanton.
- Amarantus mangostanus Linn., Blanco, ed. 1, 711; ed. 2, 492. Probably correct, although Villar reduces Blanco's species to Amarantus melancholicus Linn., which is a synonym of Amarantus gangeticus Linn. T., Halon.
- Illecebrum lanatum Linn., Blanco, ed. 1, 190; Celosia lanata Linn., Blanco, ed. 2, 134 = Aerua lanata (Linn.) Juss.!
- Achyranthes aspera Linn., Blanco, ed. 1, 188; ed. 2, 133.! T., Hangor, Hangot, Docotdocot, Libai. P., Angud. V., Hangor.
- Achyranthes villosa Blanco, ed. 1, 189; ed. 2, 134, non Forsk. Reduced by Villar to Alternanthera denticulata R. Br., probably however only a form of the common Alternanthera sessilis R. Br. T., Bonga bonga.
- Gomphrena globosa Linn., Blanco, ed. 1, 198; ed. 2, 139.! T., Buquingan

CHENOPODIACEÆ.

- Chenopodium ambrosioides Linn., Blanco, ed. 1, 200; ed. 2, 140.! T., V., P., Aposotis.
- Basella rubra Linn., Blanco, ed. 1, 215; ed. 2, 151 = Basella alba Linn.! T., Libato. I., Ilaibaquir.
- Basella lucida Linn., Blanco, ed. 1, 151; ed. 2, 273 = the preceding.! T., Grana.
- Gomphrena volubilis Blanco, ed. 1, 199; ed. 2, 140 = Anredera scandens Juss.!

POLYGONACEÆ.

- Polygonum stoloniferum Blanco, ed. 1, 314; ed. 2, 219. Reduced by Villar to Polygonum barbatum Linn., var. vulgare Meisn., which is probably correct. T., Subsuban. P., Canubsuban.
- * Polygonum bellardi Blanco, ed. 1, 314; ed. 2, 219, non Boiss. Reduced by Villar to *Polygonum orientale* Linn., which is probably an error. Blanco's specimens were from Taguig, Province of Rizal, flowering in March. T., Subsuban. P., Canucsuban.

Rheum muricatum Blanco, ed. 1, 327; Rumex muricatus Blanco, ed. 2, 195 = Rumex maritimus Linn.! Blanco's specimens were from cultivated plants.

NEPENTHACEÆ.

Nepenthes alata Blanco, ed. 1, 805; ed. 2, 555. A good species.! Nepenthes ventricosa Blanco, ed. 1, 807; ed. 2, 556. A good species.!

CYTINACEÆ.

Rafflesia philippinensis Blanco, ed. 2, 565; Rafflesia lagascae Blanco, ed. 2, 595 = Rafflesia manillana Teschem.!

ARISTOLOCHIACEÆ.

- * Aristolochia sericea Blanco, ed. 1, 283; ed. 2, 198. Reduced by Villar to *Bragantia corymbosa* Griff., which may be correct, although the latter species is at present not known from the Philippines. Blanco's specimens were from Agoo, Province of Union.
- * Aristolochia indica Blanco, ed. 1, 282; ed. 2, 197, non? L. Considered by Villar to represent the Linnean species, but probably only a form of Aristolochia tagala Cham. T., Timbagan, Malaubai.
- Aristolochia subsagittata Blanco, ed. 1, 283; ed. 2, 197 = Aristolochia tagala Cham.! T., Timbangan, Timbangantimbangan, Malaubi. I., Taointaoin.

PIPERACEÆ.

- Piper parvifolium Blanco, ed. 1, 23; ed. 2, 17. Reduced by Villar to Piper caninum. A. Dietr., which is probably correct. T., Sabia.
- * Piper obliquum Blanco, ed. 1, 22; ed. 2, 16. Reduced by Villar to Piper corylistachyon C. DC. If this is correct, Blanco's name is the earlier. T., V., Litlit, Litbit, Saog, Sangilo. B., Nopau.
- Piper betel Blanco, ed. 1, 22; ed. 2, 16 = Piper betle Linn.! T., B., Itmo, Mamin. V., Buyo, Mamon. P., Samat.
- Piper nigrum Linn., Blanco, ed. 1, 21; ed. 2, 16.! Sp.-Fil., Pimienta. T., Malisa.
- * Piper anisumolens Blanco, ed. 1, 23; Piper anisodorum Blanco, ed. 2, 16. Reduced by Villar to Piper marginatum Jacq., an American species, which is probably correct. T., Buyo de anis.

CHLORANTHACEÆ.

Chloranthus inconspicuus Blanco, ed. 2, 54, non Swartz. Reduced by Villar to Chloranthus officinalis Blume, which may be correct. Chloranthus brachystachyus Blume, is, however, the more common species in the Philippines.

MYRISTICACEÆ.

- Myristica luzonica Blanco, ed. 1, 664; ed. 2, 462; 463 = Myristica philippinensis Lam.! T., Talangtalang, Anis moscada, Anis cahoi, Duguan.
- * Sterculia glomerata Blanco, ed. 1, 764; ed. 2, 525. This species is reduced by Villar to *Myristica corticosa* Hook. f. et Thoms., a species that does not extend to the Philippines, considered by Warburg to be possibly the same as *Knema heterophylla* (F.-Vill.) Warb. T., *Hindurugo*.
- * Sterculia decandra Blanco, ed. 1, 766; ed. 2, 525. Reduced by Villar to *Myristica intermedia* Blume, a species that does not extend to the Philippines. Probably a species of *Myristica*. Blanco's specimens were from Balanga, Province of Bataan. T., *Banabanalo*.

LAURACEÆ.

- * Laurus martinicensis Blanco, ed. 1, 317; ed. 2, 223, non alibrum. Reduced by Villar to *Beilschmiedia madang* Blume, which is certainly an error. Probably a species of *Lauracea*, but the genus is doubtful. A tree flowering in December. T., *Dinglas*.
- * Salgada laurifolia Blanco, ed. 2, 221. Reduced by Villar to Eusideroxylon borneense T. et B., which must be considered very doubtful. Blanco's specimens were from Angat, Province of Bulacan.
- * Laurus cassia Blanco, ed. 1, 317; ed. 2, 223, non Linn. Considered by Villar to represent Laurus cassia Burm, and by Villar reduced to a variety of Cinnamomum zeylanicum Breyn. Certainly erroneously referred. Blanco's specimens were from Balanga, Province of Bataan T., Samilin.
- Laurus cinnamomum Blanco, ed. 1, 319; ed. 2, 225, non Linn. Referred by Villar to *Cinnamomum burmanni* Nees, which may be correct, as Blanco's specimens were from a cultivated plant.
- Laurus culilaban Blanco, ed. 1, 315; ed. 2, 222, non aliorum. This species was reduced by Villar to Cinnamomum pauciflorum Nees, which is certainly an error, as that species does not extend to the Philippines. Probably identical with the form described by Vidal as Cinnamomum mercadoi. T., Calingag, Macalingag. V., Calingag, Calinga. P., Calingad. I., Candaroma.
- Laurus persea Linn., Blanco, ed. 2, 224 = Persea gratissima Gaertn., an American species, cultivated. Blanco's specimens were from Santa Ana, a suburb of Manila.
- Sebifera glutinosa Blanco, ed. 1, 819; ed. 2, 566, non Lour. Reduced by Villar to *Litsea chinensis* Lam., which is probably correct. T., *Pusopuso*. I., *Sablut*.

- Sebifera balongai Blanco, ed. 1, 820; ed. 2, 567. Reduced by Villar to Litsea literalis Blume, but Vidal refers Litsea literalis F.-Vill. to Litsea chinensis Lam. Probably only a form of the latter species. T., Balongai.
- * Olax baticulin Blanco, ed. 2, 586 = Litsea obtusata F.-Vill.? Impossible to determine definitely if Blanco's species is the same as Villar's. T., Baticulin.
- Laurus lanosa Blanco, ed. 1, 318; ed. 2, 224. Reduced by Villar to Litsea villosa Blume, which may be correct. T., V., Batobato.
- * Ajovea malabonga Blanco, ed. 1, 233; Laurus hexandra Blanco, ed. 2, 222, non aliorum. Reduced by Villar to *Iteadapne confusa* Blume, which must be considered doubtful, as the latter species is unknown from the Philippines. A tree flowering in August. T., *Malabonga*.
- Cassytha filiformis Linn., Blanco, ed. 1, 321; ed. 2, 226.! T., Malabohoo.
 Hernandia sonora Blanco, ed. 1, 689; ed. 2, 478, non Linn. = Hernandia peltata Meissn.! T., Colongcolong.

THYMELAEACEÆ.

- * Daphne indica Blanco, ed. 1, 309; ed. 2, 215, non? Linn. Considered by Villar to represent the Linnean species, probably, however, the form described as Wickstroemia viridiflora Meissn. T., Salago, Malasampaga.
- Daphne foetida Linn. ? Blanco, ed. 1, 308; ed. 2, 214. Certainly only a form of the preceding.! T., Salago, Malasampaga.
- * Daphne aquilaria Blanco, ed. 1, 310; ed. 2, 216. Reduced by Villar to Wickstroemia ovata Mey. Vidal, however, states that Wickstroemia ovata F.-Vill., is the same as W. indica Mey. I., Arandon.
- Dais laurifolia Blanco, ed. 1, 375; ed. 2, 263, non Jacq. = Phaleria perrottetiana (Decne.) F.-Vill.!

ELAEAGNACEÆ.

Elaeagnus angustifolia Blanco, ed. 1, 74; ed. 2, 53, non Linn. = Elaeagnus latifolia Linn.! T., Alingaro.

LORANTHACEÆ.

- Lonicera symphoricarpos Blanco, ed. 1, 161; Loranthus philippinensis Cham. et Schlecht., Blanco, ed. 2, 164. The second name is probably correct. T., Dapo.
- *Loranthus paucifiorus Blanco, ed. 1, 235; Lonicera tomentosus Blanco, ed. 2, 164, non aliorum. Considered by Villar to be a distinct species and retained under the former name. Blanco's specimens were parasitic on Spondias. T., Dapo sa cahoy. I., Binbinguay.

- * Hillia longiflora Blanco, ed. 1, 235; ed. 2, 165. Reduced by Villar to Loranthus ampullaceus Linn., which is probably an error, as the latter species is not found in the Philippines. Blanco's specimens were parasitic on Mangifera altissima Blanco, a tree known to the natives as pajo. Flowering in April. T., Dapo sa pajo.
- *Brabejum ? caliculatum Blanco, ed. 2, 39. Reduced by Villar to Loranthus, as a distinct species, Loranthus blancoanus F.-Vill. Possibly a species of Loranthus. Blanco's specimens were from Cebu. V., Malabachao.
- Fusanus? parasitus Blanco, ed. 2, 53. Reduced by Villar to Viscum orientale Willd., which is probably correct. Blanco's specimens were parasites on a species of Pterocarpus in Angat, Province of Bulacan, flowering in February.

EUPHORBIACEÆ.

- Euphorbia dulcis Blanco, ed. 1, 412; ed. 2, 287, non Linn. Reduced by Villar to Euphorbia laevigata Vahl. = Euphorbia atoto Forst.!
- Euphorbia parannaquensis Blanco, ed. 2, 286 = Euphorbia serrulata Reinw.!
- Euphorbia capitata Lam., Blanco, ed. 1, 411; Euphorbia hirta Linn.,
 Blanco, ed. 2, 286 = Euphorbia pilulifera Linn.! T., Batobatonis,
 Sayican, Gatasgatas, Golandrina. V., Gatasgatas, Buyayava, Tavava,
 Bovi. P., Malismalis, Sisiohan, Bolobtones, Magatas.
- Euphorbia pentagona Blanco, ed. 1, 413; ed. 2, 287, non Haw. Reduced by Villar to Euphorbia neriifolia Linn., which is probably correct. B., T., P., Sorosoro, Sorog sorog, Bait, Lengua de perro. I., Carambuaya.
- Euphorbia tirucalli Linn., Blanco, ed. 1, 412; ed. 2, 287.! T., Catuit. Sp. Fil., Consuelda.
- Clutia stipularis Linn., Blanco, ed. 1, 818; ed. 2, 564 = Bridelia stipularis Blume.! T., Hingongoto, Lobalob. I., Calondagui, Carabaui.
- Gluta orgyalis Blanco, ed. 2, 451 = CLEISTANTHUS ORGYALIS (Blanco). (Cleistanthus blancoi Rolfe, Journ. Linn. Soc. Bot. 21: 315; Vidal, Rev. Pl. Vasc. Filip. 234.) Erroneously reduced by Villar to Cleistanthus ferrugineus Muell. Arg. Blanco's specimens were from Angat, Province of Bulacan, flowering in April.
- Kirganelia triandra Blanco, ed. 1, 711; ed. 2, 492 = Phyllanthus triandrus Muell. Arg.!
- Kirganelia alba Blanco, ed. 1, 713; ed. 2, 494 = Glochidion album (Blanco) Boerl. (Phyllanthus albus Muell. Arg.) Blanco's specimens were from Angat, Province of Bulacan. It is represented by Nos. 3162 and 3797 Merrill.
- Kirganelia villosa Blanco, ed. 1, 712; ed. 2, 493, non Phyllanthus villosus Poir. = Glochidion llanosii Muell. Arg.!

- * Kirganelia nigrescens Blanco, ed. 1, 712; ed. 2, 493. Reduced by Villar to Glochidion molle Blume, which may be correct. A shrub flowering in May.
- Cicca acidissima Blanco, ed. 1, 700; ed. 2, 486 = Phyllanthus distichus Muell. Arg.! T., Iba, Banquilin. V., Poras, Layohan. P., Iba.
- Cicca decandra Blanco, ed. 1, 701; ed. 2, 487 = Phyllanthus reticulatus Muell. Arg.! T., Tinatinaan, Tintatintahan, Malatinta.
- Phyllanthus carolinianus Blanco, ed. 1, 691, non Walt. Phyllanthus kirganelia Blanco, ed. 2, 480, non Willd. = Phyllanthus niruri Muell. Arg.!
- Kirganelia pumila Blanco, ed. 1, 712; ed. 2, 493. Reduced by Villar to *Phyllanthus niruri* Muell. Arg., as a variety, but more probably a form of *P. urinaria* Linn., or *P. simplex* Muell. Arg.
- *Phyllanthus niruri Blanco, ed. 1, 690, non Linn.; Phyllanthus tetrander Blanco, ed. 2, 480, non Roxb. Considered by Mueller to represent a distinct species, which he calls Phyllanthus blancoanus. This species is known only from Blanco's description. Blanco's specimens were from Bauang, Province of Batangas, a shrub 5 feet high, flowering in May.
- Cicca pentandra Blanco, ed. 1, 701; ed. 2, 486 = Fluggea obovata Wall.! T., Botolan.
- Cansiera pentandra Blanco, ed. 1, 73; ed. 2, 53 = Antidesma leptocladum genuinum Muell. Arg.! Erroneously reduced by Villar to Antidesma cumingiana Muell. Arg. T., Bignay pogo.
- Cansiera grossularioides Blanco, ed. 1, 73; ed. 2, 53. Reduced by Villar to Antidesma ghaesembilla Gaertn., var. paniculatum Muell. Arg. Certainly a form of this species! T., Cabogbog.
- Cansiera rheedii Blanco, ed. 1, 73; ed. 2, 52, non aliorum = Antidesma ghaesembilla Gaertn,! the pubescent form. T., Bignai pogo.
- Antidesma spicata Blanco, ed. 1, 794; Antidesma alexiteria Blanco, ed. 2, 547, non alior. Reduced by Villar to Antidesma ghæsembilla Gaertn., which is certainly correct.! T., Calamantao.
- Stilago bunius Linn., Blanco, ed. 1, 782; ed. 2, 539 = Antidesma bunius (Linn.) Spreng.! T., Bignai, Bugnai.
- Jatropha curcas Linn., Blanco, ed. 1, 759; ed. 2, 522.! T., Tuba.
 V., Casla. I., Tavatava.
- Jatropha janipha Blanco, ed. 1, 758; ed. 2, 521, non Linn. = Jatropha
 multifida Linn.!
- Aleurites Iobata Blanco, ed. 1, 756; Aleurites triloba Forst., Blanco, ed. 2, 520 = Aleurites moluccana Willd.! T., Lumban, Capili.
- Aleurites lanceolata Blanco, ed. 1, 757; ed. 2, 521 = the preceding.!
- Aleurites trisperma Blanco, ed. 1, 755; Aleurites saponaria Blanco, ed. 2, 519. A distinct species.! T., Balocanad, Baguilumban, Calumban. V., P., Balucanag.

- Croton glandulosum Blanco, ed. 1, 754; Croton muricatum Blanco, ed. 2, 518 = Croton tiglium Linn.! T., Tuba, Camaisa, Balanti.
- * Croton drupaceus Blanco, ed. 2, 519 non Roxb. Reduced by Villar to Croton luzonensis Muell. Arg., probably, however, only a form of the preceding. Blanco's specimens were from Guadalupe, near Manila, flowering in October. T., Balanti.
- Croton variegatus Linn., Blanco, ed. 1, 751; ed. 2, 517 = Codiaeum variegatum (Linn.) Muell. Arg.! T., Saguilala. V., Calipayang. Sp.-Fil., Buena vista, San Francisco.
- Adelia acidoton Blanco, ed. 1, 815; ed. 2, 562, non Linn. = Sumbavia rottleroides Baill.! T., Aliparo.
- Jatropha manihot Linn., Blanco, ed. 1, 760; ed. 2, 522 = Manihot utilissima Pohl.! T., Camoting cahoy.
- * Adelia resinosa Blanco, ed. 2, 562. Reduced by Villar to Claoxylon wallichianum Muell. Arg., which is probably an error. This species is described in the first edition under the native name Balilangouag. Blanco's specimens were from Pasig, Province of Rizal.
- * Niota globosa Blanco, ed. 2, 214. Reduced by Villar to Claoxylon indicum Muell. Arg., which may be correct. Blanco's specimens were from Angat, Province of Bulacan.
- Acalypha angatensis Blanco, ed. 1, 750; ed. 2, 516. Considered by Villar to be a distinct species, probably, however, only a form of Acalypha stipulacea Klotzsch. Blanco's specimens were from Angat, Province of Bulacan, a shrub, flowering in December.
- Acalypha tomentosa Blanco, ed. 1, 750. Reduced by Villar to Acalypha grandis Muell. Arg., which may be correct, or it may be a form of the next species. This species is not included in the second edition.
- Acalypha glandulosa Blanco, ed. 1, 749; ed. 2, 516, non Cav. = Acalypha stipulacea Klotzsch.!
- Acalypha caroliniana Blanco, ed. 1, 748; ed. 2, 515, non Walt. = Acalypha indica Linn.!
- Adelia glandulosa Blanco, ed. 1, 814. Reduced by Villar to Alchornea javensis Muell. Arg., which is probably correct. This species is not included in the second edition. Blanco's specimens were from Santa Maria, Province of Ilocos Sur.
- *Tragia innocua Blanco, ed. 2, 479. Reduced by Villar to Alchornea villosa Muell. Arg., which is probably an error. Blanco's specimens were from Angat, Province of Bulacan, flowering in February.
- * Excaecaria sicca Blanco, ed. 1, 787; ed. 2, 542. Reduced by Villar to Alchornea mollis Muell. Arg., which is certainly an error. Blanco states that his species is poisonous, and common along the margins of streams. T., Balanti.
- * Prockia albicans Blanco, ed. 1, 430; ed. 2, 299. Reduced by Villar to Alchornea blumeana Muell. Arg., which is certainly an error. T., Busilac.

- Adelia monoica Blanco, ed. 2, 561 = Mallotus moluccanus (Linn.) Muell. Arg.!
- Adelia bernardia Blanco, ed. 1, 814, non Linn.; Adelia barbata Blanco, ed. 2, 561 = Mallotus ricinoides (Pers.) Muell. Arg.! T., Alim, Taquip asin. V., Alum, Arum. P., Pacalcal, Vilos. C., Tapit. Sp.-Fil., Tres puntas.
- * Adelia papillaris Blanco, ed. 2, 562. Reduced by Villar to *Mallotus zollingeri* Muell. Arg., which is certainly an error, as the latter species does not extend to the Philippines. Blanco's specimens were from Guadalupe, near Manila.
- Croton lacciferum Blanco, ed. 1, 752; ed. 2, 517 = Macaranga tanarius Muell. Arg., var. tomentosa Muell. Arg.! T., Binonga, Minonga. V., P., Binonga. P., Bilau. I., Ginabang, Samac.
- Croton grandifolium Blanco, ed. 1, 753; ed. 2, 518 = Macaranga mappa (Linn.) Muell. Arg.! T., Bingabing.
- Ricinus communis Linn., Blanco, ed. 1, 761; ed. 2, 523.! T., Tangantangan, Lingasina. I., Tangantangan, Tavatavangsina.
- Lumanaja fluviatilis Blanco, ed. 1, 821; ed. 2, 568 = Homonoia riparia Lour.! T., Lomonai, Agoioi.
- Excæcaria lævis Blanco, ed. 1, 788; ed. 2, 543 = Homalanthus populneus (Geisel.) Pax.! T., Botong gubat.
- * Tragia bracteata Blanco, ed. 2, 481. Reduced by Villar to Pimelodendron amboinicum Hassk., which is certainly an error. Blanco's specimens were from Angat, a shrub, flowering in February. T., Tablon.
- Excæcaria agallocha Linn., Blanco, ed. 1, 786; ed. 2, 541.! T., Buta, Butabuta. V., Alipata, Lipata, Himbabao, Siac. P., Butabuta.

URTICACEÆ.

- Celtis philippinensis Blanco, ed. 1, 197; ed. 2, 139. A good species.! T., Malaitmo.
- Celtis lima Blanco, ed. 1, 197; ed. 2, 139 = Trema amboinensis Blume.! T., Hanarion.
- Calius lactescens Blanco, ed. 1, 698; ed. 2, 485 = Streblus asper Lour.! T., Calios. I., Alodig.
- Malaisia tortuosa Blanco, ed. 1, 789; ed. 2, 543.! A good genus and species. T., Malaisis.
- Morus Iuzonica Blanco, ed. 1, 703; Broussonetia Iuzonensis Blanco, ed. 2, 488 = Allaanthus Iuzonicus (Blanco) F. Vill.! T., Himbabao.
- Morus alba Linn., Blanco, ed. 1, 705; ed. 2, 489.! T., Morera.
- Ficus indica Linn., Blanco, ed. 1, 681; ed. 2, 473. Reduced by Villar to *Ficus saxophila* Blume, but Blanco probably correctly interpreted the Linnean species. *Ficus saxophila* Blume is not known from the Philippines. T., P., *Baliti*. V., *Dalaguit*, *Nonoc*.

- * Ficus payapa Blanco, ed. 1, 683; ed. 2, 475. Reduced by Villar to Ficus microcarpa Linn. f., which is a synonym of Ficus retusa Linn. Possibly correctly referred. T., Payapa.
- * Ficus dicarpa Blanco, ed. 1, 682; Ficus nepalensis Blanco, ed. 2, 474 non Spreng. Reduced by Villar to Ficus hæmatocarpa Blume, which is a synonym of Ficus benjamina Linn., which may be correct. Planco's specimens were from Cebu. V., Talicot.
- Ficus hispida odorata Blanco, ed. 1, 686; ed. 2, 476 = Ficus odorata (Blanco) Merrill, Govt. Lab. Publ. 17: 15. 1904. Erroneously reduced by Villar to Ficus pungens Reinw. T., Agosos.
- Ficus aspera Forst., Blanco, ed. 1, 676; ed. 2, 476. Blanco gives only a partial translation of the description of this species, without definitely referring any form to the species. Villar erroneously reduces it to Ficus hirta Vahl.
- Ficus hispida linearis Blanco, ed. 1, 685; ed. 2, 476 = Ficus cumingii Miq.!. Erroneously reduced by Villar to Ficus heterophylla Linn. T., V., P., Isis, Asis, Issio, Isioisio.
- Ficus hispida hastata Blanco, ed. 1, 685; ed. 2, 476 = Ficus quercifolia Roxb.!
- Ficus heterophylla Blanco, ed. 1, 685; ed. 2, 476, non Linn. = Ficus quercifolia Roxb.! T., Asis, Isis.
- * Ficus argentea Blanco, ed. 1, 681; ed. 2, 473. Reduced by Villar to Ficus polycarpa Wall., which is a synonym of Ficus ribes Reinw., and which is certainly an error.! Blanco's specimens were from the seashore at Mariveles, Province of Bataan.
- * Ficus rostrata Blanco, ed. 1, 679; ed. 2, 472. Reduced by Villar to Ficus radicans Roxb., which is a synonym of Ficus rostrata Lam. Possibly correctly referred.
- * Ficus aspera volubilis Blanco, ed. 1, 676; ed. 2, 472. Erroneously reduced by Villar to Ficus hederacea Roxb., which is probably a synonym of Ficus scandens Roxb. Blanco's specimens were from Calapan, Mindoro, a small vine with very rough leaves. T., Hagupit.
- Ficus aspera nota Blanco, ed. 1, 677; Ficus scabra Blanco ed. 2, 471, non Forst. = Ficus nota (Blanco) Merrill, Govt. Lab. Publ. 17: 10. 1904.! Erroneously reduced by Villar to Ficus racemifera Roxb. T., Tibig.
- Ficus glomerata Blanco, ed. 1, 683; ed. 2, 475, non Roxb. = Ficus minahassæ Miq.! Erroneously considered by Villar to be identical with Roxburgh's species. T., Tibig na lalaqui. V., Haguimit, Aimit.
- Ficus laevigata Blanco, ed. 1, 682; ed. 2, 474 = Ficus variegata Blume.! Erroneously reduced by Villar to Ficus cuneata Miq. T., Tangisang bayauac.
- Ficus hauili Blanco, ed. 1, 684; Ficus laurifolia Blanco, ed. 2, 475. Apparently a distinct species. Reduced by Villar to Ficus leucopleura Blume, which is a synonym of Ficus leucantatoma Miq. A shrub or small tree. T., Hauili.

- * Ficus laccifera Blanco, ed. 1, 673; ed. 2, 468, non Roxb. Reduced by Villar to Ficus radiata Decne., which is also a synonym of Ficus leucantatoma Poir. Villar's reduction is probably an error. Blanco's specimens were from Cebu and Negros, said to be a very common shrub. V., Lagnob.
- Ficus pseudopalma Blanco, ed. 1, 680; ed. 2, 473. A distinct species.! This species is not listed in Index Kewensis. T., Niogniogan.
- Morus tinctoria Blanco, ed. 1, 404, non Linn.; Broussonetia tinctoria Blanco, ed. 2, 488, non Spreng. = Cudrania obovata Trec.! T., Tahid labuyo. I., Talolong.
- Artocarpus incisa Linn. f., Blanco, ed. 1, 668; ed. 2, 465.! The earliest name for this species is Artocarpus communis Forst. T., V., P., Antipolo, Tipolo.
- Artocarpus rima Blanco, ed. 1, 671; ed. 2, 467. Reduced by Villar to Artocarpus incisa Linn. f., var. laevis Miq. From Blanco's description certainly not this variety. A form or variety of the preceding species.! T., Rima. V., Colo. C., Ogob. ?
- Artocarpus camansi Blanco, ed. 1, 670; ed. 2, 467. A form or variety of Artocarpus incisa Linn. f.! T., V., Dalangain, Dolongain, Camongsi. I., Pacac.
- Artocarpus odoratissima Blanco, ed. 1, 671; ed. 2, 467. Certainly closely related to Artocarpus incisa Linn. f., and probably only a variety of that species. T., Loloi.
- Artocarpus maxima Blanco, ed. 1, 669; Artocarpus integrifolia Linn. f., Blanco, ed. 2, 466. The latter name is correct. T., Nangca.
- Artocarpus ovata Blanco, ed. 1, 666; ed. 2, 464, non Noronha = Artocarpus cumingiana Trec.! T., Anobing, Anobing. P., Anobion.
- Artocarpus lamellosa Blanco, ed. 1, 667; ed. 2, 465. Reduced by Villar to Artocarpus nitida Trec., but Blanco's name is the older and should be retained. T., Anobing caguiosing, Anobling.
- Procris violacea Blanco, ed. 1, 706; ed. 2, 490 = Conocephalus violaceus (Blanco) (Conocephalus ovatus Trec.!). Erroneously reduced by Villar to the related Malayan species, Conocephalus suaveolens Blume. T., Lagna.
- * Procris erecta Blanco, ed. 1, 707; ed. 2, 490. Reduced by Villar to Conocephalus, as a distinct species, C. erectus (Blanco) F.-Vill. Known only from Blanco's description. T., Hanopol.
- Urtica sessiliflora Blanco, ed. 1, 696, non Swartz; Urtica capitata?

 Blanco, ed. 2, 483, non L. = Fleurya interrupta Gaud.! T., Lipang
 Castila. V., Langala, Damoro, Dalamo, Daodaua. P., Lopa.
- Urtica umbellata Blanco, ed. 1, 696, non Bory; Urtica ferox Blanco, ed. 2, 484, non Forst. = Laportea gaudichaudiana Wedd.! T., V., Lipa, Lipai, Lingaton, Apariagua. P., Lipang doton.
- * Dorstenia pubescens Blanco, ed. 1, 692; ed. 2, 481, non Forst. Reduced by Villar to *Elatostema obtusum* Wedd., which is probably an error. Certainly, however, a species of *Elatostema*.

- Urtica nivea Linn., Blanco, ed. 1, 697; ed. 2, 484 = Boehmeria nivea (Linn.) Hook. et Arn.! I., Ig., Amarai, Arimai, Labnis.
- Urtica villosa Blanco, ed. 1, 695; ed. 2, 483, non Salzm. Reduced by Villar to *Pouzolzia indica* Gaud., which is probably correct.
- * Urtica japonica Blanco, ed. 1, 684; ed. 2, 482, non Linn. Reduced by Villar to *Pouzolzia viminea* Wedd., which is probably an error. Probably, however, a species of *Pouzolzia*.
- Urtica baccifera Blanco, ed. 1, 695, non Linn.; Urtica arborescens Blanco, ed. 2, 483, non Poir. = Pipturus asper Wedd.! T., Dalonot. V., Dalonotan, Handaramai, Hindalamai. P., Dalonot.

JUGLANDACEÆ.

Gyrocarpus pendulos Blanco, ed. 2, 55 = Engelhardtia spicata Blume.! T., Pingol.

CASUARINACEÆ.

Casuarina equisetifolia Forst., Blanco, ed. 1, 661; ed. 2, 460. Certainly this species, although Villar refers it to Casuarina sumatrana Jungh. T., Agoho. V., Agoho, Malabohoc. P., Agoho. I., Aro, Caro, Agoo.

CUPULIFEREÆ.

- Quercus molucca Blanco, ed. 1, 726, non Rumph.; Quercus concentrica Blanco, ed. 2, 502, non Lour. = Quercus llanosii A. DC.! T., Alayan, Hayopag.
- Quercus glabra Blanco, ed. 1, 727, non Thunb. Quercus ovalis Blanco, ed. 2, 502. A distinct species; the second name should be retained. Quercus blancoi A. DC., is a synonym. T., Macabingao, Mangasiriqui, Hayopag.
- * Quercus cooperta Blanco, ed. 2, 503. Reduced by Villar to Castanopsis costata A. DC., probably, however, some species of Quercus. Blanco's specimens were from Angat.
- * Quercus cerris Blanco, ed. 1, 727; ed. 2, 503. Reduced by Villar to Castanopsis javanica A. DC., probably, however, some species of Quercus. Blanco's specimens were from Angat.
- Fagus philippinensis Blanco, ed. 2, 503 = Castanopsis philippinensis (Blanco) Vidal.! Erroneously referred by Villar to Castanopsis sumatrana A. DC. T., Tacatac, Talacatac, Lovian.

SALICACEÆ.

Salix azaolana Blanco, ed. 2, 539. Reduced by Villar to Salix tetrasperma Roxb., but apparently a distinct species, represented by Nos. 1987 and 2139 Forestry Bureau, Ahern's collector. T., Tiaun.

21896 - - 6

GNETACEÆ.

Gnetum gnemon Linn., Blanco, ed. 1, 747; ed. 2, 514.! T., Culiat.

Thoa pendula Blanco, ed. 1, 746; Thoa edulis Blanco, ed. 2, 514. Reduced by Villar to Gnetum scandens Roxb., but probably the same as Gnetum latifolium Blume. T., Culiat, Lamparahan, Biasbias, Taval, Malaitmo, Bangal, Itmong ouac.

CONIFEREÆ.

Agathis Ioranthifolia Blanco, ed. 2, 528, non Salisb. = Agathis philippinensis Warb.!

Pinus taeda Blanco, ed. 1, 767; ed. 2, 528, non Linn. = Pinus insularis Endl.! I., Saleng.

CYCADACEÆ.

Cycas circinalis Linn., Blanco, ed. 1, 745; ed. 2, 513.! T., Patúbo, Pitogo, Bitógo.

HYDROCHARIDEÆ.

Ottellia alismoides Pers., Blanco, ed. 1, 461; ed. 2, 321.! T., Calaboa.

Ottellia ensiformis Blanco, ed. 1, 460; ed. 2, 320. Retained by Naves as a distinct species, citing *Damasonium lancifolium* Presl., as a synonym. Certainly a form of the preceding species.! Blanco's specimens were from Pasig, Province of Rizal, flowering in March. T., *Espada*.

Vallisneria spaerocarpa Blanco, ed. 1, 780; ed. 2, 538 = Enhalus koenigii Rich.! T., Isai.

Vallisneria spiralis Linn., Blanco, ed. 1, 781; ed. 2, 538.! T., Cintascintasan.

SCITIMINEÆ.

Musa. Blanco describes eighteen varieties of the Banana, which, with the exception of four, are being placed by him under the species Musa paradisiaca Linn. He states that he saw in the Philippines fifty-seven varieties, and divides the various forms and species into two groups, those with thick-skinned fruit, and those with thin-skinned fruit. The many varieties described by Blanco are mostly entirely ignored by Schumann¹ in his recent monograph of the family, although many of them are very characteristic. It is quite impossible with our present information to correlate Blanco's varieties with those of other authors from the Malayan region, and accordingly the following list is copied from the two editions of this Flora, without reference to the identifications of Naves. The common native name for Bananas is Saguing, the Spanish names Platanos and Plantanos being also frequently used.

Musa paradisiaca Linn., Blanco, ed. 1, 239; ed. 2, 168.

¹ Engler, Pflanzenreich 4, 1900.

BANANAS WITH THICK-SKINNED FRUITS.

- * M. paradisiaca compressa Blanco, ed. 1, 240; ed. 2, 168. T., Bisco, Saba. Sp.-Fil., Obispo.
- * M. paradisiaca ternatensis Blanco, ed. 1, 243; ed. 2, 170. T., Ternate, Gloria.
- * M. paradisiaca lacatan Blanco, ed. 1, 243; ed. 2, 170. T., Lacatan.
- * M. paradisiaca pumila Blanco, ed. 1, 244; ed. 2, 171. T., Tampohin.
- * M. paradisiaca suaveolens Blanco, ed. 1, 244; ed. 2, 171. T., Bungulan.
- * M. paradisiaca magna Blanco, ed. 1, 244; ed. 2, 171. T., Tondoc.
- * M. paradisiaca subrubea Blanco, ed. 1, 245; ed. 2, 171.
- * M. paradisiaca violacea Blanco, ed. 1, 245; ed. 2, 171.
- * M. paradisiaca glaberrima Blanco, ed. 1, 245; ed. 2, 171. T., Galamai señora, Dinugoan.
- * M. paradisiaca maxima Blanco, ed. 1, 245; ed. 2, 171. T., Batavia, Matavia, Songai, Anuang.
- * M. paradisiaca longa Blanco, ed. 1, 245; ed. 2, 171. T., Quinauayan.
- * M. paradisiaca tombac Blanco, ed. 1, 246; ed. 2, 171. T., Tinumbaga, Goyoran.
- * M. paradisiaca ulnaris Blanco, ed. 1, 246; ed. 2, 172. This variety should be entirely discarded, as Blanco described it entirely from hearsay, and never saw specimens. "Los que han visto y comida el fruto, dicen que es del grueso de la pantorrilla, y que llega á tener hasta una braza de largo.!"
- M. troglodytarum Linn., Blanco, ed. 1, 246; ed. 2, 172 = Musa paradisiaca Linn., subsp. troglodytarum (Linn.) Baker. T., Botohan, Botoan.
- M. trogloditarum errans Blanco, ed. 1, 247; ed. 2, 172. Certainly a form or variety of Musa textilis Nec.! T., Saguing machin, Saguing na ligao.
- * M. troglodytarum dolioliformis Blanco, ed. 1, 855; ed. 2, 174.
- M. trogloditarum textoria Blanco, ed. 1, 247; ed. 2, 173 = Musa textilis Neé.! T., Abacá.

BANANAS WITH THIN-SKINNED FRUITS.

- * M. paradisiaca glauca Blanco, ed. 1, 250; ed. 2, 175. T., Bingticohol, Tinalong.
- * M. paradisiaca cinerea Blanco, ed. 1, 250; ed. 2, 175. Said by Blanco to have been introduced from India. T., Letondal.
- Amomum zingiber Linn., Blanco, ed. 1, 2; ed. 2, 2 = Zingiber officinale Rosc.! T., Luya. C., Laya.
- Amomum zerumbet Linn., Blanco, ed. 1, 2; ed. 2, 2 = Zingiber zerumbet (Linn.) Rosc.! T., Luyang osiu, Lampuyan.?
- Curcuma longa Linn., Blanco, ed. 1, 5; ed. 2, 4.! T., Dilao. V., Dulao, Calavaga, Quinamboy. P., Angai.

- *Costus luteus Blanco, ed. 1, 4; ed. 2, 3 = Curcuma xanthorrhiza Roxb. (?) T., Tamahilan, Tamohilan, Tamo, Tamocansi, Tumahila, Bolon. P., Tamo.
- * Costus nigricans Blanco, ed. 1, 3; ed. 2, 3. Reduced by Naves to Curcuma zerumbet Roxb., which is a synonym for Curcuma zedoaria Rosc. The latter name is probably the correct one for Blanco's species. T., Barac, Lampuyang.
- Kaempferia rotunda Blanco, ed. 1, 5; ed. 2, 4, non Linn. Reduced by Naves to Kaempferia galanga Linn., var. Probably a form of this species. T., Dusog, Duso, Dusod, Guisol. P., Cusol, Guisol, Dusol.
- * Amomum echinatum Willd., Blanco, ed. 2, 3. Reduced by Naves to Amomum aculeatum Roxb., var. majus. This may be correct. Blanco's specimens were from Angat, Province of Bulacan.
- Renealmia gracilis Blanco, ed. 1, 1; Renealmia exaltata Blanco, ed. 2, 1, non Linn. = Alpinia elegans (Presl) K. Sch. (Kolowratia elegans Presl; Alpinia gracilis Rolfe). Erroneously reduced by Naves to Alpinia gigantea Blume. T., Tagbac. P., V., Catcatan, Catotang, Tagbac, Salbac, Bagombon.
- Maranta arundinacea Blanco, ed. 1, 7; ed. 2, 5, non Linn. = Donax arundastrum Lour.! T., Tagbactagbac, Bamban.
- Canna Indica Linn., Blanco, ed. 1, 6; ed. 2, 5.! T., Cacuentasan, Tiquis tiquis, Ticas, Ticas ticas.

ORCHIDACEÆ.

- * Epidendrum equitans Blanco, ed. 1, 645; ed. 2, 449. Reduced by Naves to *Eria aporoides* Lindl., but the identification is very doubtful. Said by Blanco to be common in the Tala Mountains, Province of Rizal or Bulacan.
- * Epidendrum lineare Blanco, ed. 1, 644; ed. 2, 449, non Linn. Reduced by Naves to *Cleisostoma ionosmum* Lindl., which may possibly be correct.
- * Orchis lanigera Blanco, ed. 1, 641; ed. 2, 446. Reduced by Villar to Aerides virens Lindl., which is probably an error. Flowering in July.
- Arethusa glutinosa Blanco, ed. 1, 641; ed. 2, 446. Reduced by Naves to Geodorum semicristatum Lindl., which is probably correct. T., Cebollas del monte, Fariqtan, Camaog.
- * Cypripedium bulbosum Blanco, ed. 1, 637; ed. 2, 444, non Linn. Reduced by Naves to *Eulophia sumatrana* Blume, which is probably an error. A small plant, flowering in July.
- * Dendrobium teretifolium Spreng. ? Blanco, ed. 2, 444; D. teres Blanco, ed. 1. 638. Naves considers that Blanco correctly interprets Sprengel's species, and reduces it to Luisa teretifolia Gaud. Blanco's specimens were from Angat, Province of Bulacan, flowering in January or March.
- Phalaenopsis amabilis Blume, Blanco, ed. 2, 592. Probably correctly determined by Blanco. T., Dapo.

- *Ophrys cernua Blanco, ed. 1, 639; ed. 2, 445. Reduced by Naves to Grammatophyllum scriptum Blume, which is probably an error. A terrestrial plant, growing near the sea, at Bauang, Province of Batangas, flowering in March.
- * Vanilla ovalis Blanco, ed. 2, 448; Epidendrum vanilla Blanco, ed. 1, 643, non L. Naves considers that Blanco's species is identical with Vanilla aromatica Sw., but this is probably an error. Blanco's specimens were from Majayjay, Province of Laguna.
- * Vanilla majaijensis Blanco, ed. 2, 593. Reduced by Naves to Vanilla planifolia Ait., which is probably an error. Probably the same as the species described by Blanco as Vanilla ovalis, whatever that may be. Blanco's specimens were from the same locality as the preceding species.
- *Thelymitra malintana Blanco, ed. 1, 462; ed. 2, 447. Reduced by Naves to *Habenaria trinervis* Wight., which is probably an error, at least as to the species. Said by Blanco to be very common in Malinta, near Manila, flowering in October.

IRIDACEÆ.

Antholyza meriana Blanco, ed. 1, 24; ed. 2, 18, non Linn. Reduced by Naves to Sisyrinchium palmifolium Linn., an American species, which may be correct. Blanco's specimens were from cultivated plants flowering in November. Sp.-Fil., Cebollas de Persia.

AMARYLLIDACEÆ.

- Gethyllis acaulis Blanco, ed. 1, 260; ed. 2, 181. Reduced by Naves to *Hypoxis franquevillei* Miq. = *Hypoxis flava* Lour., which is certainly correct. Blanco's specimens were from Malinta, near Manila, flowering in October.
- Agave americana Blanco, ed. 1, 258; ed. 2, 180 = Agave rigida Mill. var.! T., Magui.
- Polianthes tuberosa Linn., Blanco, ed. 1, 259; ed. 2, 181. Correctly identified by Blanco, a Mexican species introduced and cultivated, flowering in May. Sp.-Fil., Azucena.
- Crinum giganteum Blanco, ed. 2, 175, non Andr.; Haemanthus pubescens Blanco, ed. 1, 253. Reduced by Naves to Crinum asiaticum Linn., which may be correct. T., Bacong.
- Crinum asiaticum Blanco, ed. 1, 251; ed. 2, 175, non Linn. Reduced by Naves to *Crinum gracile* E. Meyer, which is probably correct. Blanco's specimens were from Mandaloyon, near Manila, flowering in July.
- * Amaryllis atamasco Blanco ed. 1, 254; ed. 2, 178, non Linn. Reduced by Naves to *Hippeastrum miniatum* Herb., an American species, which may be correct, although Blanco does not state whether or not the plant was cultivated. Flowering in February. T., *Bacong, Talicoran*.

- Pancratium amboinense Linn., ? Blanco, ed. 1, 252; ed. 2, 177. Reduced by Naves to Eurycles amboinensis Herb. = Eurycles sylvestris Salisb., which may be correct. Common along the seashore, flowering in April. T., Catongal. V., Abur, Panabor, Catangal, Talaonor, Tonuar, Dausum.
- * Pancratium maritimum Blanco, ed. 1, 252; ed. 2, 177, non Linn.? The genus is probably correct, but the species is very doubtful. Pancratium maritimum Linn., is an American species. Blanco's specimens were from the Province of Batangas, there called Cebollas del Monte, from which it is evident that the plant that Blanco describes was probably from the forests, and not from the seashore.
- Pancratium zeylanicum Linn., Blanco, ed. 1, 253; ed. 2, 177. Probably correctly identified by Blanco. Flowering in April. T., Lunas, Catongal. Sp.-Fil., Rosas de Persia.
- * Pancratium illyricum Blanco, ed. 1, 251; ed. 2, 176, non Linn. Reduced by Naves to *Hymenocallis adnata* Herb., an American species which may be correct, although Blanco does not state whether or not his material was from cultivated specimens. Flowering in June. T., Tagalhi.

BROMELIACEÆ.

Bromelia ananas Linn., Blanco, ed. 1, 230; ed. 2, 162 = Ananassa sativa Lindl.! Sp. Fil., Piña.

TACCACEÆ.

- Tacca pinnatifida Forst., Blanco, ed. 2, 182; Tacca gaogao Blanco, ed. 1, 262, 856. The first name is correct. Z., Tambobon. I., Panarien. T., Gaogao.
- Tacca vesicaria? Blanco, ed. 1, 261; T. palmata Blanco, ed. 2, 182.
 Considered by Naves to be identical with Tacca palmata Blume.
 Blanco's specimens were from Malinta, near Manila, flowering in July. V., Magsaloro. Sp.-Fil., Corazon de Angel.

DIOSCOREACEÆ.

- * Dioscorea divaricata Blanco, ed. 1, 797; ed. 2, 550. Reduced by Naves to Dioscorea batatas Decne., but even if correctly referred, Blanco's name is the older. A species flowering in October, the tubers very deep in the ground. T., Paquit, Cobag. V., Baliacag. I., Dulian.
- Dioscorea alata Linn., Blanco, ed. 1, 799; ed. 2, 550.! T., Ubi. V., Quinampai.
- Dioscorea pentaphylla Linn., Blanco, ed. 1, 802; ed. 2, 552.! T., Limalima.
- Dioscorea papillaris Blanco, ed. 1, 801; ed. 2, 552. Reduced by Naves to Dioscorea sativa Linn., which is probably correct. T., Tongo, Quiroi.
- Dioscorea triphylla Blanco, ed. 1, 799; ed. 2, 551, non Linn = Dioscorea daemona Blume.! Reduced by Naves to Dioscorea hirsuta Blume. T., Nami. V., Corot, Gayos. P., Calut.

Dioscorea tugui Blanco, ed. 1, 800; Dioscorea sativa Blanco, ed. 2, 551, non Linn. = Dioscorea fasciculata Roxb.! T., Toqui.

LILIACEÆ.

- Smilax pseudochina Blanco, ed. 1, 795; ed. 2, 548, non Linn. = Smilax bracteata Presl, for most part, although Naves considers that Blanco's description also applies in part to Smilax china, S. laevis and Heterosmilax bornensis A. DC. T., Obat, Hampas tagbalang, Camagsa obat. V., Banag, Ubiubihan, Ronas. P., Barag. Sp.-Fil., Zarzaparrilla.
- * Smilax fistulosa Blanco, ed. 1, 796; ed. 2, 549. Reduced by Naves to Smilax indica Vitm., which may be correct. Blanco's description is very imperfect.
- Smilax latifolia Blanco, ed. 2, 548. Reduced by Naves to Smilax macrophylla Roxb., which may be correct. T., Sipit olang. V., Ronas.
- * Smilax divaricata Blanco, ed. 1, 795; ed. 2, 548. Reduced by Naves to Dioscorea myriantha Kunth. From Blanco's description the species may be either a Smilax or a Dioscorea, but from the native names it is apparently a species of Smilax. T., Obat, Camagsa obat, Hampas tagbalang or tigbalang.
- Flagellaria indica Linn., Blanco, ed. 2, 196.! T., Balinguai.
- Dracaena terminalis Linn., Blanco, ed. 1, 263; ed. 2, 183 = Cordyline terminalis (Linn.) Kunth.! T., Varas hari, Varas ni Jesé, Toncod obispo.
- Aloe humilis Blanco, ed. 1, 256; ed. 2, 179, non? Linn. Reduced by Villar to Aloe barbadensis Mill. A cultivated species. T., Sabila. V., Dilang boaya.
- Allium tricoccum Blanco, ed. 1, 239; ed. 2, 167, non Ait. Reduced by Villar to Allium uliginosum Don., which is cited by Hooker as a synonym of Allium tuberosum Roxb. T., Cuchai. V., Ganda.

PONTEDERIACEÆ.

Pontederia vaginalis Blanco, ed. 1, 255; ed. 2, 178, non Linn. = Monochoria hastaefolia Presl.! T., Calaboa.

COMMELINACEÆ.

- Lechea minor Blanco, ed. 1, 52; ed. 2, 35, non Linn. = Pollia sorzogonensis Presl.! V., Lohod lohod.
- Commelyna polygama Blanco, ed. 1, 25; ed. 2, 18, non Roth. = Commelina benghalensis Linn.! T., Alichangon. V., Sabilao. P., Biasbias.
- Commelyna cristata Linn., Blanco, ed. 1, 231; ed. 2, 163 = Cyanotis cristata (Linn.) R. et S.!
- Tradescantia geniculata Blanco, ed. 1, 232, non Jacq. = Floscopa scandens

 Lour.! This species is excluded from the second edition.
- Tradescantia discolor Smith, Blanco, ed. 1, 232; ed. 2, 163 = Rhaeo discolor Hance.!

PALMÆ.

- Calamus usitatus Blanco, ed. 1, 265; ed. 2, 185 = Daemonorops gaudichaudii Mart.! Erroneously reduced by Naves in part to Daemonorops rumphii Mart., and in part to Calamus pisicarpus Blume, species that do not extend to the Philippines. T., Yantoc, Oayi.
- Calamus gracilis Blanco, ed. 1, 267; ed. 2, 186, non Roxb. = Calamus blancoi Kunth.! Erroneously reduced by Naves to Calamus buroensis Mart., a species that does not extend to the Philippines. T., Talola.
- Calamus mollis Blanco, ed. 1, 664; ed. 2, 184. A good species; Calamus haenkeanus Mart., is a synonym. T., Dit-an.
- Calamus maximus Blanco, ed. 1, 266; ed. 2, 185. This is a distinct species (No. 1893 Merrill). It certainly is not a synonym of Calamus albus Pers., to which Naves reduces it, as the latter species is not found in the Philippines. T., Palasan. P., Labni. V., Parasan, Calapi. C., Labnit, Calapi.
- Areca catechu Linn., Blanco, ed. 1, 714; ed. 2, 494.! T., V., Bonga. P., Luyos.
- Areca catechu humilis Blanco, 11. cc., may be a valid variety. It is reduced by Naves to Areca catechu Linn., var. pumila Miq. T., Mangipod, Sacsic.
- Caryota urens Blanco, ed. 1, 740; ed. 2, 510, non Linn. = Caryota cumingii Lodd.! T., Taquipan, Taguipan.
- Caryota palindan Blanco, ed. 2, 513 = Orania Palindan (Blanco) (Orania philippinensis Scheff.). Erroneously reduced by Naves to Orania regalis Zipp. T., Palindan, Barangoi.
- Caryota tremula Blanco, ed. 1, 744; ed. 2, 512. Reduced by Naves to Wallichia tremula, which may be correct. T., Dumayaca.
- Caryota onusta Blanco, ed. 1, 741; ed. 2, 511 = Arenga saccharifera Lab.! T., Cauon, Iroc, Pugahan.
- Corypha umbraculifera Linn., Blanco, ed. 1, 228; ed. 2, 160.! T., B., Buri, Buli.
- Corypha minor Blanco, ed. 1, 229; ed. 2, 161, non Jacq. = Livistona merrillii Beccari.! Erroneously reduced by Naves in part to Livistona rotundifolia Mart., and in part to L. papuana Beccari, neither species being known from the Philippines. V., T., Anahao. P., Anao, Bolong ulong.
- Cocos nuclfera Linn., Blanco, ed. 1, 716; ed. 2, 495. T., I., C., Niog. P., Ngongot.
- Cocos mammilaris Blanco, ed. 1, 722; ed. 2, 499. A form or variety of the preceding species reduced by Naves to the var. lansiformis Miq. T., Adiavan.
- Nipa littoralis Blanco, ed. 1, 662; Nipa fructicans Wurmb., Blanco, ed. 2, 461. The second name is correct. T., Nipa, Sasa.

PANDANACEÆ.

- Pandanus spiralis Blanco, ed. 1, 777; ed. 2, 535 = Pandanus tectorius Soland.! T., Pandan.
- Pandanus gracilis Blanco, ed. 1, 778; ed. 2, 536. Erroneously reduced by Naves to *Pandanus humilis* Rumph., a species that does not extend to the Philippines. A species closely related to, if not identical with *Pandanus whitfordii* Merrill, although Blanco describes his species as unbranched. Blanco's specimens were from the Tala Mountains, Province of Rizal or Bulacan, flowering in May.
- Pandanus malatensis Blanco, ed. 2, 536. Reduced by Naves to Pandanus littoralis Jungh. Certainly a form or variety of Pandanus tectorius Soland.! Blanco's specimens were from Malate, a suburb of Manila, flowering in November.
- * Pandanus exaltatus Blanco, ed. 1, 778; ed. 2, 536. Erroneously reduced by Villar to Pandanus fascicularis Lam. Blanco evidently includes two species in his description, one form growing in the mountains, the other on the seashore. The latter is certainly Pandanus tectorius Soland., what the former is can not be determined from Blanco's description. T., Alas-as. In the Tala Mountains, Carahumai.
- * Pandanus sabotan Blanco, ed. 1, 779; ed. 2, 537. Erroneously reduced by Naves to *Pandanus dubius* Spreng. Certainly a distinct species, but known only from Blanco's description. Blanco's specimens were from the Province of Laguna. T., Sabotan.
- Pandanus radicans Blanco, ed. 1, 780; ed. 2, 537. Reduced by Naves to Pandanus bagea Miq., which is a synonym of Pandanus dubius Spreng., and which is probably correct. Pandanus dubius has recently been discovered in the southern Philippines. Blanco's specimens were from Leyte. V., Olango.
- * Pandanus inermis Blanco, ed. 2, 537, non Roxb. Erroneously reduced by Naves to *Pandanus moschatus* Rumph. Probably a good species, but Blanco describes the leaves only. A species from the mountains near Angat, Province of Bulacan, the leaves without spines.
- *Tillandsia pseudo-ananas Blanco, ed. 1, 853; ed. 2, 162. Reduced by Naves to Freycinetia insignis Blume, a species that does not extend to the Philippines. Probably a species of this genus, but it is impossible to determine which from Blanco's description. Blanco's specimens were from Angat, Province of Bulacan, flowering in August.

ARACEÆ.

* Batis hermaphrodita Blanco, ed. 1, 791; ed. 2, 544. Reduced by Naves to *Pothos chapelieri* Schott., which is certainly an error, as that species is known only from Madagascar. The specimen of Vidal's collection, cited by Naves, is *Pothos scandens* Linn. Blanco's specimens were from Bauang, Province of Batangas, a scandent plant, flowering in December. T., Apís.

- Pothos pinnata Blanco, ed. 1, 646; ed. 2, 450, non Linn. Reduced by Naves to Epipremnum medium Engl., which is probably correct. Flowering in February. T., Tibatib, Tampin banal, Malapacpac balavay. V., Bisaco, Tirbatib, Dibatib, Balicupcup, Daila, Garban, Tacolin, Horog. I., Amolong.
- Arum decurrens Blanco, ed. 1, 656; ed. 2, 457 = Amorphophallus campanulatus Blume.! T., Apon, Pongapon. I., Carot, Corot.
- Calla gaby Blanco, ed. 1, 659; Caladium esculentum Blanco, ed. 2, 459 = Colocasia antiquorum Schott.! var. T., V., P., Gaby, Gávay, Lagvay, Dagmay, Gandus.
- Calla badian Blanco, ed. 1, 658. Reduced by Villar to Alocasia indica Schott. var. variegata Engl., which is certainly an error. Blanco's species is certainly a form of the preceding species.! Included in the second edition only in a note following Caladium esculentum. T., V., Galiang, Biga, Badiang.
- Calla maxima Blanco, ed. 1, 658; Arum grandifolium Blanco, ed. 2, 458 = Alocasia indica Schott.! T., V., P., Biga.
- Arisaema polyphylla Blanco, ed. 1, 659; Caladium? digitatum Blanco, ed. 2, 459 = Arisaema polyphylla (Blanco) (Arisaema cumingii, Schott.!). Blanco's specimens were from Agoo, Province of Union.
- Arum divaricatum Linn., Blanco, ed. 1, 657; ed. 2, 458. Reduced by Naves to *Typhonum divaricatum* Decaisne, which is probably correct. T., Gabigabihan, Gabingouac.
- Pistia stratiotes Linn., Blanco, ed. 1, 651; ed. 2, 454.! T., Quiapo. I., Loloan.

LEMNACEÆ.

Lemna gibba Blanco, ed. 1, 672; ed. 2, 468, non Linn. Reduced by Naves to Lemna oligorrhiza Kurz, which may be correct. T., Inalai.

TYPHACEÆ.

Typha angustifolia Linn., Blanco, ed. 1, 687; ed. 2, 477 = Typha angustifolia Linn., subsp. javanica.! T., Balangot.

NAIADACEÆ.

* Najas palustris Blanco, ed. 1, 660; Najas lobata Blanco, ed. 2, 459. Reduced by Naves to *Naias tenuifolia* R. Br., which is probably an error. Probably the same as *Naias minor* All. A plant growing in stagnant fresh water, flowering in March.

CYPERACEÆ.

- Kyllinga triceps Blanco, ed. 1, 34; ed. 2, 23, non Vahl. = Kyllinga monocephala Rottb.! T., Muthang, Anuang.
- *Cyperus difformis Blanco, ed. 1, 32; ed. 2, 22, non Linn. Reduced by Naves to Cyperus ornatus R. Br., which is a synonym of Cyperus procerus Rottb., probably an error. T., Titio, Tiquio.

- Cyperus rotundus Linn., Blanco, ed. 1, 31; ed. 2, 21.! T. Mutha. V., Botobotones. P., Mota, Malaapolid, Sursur, Onoran, Casung, Omadiung, Galonalpas.
- * Carex tuberosa Blanco, ed. 1, 35; ed. 2, 24. Reduced by Naves to Eleocharis tuberosa Schult., which is probably an error, at least as to the species. Blanco's specimens were from Hagonoy, Province of Bulacan, the tubers edible. T., Potoc, Cibicibollasan, Apulid. Sp. Fil., Cabezas de Negrito.
- Carex glomerata Blanco, ed. 2, 24, non Thunb. Reduced by Villar to Fimbristylis ferruginea Vahl., which may be correct. Blanco's specimens were from Mandaloyon, near Manila, flowering in January. T., Apulid.
- Scirpus niloticus Blanco, ed. 1, 33; ed. 2, 23, non Gmel. = Fimbristylis miliacea Vahl.! T., Agor.

GRAMINEÆ.

- * Paspalum villosum Blanco, ed. 1, 40; ed. 2, 28, non Thunb. Reduced by Villar to Paspalum mollicomum Kunth, which is a synonym of Panicum muticum Forsk. Blanco's description hardly applies to the latter species, which, moreover, is not definitely known from the Philippines. T., Parag-is.
- Panicum miliaceum Blanco, ed. 1, 39; ed. 2, 28, non Linn. = Chaetochloa italica (Linn.) Scribn.! (Setaria italica Kunth.) T., Dava. I., Bucacao.
- Cenchrus hexaflorus Blanco, ed. 1, 36; ed. 2, 24 = Pennisetum macrostachyum Brongn.! Reduced by Villar to Pennisetum nigricans Miq., which is a synonym of Pennisetum compressum R. Br. Blanco's specimens were from the Province of Batangas, flowering in June. T., Aguingai.
- Stipa spinifex Linn., Blanco, ed. 1, 46; ed. 2, 31 = Spinifex squarrosus Linn.!
- Spinifex squarrosus Linn., Blanco, ed. 1, 46; ed. 2, 31.!
- Ægilops fluviatilis Blanco, ed. 1, 47; ed. 2, 32. Reduced by Villar to Rottboellia muricata Retz., which equals Eremochloa muricata (Retz.) Hack. Blanco's species probably is the same as Manisuris exaltata (Linn. f.) O. Ktze. T., Timsim.
- Zea mays Linn., Blanco, ed. 1, 686; ed. 2, 476.! Sp.-Fil., Maiz.
- Coix lachryma Linn., Blanco, ed. 1, 688; ed. 2, 478 = Coix lachryma-jobi Linn.! T., Tigbi.
- Andropogon contortus Linn., Blanco, ed. 1, 38; ed. 2, 26 = Heteropogon contortus (Linn.) R. et S.!
- * Andropogon ramosus Blanco, ed. 1, 37; ed. 2, 25, non Forsk. Reduced by Villar to *Ischaemum ciliare* Retz., which is certainly an error, at least as to the species. A species growing in the rice paddies in November, in appearance somewhat resembling wheat. T., *Tinitrigo* Sp.-Fil., *Trigo*.

- *Andropogon schoenanthus Blanco, ed. 1, 39; ed. 2, 27, non Linn. Considered by Villar to have been correctly identified by Blanco, but it is very doubtful if Blanco describes the Linnean species. T., Salai, Tanglad, Paja de meca. V., Baliyoco.
- Andropogon nardus Blanco, ed. 1, 39; ed. 2, 27, non Linn = Andropogon squarrosus Linn.! Andropogon muricatus Retz., to which Villar reduces Blanco's species as a synonym of Andropogon squarrosus Linn. Sp. Fil., Raiz de mora, Yerva mora.
- Saccharum koenigii Blanco, ed. 1, 44; ed. 2, 30, non Retz. = Saccharum spontaneum Linn.! Erroneously reduced by Villar to Imperata arundinacea Cyr. T., V., Cogon. P., Ilib, Bálili. C., Cogon cogon.
- Rhaphis trivialis Lour., Blanco, ed. 1, 45; Andropogon aciculatus Retz., Blanco, ed. 2, 26 = Chrysopogon aciculatus (Retz.) Trin.! T., Amores secos, Marisicos, Mariscos.
- Holcus saccharatus Linn., Blanco, ed. 1, 47; ed. 2, 32 = Sorghum saccharatum (Linn.) Pers. var.! T., Batad.
- Saccharum officinarum Linn., Blanco, ed. 1, 42; ed. 2, 29.! T., Tubo. Sp.-Fil., Caña dulce.
- Anthistiria gigantea Cav., Blanco, ed. 1, 49; ed. 2, 33.!

SPECIES OF ORYZA GROWING IN WATER, LOWLAND RICE.

- Oryza sativa binamban Blanco, ed. 1, 273; ed. 2 = Oryza sativa Linn.! T., Binambang.
- Oryza sativa lamuyo Blanco, ed. 1, 273; ed. 2, 190 = a variety of the preceding species.! T., Lamuyo.
- Oryza sativa glutinosa Blanco, ed. 1, 273; ed. 2, 190. Retained by Villar as a distinct species, Oryza glutinosa Lour. Certainly a variety of Oryza sativa Linn. T., Malagquit.
- Oryza aristata Blanco, ed. 1, 274; ed. 2, 190. Reduced by Villar to Oryza montana Lour. Certainly a variety of O. sativa Linn.!

SPECIES OF ORYZA GROWING IN DRY PLACES, UPLAND RICE.

- Oryza sativa praecox Blanco, ed. 1, 274; ed. 2, 190. Retained by Villar as a distinct species, Oryza praecox Lour. Certainly a variety of O. sativa Linn.! T., Dumali.
- Oryza sativa quinanda Blanco, ed. 1, 274; ed. 2, 191 = Oryza sativa Linn., var.! T., Quinanda.
- Oryza sativa pilosa Blanco, ed. 1, 275; ed. 2, 191. Retained by Villar as a distinct species, Oryza latifolia Desv. Certainly a variety of Oryza sativa Linn.! T., Bolohan.
- Oryza sativa rubra Blanco, ed. 1, 275; ed. 2, 191 = Oryza sativa Linn. var. T., Malagquit.
- Oryza sativa violacea Blanco, ed. 1, 275; ed. 2, 191 = Oryza sativa Linn. var.! T., Tangi.

- Arundo tecta Blanco, ed. 1, 48; ed. 2, 33, non Walt. = Phragmites communis Trin.! Reduced by Villar to Phragmites roxburghii Steud., which is a synonym of Phragmites karka Trin. T., Tambo.
- Cyperus paniculatus Blanco, ed. 1, 32; ed. 2, 22, non aliorum = Eragrostis tenella R. et S.! T., Caliraorao.
- Bambus pungens Blanco, ed. 1, 270; Bambus arundo Blanco, ed. 2, 188. Reduced by Villar to *Bambusa arundinacea* Willd., which is probably correct. T., Cauayangtotoo.
- *Bambus lima Blanco, ed. 1, 271; ed. 2, 189. Reduced by Villar to Bambusa longinodis Miq., but without good reason. Plant ten to fifteen feet high. T., Anos.
- * Bambus textoria Blanco, ed. 1, 270; ed. 2, 188. Reduced by Villar to Gigantochloa ater Kurz, but without good reason. T., Calbang.
- Bambus diffusa Blanco, ed. 1, 269; ed. 2, 187 = DINOCHLOA DIFFUSA (Blanco). Reduced by Villar to Schizostachyum acutiflorum Munro. T., Osiu, Bocavi.
- * Bambus monogyna Blanco, ed, 1, 286; ed. 2, 187. Reduced by Villar to *Dendrocalamus strictus* Nees, which may be correct. T., *Cauayang quiling*.
- * Bambus mitis Blanco, ed. 1, 271; ed. 2, 188. Reduced by Villar to Dendrocalamus sericeus Munro, which is certainly an error. T., Tiauanac.
- * Bambus lumampao Blanco, ed. 1, 272; ed. 2, 189. Reduced by Villar to *Dendrocalamus membranaceus* Munro, which is probably an error. T., *Lumampao*, *Bocaui*.
- * Bambus levis Blanco, ed. 1, 272; ed. 2, 189. Reduced by Villar to Dendrocalamus flagellifer Munro, which is probably an error. T., Cauayang boo.

LYCOPODIACEÆ.

- *Lycopodium gnidioides Blanco, ed. 1, 824; ed. 2, 569, non Linn. Reduced by Villar to Lycopodium selago Linn., which is certainly an error. An epiphyte. T., Pacayomcom.
- * Lycopodium dichotomum Blanco, ed. 2, 570. Reduced by Villar to Lycopodium cernuum Linn., which is probably an error. Blanco's specimens were from plants cultivated in Manila.

MARSILEACEÆ.

- * Marsilea minuta Blanco, ed. 1, 834; ed. 2, 577, non? Linn. Considered by Villar to be the Linnean species, which is doubtful. Blanco's specimens were from Pasig, near Manila, and from the Province of Batangas and other places, March.
- Marsilea trifolia Blanco, ed. 1, 835; ed. 2, 577. Reduced by Villar to Pilularia globulifera Linn., which may be correct. January.

FILICES.

- * Myriotheca arborescens Blanco, ed. 1, 831; ed. 2, 575. Reduced by Villar to Cyathea integra J. Sm. If this is correct Blanco's name is the earlier. T., Calatondon.
- * Adiantum falcatum Blanco, ed. 1, 833; ed. 2, 576, non Sw. Reduced by Villar to *Lindsaya retusa* Mitten., which may be correct. Blanco's specimens were from Mandaloyon, near Manila.
- * Adiantum lancea Blanco, ed. 1, 833; ed. 2, 576, non aliorum. Reduced by Villar to *Lindsaya ensifolia* Swartz, which may be correct.
- Adiantum philippinense Blanco, ed. 1, 831; ed. 2, 575, non Linn. = Adiantum lunulatum Burm.! T., Culantrillo, Lamot lamotan, Gayoman manoc, Caicai. P., Culantrillo. I., Dalipaco.
- Adiantum tenuifolium Blanco, ed. 1, 832; ed. 2, 575. Reduced by Villar to a variety of the preceding species, which is probably correct.
- Adiantum lyratum Blanco, ed. 1, 832; ed. 2, 575. Reduced by Villar to Adiantum caudatum Linn., which may be correct.
- Coenopteris quadripinnata Blanco, ed. 1, 833; ed. 2, 576 = Onychium auratum Kaulf.! T., Pacong anuang, Diladila.
- * Pteris grandifolia Blanco, ed. 1, 829; ed. 2, 574. Reduced by Villar to *Pteris opaca* J. Sm., which may be correct. T., V., *Dagabas*, *Darias*, *Tagabas*.
- Naias obvoluta Blanco, ed. 2, 460 = Ceratopteris thalictroides Brongn.!
- * Allantodia pinnata Blanco, ed. 2, 571. Reduced by Villar to Asplenium brackenridgei Baker, which may be correct. Blanco's specimens were from Mandaloyon, near Manila, where he states that it is very common.
- Hemionitis incisa Blanco, ed. 1, 829; ed. 2, 574 = Asplenium esculentum Presl.! T., Paco.
- Pteris piloselloides Blanco, ed. 1, 830; ed. 2, 574. Reduced by Villar to Nephrolepis acuta Presl., which is probably correct.
- * Pteris trichomanoides Blanco, ed. 1, 830; ed. 2, 574. Reduced by Villar to Nephrodium ramosa Moore, which is certainly an error. A foot high, growing on roofs.
- * Polypodium serratum Blanco, ed. 1, 827; ed. 2, 572. Reduced by Villar to Nephrodium floccigera Moore, which may possibly be correct. Said by Blanco to be very common.
- * Pteris sinuata Blanco, ed. 1, 830; Polypodium adiantiforme Blanco, ed. 2, 573. Reduced by Villar to Nephrodium davallioides Kunze, which may be correct. A fern three feet high.
- Polypodium phymatodes Linn. Blanco, ed. 1, 827; ed. 2, 572.!
- Polypodium quercifolium Linn., Blanco, ed. 1, 827; ed. 2, 572.! T.,
 Pacpac lauin, Paipai amo. V., Cabacbun, Cabcab, Cabcaban. P., Gona,
 Tibatib. I., Capacapa.

- * Polypodium signatum Blanco, ed. 2, 572; Pteris signata Blanco, ed. 1, 830. Reduced by Villar to Polypodium albo-squamatum Blume, which must be considered as doubtful. Blanco's specimens were from Mandaloyon, near Manila.
- Ugena semihastata Blanco, ed. 1, 822; ed. 2, 568. Reduced by Villar to Lygodium dichotomum Swartz, which is probably correct. T., Nito.
- Ugena alba Blanco, ed. 1, 823; ed. 2, 569. Reduced by Villar to Lygodium scandens Swartz, which is probably correct. T., Nitong puti.
- Helminthostachys dulcis Kaulf, Blanco, ed. 2, 596 = Helminthostachys zeylanica Hook.! T., Tocorlanguit.
- * Achrostichum simplex Blanco, ed. 1, 826; ed. 2, 571. This species is not cited by Villar by name but from the page citations, it is the species he intends to refer to Achrostichum latifolium Swartz, which reference is very doubtful.
- * Achrostichum tripinnatum Blanco, ed. 1, 826; ed. 2, 571. Reduced by Villar to Achrostichum apiifolium Hook., which may be correct. Blanco's specimens were from Malinta, near Manila.
- Achrostichum lagolo Blanco, ed. 1, 826; ed. 2, 720 = Achrostichum aureum Linn.! T., Lagolo.
- Blechnum colubrinum Blanco, ed. 1, 834; ed. 2, 576 = OLEANDRA COL-UBRINA (Blanco) Copeland.! A species related to Oleandra neriiformis Cav., to which it was reduced by Villar. It is represented by No. 3238 Merrill. T., Lunas.

CELLULAR CRYPTOGAMS.

Villar makes no attempt to identify the few species of cellular cryptogams described by Blanco, but for completeness the list is here given.

- * Marchantia ed. 1, 835; ed. 2, 577. Generic description only.
- * Tremella, ed. 1, 837; ed. 2, 578. Generic description only.
- * Agaricus, ed. 1, 845; ed. 2, 583. Generic description only, which also includes Polyporus. I., Colatcolat.
- * Scierotium subterraneum Blanco, ed. 1, 845; ed. 2, 584. A fungus found in the nests of the anay, white ant.
- * Fucus prolifer Blanco, ed. 1, 838; ed. 2, 579.
- * Fucus denticulatus Blanco, ed. 1, 839; Fucus natans Blanco, ed. 2, 579.
- * Fücus gulaman Blanco, ed. 1, 839; Fucus edulis Blanco, ed. 2, 580. T., Gulaman.
- * Ulva umbilicalis Blanco, ed. 1, 852; ed. 2, 581.
- * Ulva compressa Blanco, ed. 1, 842; ed. 2, 581.
- * Ulva reticulata Blanco, ed. 1, 842; ed. 2, 582.
- * Ulva intestinalis Blanco, ed. 1, 842; ed. 2, 582.
- * Conferva littoralis Blanco, ed. 1, 843; ed. 2, 582.
- * Conferva setosa Blanco, ed. 1, 844; ed. 2, 582.

UNDETERMINED SPECIES.

- * Cynomorium philippinense Blanco, ed. 1, 665; ed. 2, 464. Blanco's specimens were from Cebu. V., T., Capulao.
- * Erotium lanigerum Blanco, ed. 1, 440; ed. 2, 307. Blanco's specimens were from Cebu. V., Hagonoi sa lasang.
- * Kleinhovia serrata Blanco, ed. 1, 653; ed. 2, 456. Blanco's specimens were from Pasig, Province of Rizal, a shrub or small tree, flowering in September.
- *Legnotis lanceolata Blanco, ed. 1, 445. This species is not listed in Index Kewensis, and I have been unable to find the species in the second or third editions of the Flora de Filipinas or in the Novissima Appendix. Blanco's specimens were from Cebu, a tree flowering in February. V., Barit.
- * Nama jamalcensis Blanco, ed. 1, 211; ed. 2, 148, non Linn. Perhaps Blanco's material was from some cultivated species.

INDEX TO NATIVE NAMES.

•	Page.		Page.
Abacá	83	Alocasoc	69
Abangabang	33, 66	Alodig	78
Abilo	30	Alpai	35
Abur	86	Alum	78
Acapulco	41	Alupai	35
Achiote	18	Ајуоруор	67
Acle	42	Amadiong	91
Adaan	42	Amaga	
Adelfa	59	Amarai	81
Adgao	68	Amargoso	49
Adiavan	. 88	Amarillo	56
Agaiyangyang	38	Amoguis	36
Agapanga	31	Amolong	90
Agboligan	68	Amores secos	92
Agoho	•	Ampalaya	49
Agoioi	78	Amuyon	17
Agonoi	55	Anabo	23
Agoo	81	Anabon	25
Agor	91	Anagalti	30
Agosos	79	Anagap	43
Aguason	61	Anagatli	30
Aguingai		Anahao	88
Aguiu	30	Ananangtang	30
Ahihiro		Ananaplas	
Aimit		Anao	88
Alagao		Angai	83 43
Alambihor		Angelico	71
Alangilang		Angud	29
Alangilang sonson		Aniatan	25 25
Alangitngit		Anibiong	25 25
Alanguit		Anii	39
Alayan		Anilao	26
Albaricoques		Anis cahoi	73
Alcaparras		Anis moscada	73
Aldonises		Anitap	42
Alibanban		Anito	69
Alibihil		Anobing	80
Alibun		Anobing caguiosing	80
Alibutbut		Anobion	80
Alichangon		Anobling	
Alim		Anobran	68
Alingaro		Anonang	62
Alipai		Anonas	16
Aliparo		Anos	93
Alipata		Ansohan	
218967		97	

	Pa	ge.		. Pa	ıge.
Anten		30	Bahai		39
Antipolo		80	Baino		17
Antolang		67	Bait		75
Antolangan		23	Balabalanoyan	17.	18
Anuang	83.	90	Balabalayan		68
Apalaya		49	Balacat		33
Apalia		49	Balacbac		48
Apalit		40	Baladre		59
Apariagua		80	Balago		19
Apatot		54	Balaibai		69
Apis		89	Balai-lamoc		53
Apiton		21			18
-	-1 17		Balai namoc		
Apoiapoyan	11,		Balangot		90
Apon		90	Balanoi		69
Aposotis		71	Balanti		77
Apulid		91	Balao		21
Arandon		74	Balasbas		67
Araña		18	Balasbas malomay		67
Ardotag		55	Balasin		17
Arimai		81	Balatong		39
Aringit		60	Balatong aso		37
Aro		81	Balayon		41
Arodayday		62	Balayong		41
Arogangan		23	Baliacag		86
Aroma		42	Balibago		23
Aroyangyang		38	Balibanban		42
Arum		78	Balicbalic	40,	60
Asana		40	Balicupeup		90
As-as		54	Balictarin		69
Asiamo		29	Balilangouag	٠	77
Asis	•	79	Balili		92
Asuangai		69	Balimbin		27
Ataiatai		67	Balingayo		48
Ates		16	Balinguai		87
Ausiman		19	Balinhasa		36
Ayangao		42	Balitbitan	11	
Ayantoto	,	71	Baliti	ч1,	78
-		55	1		92
Ayapana	07		Baloyoco		
Ayo	21,		Balobar		45
Ayoban		55	Balocanad		76
Azucena		85	Balocas		29
Bacao			Balod		52
Bacauan	43,	~	Balogo		42
Bacong		85	Balongai		74
Bacugan		30	Balucanag		76
Badiang		90	Bamban		84
Badiara		69	Banaba		47
Badoc		55	Banabanalo	24,	73
Bagaboboi		68	Banag		87
Bagalangit		57	Banago	20,	23
Bagaolan		51	Banaibanai	15,	66
Bagarilao na itim		52	Banalo	23,	61
Bagarilat		52	Banaro		23
Bagasua		62	Banati		28
Baga tumbal		27	Bancal		52
Bagauac		68	Bancudo		54
Bagobago	20.	54	Bangal		82
Bagombon	•	84	Bangar		24
Baguilumban		76	Bangati		38

*	Pa	ge.		Page.
Bangbang		56	Bitlag	32
Bangcal		52	Bitnong	24
Bangcolon		34	Bitógo	82
Bangcoro		54	Bitongol	18
Banilac		44	Bitoon	46
Banilad		24	Bitung	46
Banilar		24	Boboa	31, 35
Banot		42	Boboi	24
Banquilin		76	Boboi gubat	23
Baquinbaquit		25	Bocaboc	56
Barac		84	Bocaui	93
Barag		87	Bocavi	93
Barangoi		88	Bognai	76
Barit	45		Bolagtob	64
Batad	10,	92	=	36
Batangbatang		17	Bolobtones	75
Batao		40		92
Batavia		83	Bolohan	
	00		Bolon	84
Baticulin	00,		Bolongcauit	31
Batino		59	Bolong eta	57
Batoan		36	Bolong tambal	30
Bato bato	63,		Bolong ulong	88
Batobatonis		75	Bonac	69
Bayabas		45	Bonga	88
Bayag cambing		41	Bonga bonga	71
Bayag usa		59	Bongalon	42
Bayangbayang		71	Bongliu	51
Bayaquiboc		54	Boocan	31
Bayati		17	Bosboron	56
Bayog		25	Boto	56
Bayogo		42	Botoan	83
Bebebabe		45	Botobotnes	91
Bella de noche		70	Botohan	83
Betis		57	Botolan	76
Biasbias 51,	82,	87	Botong	46
Bias pogo		47	Botongbotong	46
Biga		90	Botong gubat	78
Bignay pogo		76	Botong manoc	66
Bilang bilang		50	Bovi	75
Bilau		78	Boyocboyoc	49
Bilibi		62	Bucacao	91
Bilimbin		27	Buena vista	77
Bilogo		32	Bugayon	38
Binambang		92	Bugnai	76
Binbinguay		74	Bulac	24
Bingabing		78	Bulacan	63
Bingliu	30		Bulac cahoy	24
Bingticohol	00,	83	Bulac castilla	
Biniguasan		60	Bulac na monti	24
Binonga		78	Bulac na totoo	24
Binoyoc boyoc		49	Bulahan	31
Bintoco		27	Bulai	40
Binucao		20	Bulai lava	37
Biri		56		52
			Bulala	
Biringi		39	Bulaon	68
Bisaco		90	Buli	88
Bisco		83	Bungulan	83
Bitanhol		20	Buntot usa	25
Bitaog		20	Buquingan	71

· · · · · · · · · · · · · · · · · · ·		ь.		rage.
Buquiquit		38	Calayate	67
Buracan		63	Calbang	93
Buri		88	Caliantang	3 3
Busain		24	Calimbahin	45
Busilac	31,	77	Calinga	73
Buta		78	Calingad	73
Butabuta		78	Calingag	73
Buttingi		39	Calios	78
Buyayava		75	Calipayang	77
Buyo		72	Caliraorao	93
Buyochuyoc		49	Calisai	44
Buyo de anis		72	Calitcalit	33
Buyong		47	Calites	71
Cabacbun		94	Calobcob	45
Cabag		52	Calocatingan	25
Cabaicabai		40		26 26
		61	Calomata	28
Caballara			Calomata	
Caballero		41	Calondagui	75
Cabalte		26	Caloo	38
Cabatiti		33	Calumban	76
Cabcab		94	Calumpang	24
Cabcaban		94	Calumpangin	51
Cabcabron		55	Calumpang sa lati	31
Cabeza negrito		91	Calumpit	44
Cabilte		26	Calungai	36
Cablin		70	Calut	86
Cabog		16	Calutpamo	33
Cabogbog		76	Camagsa obat	87
Cabrag		39	Camagsa taquilis	36
Cachumba		56	Camaisa	77
Cacuentasan		84	Camalagui	4.1
Cadayohan		71	Camalungay	36
Cadlin		70	Camangi	37
Cadlom		70	Camanguianis	28
Café		54	Camantigui	27
Caguois		40	Camaog	84
Cahil		29	Ca Maria	56
Caicai		94	Camariang songsong	70
Calaad		17	Camatis	63
Calabazang bilog		50	Camias	27
Calaboa	82.		Camigang	62
Calacalamayan	,	17	Camiring	36
Calachuchi		59	Camit cabag	41
Calai		16	Camo camotihan	62
Calaicai		38	Camochiles	43
Calamansalai		30	Camongsi	80
Calamansanai		44	Camonsiles	43
Calamantao		76	Camoti	62
Calambibit		41	Camoting cahoy	77
Calamias		27	Campanelo	58
		40		
Calamismis			Campopot	
Calamondin		29	Camuning	28
Calangeang		51	Candaroma	73
Calantas	rn.	31	Candon	47
Calapi	o3,		Candong	47
Calapinai		35	Cangeong	62
Calasasi		59	Cangcong dapo	48
Calatondon		94	Caningag	16
Calavaga		83	Canlara	61

	Page.		Page.
Canomai	57	Cayanga	23
Canomoi	57	Cayanga rosa	23
Cansasaga	38	Cayangcan	51
Cantotai	54	Cayaocayao	59
Cantotan	58	Cayutana	27
Canubsuban	71	Cebollas del monte	84
Canucsuban	71	Cebollas de persia	85
Caña dulce	92	Chachachachahan	65, 67
Cañafistula	41	Chicos	57
Capacapa	94	Cibicibollasan	91
Capalcapal	60	Cinamomo	47
Capas	24	Cinco llagas	67
Capili	76	Cintascintasan	82
Capitcabag	41	Cobag	86
Capopot visaya	58	Cocong manoc	. 66
Capulao	96	Codamba	61
Carabaui	75	Cogon	92
Carachucha	59	Cogon cogon	92
Caragli	53	Colasiman	19
Carahumai	89	Colatcolat	95
Carambuaya	75	Coles maluco	70
Carampalit	50	Colis	47
	51		80
Carangeang	43	Colo	29
Carisquis	1	Colobot	40
Carlin	70	Colocanting	
Caro	81	Colocolog	69 69
Carot	90	Coloncogon	
Carucansoli	70	Colongcolong	74
Casabha	56	Colotan	23
Casindac	39	Colotan bilog	26
Casitas	41	Colotcolotan	23
Casla	76	Combarcombaran	69
Casopanguil	69	Comimpol	26
Casopanguil gubat	68	Comintan	44
Casoy	36	Condol	49
Castiocastiogan	23	Consuelda	75
Castoli	23	Copang	42
Casubha	56	Copcop	45
Casubhang aso	17	Corazon de Angel	86
Casung	91	Corot	
Catacataca	43	Cotingcotingan	62
Catalonga	61	Cuacuacohan	23
Catanda	-	Cuanton	71
Catandang aso	41	Cubatili	31
Catangal	86	Cubili	35
Catang catang	62	Cuchai	87
Catcatan	84	Cucubitan	49
Catiquis	34	Culantrillo	94
Catluen		Culasi	44
Catmon	15	Culatai	58
Catongal	86	Culiat	82
Catotang	84	Culit	54
Catuit	75	Cungi	27
Caturai	38	Cunti	64
Cauayang boo	93	Cupit cupit	68
Cauayang quiling	93	Curagda	
Cauayang totoo	93	Curribuetbuet	
Cauon	88	Cuscusipa	17
Cavilan	33	Cusiben	. 34

	Page.		Page.
Cusol	84	Diego de noche	70
Cuti	64	Digquit digquit	70
Dagabas	94	Diis	42
Dagamit	43	Diladila	94
Dagangdang	37	Dilang baca	50
Dagatan	65	Dilang boaya	87
Dagmay	90	Dilang butiqui	52
Daila	90	Dilang usa	55
	40	Dilan usa	62
Daitanag	16	Dilao	83
Dalaganum			
Dalagao	16	Dilivario	66
Dalaguit '	78	Dincalan	20
Dalamo	80	Dinglas 44,	
Dalandan	29	Dinuguan	83
Dalandon	68	Dirig	35
Dalangain	80	Dita	59
Dalanta	33	Dit-an	88
Dalaora	67	Docotdocot	71
Dalasa	44	Dolariu	66
Dalipaco	94	Doldol	24
Dal-lopauen	59	Dolongain	80
Dalondon	67	Dongon	24
Dalondong	57	Dool	36
Dalonot	81	Duat	46
Dalonotan	81	Duca	34
Dalugdug	41	Duclap	33
Dalupan	23	Duclitan	57
Dalupang		Ducum	23
Damonghiya		Duguan	73
Damora		Duha	24
Damoro	80	Duhat	46
Dampalit		Dulao	83
Dancagi		Dulauari	66
Dancalan		Dulian	86
Danglay		Dulitan	57
Danglin		Dumali	92
Dangling aso		Dumayaca	
Dao			37
Daodaua		Durang parang	84
		1	
Dapdap		Duso	
Dapo		Dusog	
Dapo sa cahoy		Dusol	
Dapo sa pajo		Escobang bilog	23
Daracan		Escobang haba	
Daraisig.		Espada	
Daraput		Fariqtan	
Dariangao		Flor de campana	
Darias		Flores	
Darigayos		Frijoles del Abra	
Daripay		Fruta	
Dauag		Gabigabihan	
Dausum		Gabing ouac	
Dava	91	Gabuen	55
Dayang	. 66	Gaby	. 90
Dayap	. 28	Galamai amo	. 51
Dayap na monti		Galamai señora	. 88
Decdec	. 32	Galiang	90
Dibatid		Galonalpas	
Dicuta maboloc		Gamagamatisan	64

	Page.		Page.
Gamot sa buni	41	Handaramai	81
Ganda	87	Hangor	71
Gandus	90	Hangot	71
Gaogao	86	Hanlilimocon	71
Gapasgapas	. 68	Hanopol	80
Garbanzos	39	Hapiton	21
Garban	90	Harangan	56
Gatasgatas	75	Haras	50
Gavai gavai	17	Harum	71
Gávay	90	Hauili	79
	94	1	81
Gayoman manoc	86	Hayopag	40
Gayos		Hicamas	
Gicos gicos	38	Higot balato	23
Gloria	83	Higuis manoc	55
Gogo	42	Himamao	31
Gogong bacay		Himbabao	78
Gogong bisaya		Hindalamai	81
Gogong toco	31	Hindurugo	73
Golandrina	75	Hindurugu	24
Gomamila	23	Hingongoto	75
Gona	94	Hinguiu	59
Goyoran	83	Hinlaylayon	62
Grana	71	Hintipalo	43
Granada	47	Hintotoor	41
Grano del parayso	17	Hitam	44
Guanabanos	17	Hojas moradas	67
Guicos guicos	36	Horog	90
Guijo	22	Hothot	49
Guilalas	70	Hoya cruz	66
Guinabang	78	Huampit	28
Guinguen	28	Huliganga	52
Guintenguinten			76
_	,55	Iba	
Guisan		Ibabao	41
Guisian		Ibaiban	36
Guisihan		Igasud	61
Guiso	22	Iginga	69
Guisol	84	Iguiu	30
Guisong dilao	22	Ilaibaquir	71
Gulaman	95	Ilib	92
Guling gulingab	23	Iloilo	31
Guyabas	45	Inalai	90
Guyongguyong	19	Inignin	48
Habas	39	Ipil	41
Habul	55	Iring	21
Hagonoi	55	Iroc	88
Hagonoi sa lasang	96	Isai	82
Hagorilis	71	Isioisio	79
Haguimit	79	Isis	79
Hagupit	79	Issio	79
Halahala	49	Itmo	72
Halon		Itmong ouac	82
Hamaraon	68	Labba labba	33
Hamitanag	24	Labni	88
Hamlibon	55	Labnis	81
	87		88
Hampas tagbalang		Labrit	
Hampas tigbalang	87	Labon	25
Hanambabao	36	Lacad bulan	55
Hanarion	78	Lacatan	83
Hanbabalos	52	Lacdanbulan	55

	rage.	· · · · · · · · · · · · · · · · · · ·	Lus	50.
Lactang	17	Lilitan		54
Lagayrai	62	Limalima	51,	86
Lagingi	33	Limang sugat		67
Lagini	33	Limoneito		28
Lagna	80	Limoneitos		28
Lago	56	Linatong anac		29
Lagnob	80	Linga		66
Lagolo	95	Lingalingahan		69
Lagpacum	64	Lingasina		78
Laguiolaguio	66	Lingaton		80
Lagundi	68	Lino		54
Lagunding gapang	68	Linog		56
Lagundi salasa	61	Lintang baguin		17
Lagvay	90	Linton gamai		29
Laiya	29	Lipa		80
Lalacdan	55	Līpai	38,	80
Lambayong	62	Lipang castila		80
Lamigpa	31	Lipang doton		80
Lamio	36	Lipata		78
Lamot lamotan	94	Litbit		72
Lamparahan	82	Litlit		72
Lampayong	62	Livas		42
Lampuyan	83	Loba		33
Lampuyang	84	Lobalob		75
Lamudio	50	Lobas		46
Lamuyo	92	Lobio		50
Langala	80	Locloquisin		61
Langas	36	Locoloco		69
Langil	42	Lohod lohod		87
Langis	66	Loloan		90
Lanigda	31	Loloi		80
Laniti	59	Lombayong		41
Lanotan	16	Lomonai		78
Lanotan itim	16	Longa		66
Lanoti	33	Lopa		80
Lansone	31	Lovian		81
Lansones	31	Lubilubi		
Lanta	17	Lucban		29
Lantin	70	Lugo		44
Lapnis na bolohan	23	Lumampao		93
Lapolapo		Lumban		76
Laroan		Lumboi		46
Lauaan	22	Lunas 27		
Lauan		Lupo		33
Lauas		Luya		83
Lavas		Luyang osiu		83
Laya		Luyos		88
Layo		Maasic		37
Layohan		Mabatang dicut		54
Lechias		Mabolo		58
Lengua de perro		Macabingao		81
Letondal		Macabuhay		17
Libai		Macahiya		
Libas		Macaisa		29
Libato		Macalalauang		69
Libun		Macalingag		73
Lictang		Macan		51
Ligas		Macapil		40
Ligasin	. 44	Maca sampaloc		41

	Pag	ge.		Page.
Macupa		45	Malimali	34
Madondon		53	Malis	23
Madre cacao		38	Malisa	72
Magatas		75	Malismalis	75
Magsaloro		86	Malongain	30
_		36		
Magtabig		1	Maluco	70
Magtambocao		39	Malungay	36
Magui		85	Malungit	36
Mahahiin		27	Malva rosa	26
Maisipaisa		28	Malvas de Castilla	23
Maiz		91	Mavindata	29
Malaanonan		22	Malayantoc	59
Malaapi		68	Malbas tagbalang	15
Malaapolid		91	Mamalis	
=		53		
Malaas-is			Mambog	
Malabachao		75	Mamei	57
Malabaguio		32	Mamin	72
Malabalugbug daguis		27	Mamitic	66
Malabangao		30	Mamon	72
Malabatoan '		20	Mampol	43, 52
Malabayabas		53	Mananantang	30
Malabochoe		20	Mana-naog	61
Malabohoc	74.	81	Mananquil	45
Malabonga	• -,	74	Mandalusa	67
		24	Manga	35
Malabulac			_	
Malabunot		24	Mangachapoi	22
Malacabuyao		29	Manga de anis	35
Malacadog		44	Mangadolong	38
Malacaguois		38	Manga-pico	35
Malacatmon		15	Mangasiriqui	81
Malaclac		56	Mangipod	88
Maladosos		67	Mani	38
Malagoso		50	Manimanihan	41
Malagquit		92	Manguit	38
Malaisis		78	Mantala	40
Malaitmo	78.		Manul	58
	10,		,	29
Malaiyo		36	Manungal	
Malamayana		69	Manzanilla	56
Malamolauin		28	Manzanitas	32
Malapaco		48	Marayapa	69
Malapacpac balavay		90	Margoso	49
Malapalitpit		34	Maricum	23
Malapao		21	Mariscos	92
Malapapaya		51	Marisicos	92
Malapatopat		53	Maropoto	23
Malapotocan		68	Marucum	23
Malaracopcop		45		42
	20		Marulinao	27
Malarayap			Matang arao	,
Malaruhat		46	Matang diablo	27
Malaruhat na pula		46	Matang olang	32
Malasaga		40	Matangpune	38
Malasampaga		74	Matavia	83
Malatibig		21	Mavindato	36
Malatinta		76	Mayana	69
Malatumbaga		45	Mayapis	21
Malaubai	,	72	Melindres	47
Malaubi		72	Melon daga	50
		69	1	49
Maliana			Melong ouac	
Malibago		23	Migi	31

	Page	е.		Pa	age.
Mil leguas	6	30	Pacapis		69
Minonga	7	18	Pacayomcom		93
Mocasisi	3	30	Pacayomcom Castila		41
Molauin	6	88	Paco		94
Molavin	28, 6	38	Pacong anuang		94
Molopolo	2	23	Pacopol		33
Molugoso	- 5	50	Pacpacbalang		25
Mongomongohan	4	11	Pacpac langao		38
Mongos	3	39	Pacpac lauin		94
Mopio	6	37	Pacupis		49
Morado	6	37	Pagatpat	47.	
Moradong maputi	6	37	Pagbut	,	53
Morera	7	8	Paipai amo		94
Mostaza		7	Paipaisi		70
Mota		1	Paja de meca		92
Mutha	-	1	Pajo		35
Muthang	-	00	Palac palac		57
Nabo	-	25	Paladpalad		66
Nacboligan	-	38	Palagarium		29
Naga	-	10	Palagium		29
Nami		36	Palapad		47
Nangca		30	Palasan		88
Narangitas		29	Palatpat		47
Narra		10	Paliavan		17
Nato			Palindan		88
Ngongot		88	Palis	99	
Nigui		31	Palla	40,	49
Nilad		3	Pal-lam		40
Nilar	_	3			16
Nino	_	54	Palo calai Palohambobocag		30
Niog		88	Palo Maria		20
Niogniogan			Palongpalongan		71
Nipa		88	Palo Santo	96	
Nipai		18	Palosapis	50,	22
Nitong puti		5	Palsahinhin		30
Nonoe		18	Palunai	.*	55
Noog noog		34	Pamalatanguen		53
Nopau		2	Pamalis		23
Oas		35	Pamitlain		20
Oayi		38	Pamitlatin		20
Obat		37	Pamuhat		68
Obispo		33	Pamulaclaquin		44
Ogob		30	Panabolong		56
Olango		39	Panabor		86
Olasamang aso		35	Panacla		70
Olasiman		9	Panampat		24
Onava		32	Panao		21
Onoran		1	Panaptun		67
Onti		34	Panarien		86
Oongon		4	Panayan		17
Оро		9	Pancudo		54
Oracion		0	Pandacaqui		59
Orayi		1	Pandan	44.	
Oregano			Pandapanda	,	30
Osiu		3	Pandaya		59
Ovai na panglao		5	Pangaguason		61
Paanbalivis		0	Pangangtolong		56
Pacac		30	Pangiavan		17
Pacalcal		8	Pangui		18
	•	- 1			

Pansipansi	70	Quilingiva	27
Pao	35	Quilites	71
Papaya	49	Quinalumpang	23
Papua	51	Quinamboy	83
Paquit	86	Quinampai	86
Parag-is	91	Quinanda	92
Parasan	88	Quinasaicasai	42
Paria	49	Quinauayan	83
Paronapin	24	Quindayohan	71
Pasao na bilog	26	Quiroi	86
Pasao na haba	26	Raguindi	
Pasiloag	30	Raiz de mora	92
	64	Rima	80
Pasitis		•	
Patai ood	62	Romero	70
Patani	39	Ronas	87
Pataning dagat	39	Rosas	41
Patola	49	Rosas de Persia	86
Patúbo	82	Saba	83
Payangpayang	38	Sabia	72
Payanguit	60	Sabila	87
Payapa	79	Sabilao	87
Pepinillo de San Gregorio	49	Sablut	73
Pepita	61	Sabotan	89
Pepita sa catbalogan	61	Sacsic	88
Pias	27	Saga	38
Pilani	30	Sagamamin	38
Pilapil	18	Sagmit	43
Pimienta	72	Sagnit	41
Pincapinca	66	Sagudan	20
Pincapincahan	66	Saguilala	77
Pingol	81	Saguing	
Pingol bato	50	Saguing machin	
Piña	86	Saguing na ligao	
Pipisic	57	Saguing saguing	61
Pirapit angin	33	Saguinsin	47
Piris	28	Salab	34
Pisa	30	Salago	74
Pisong	29	Salagsalag	49
Pissic	56	Salai	92
Pitogo	82	Salapao	
Platano	82	Salapong	
Platanos	82	Salaqui	
Poas	35	Salay cangay	
Pocotpocot	49		
		Salay marbar	
Polyagan	30	Salayong	
Pongapon	90	Salbac	
Ponoan	29	Saleng	
Poras	76	Salimpocot	
Potat	46, 51	Sallacapo	
Potoc	91	Sal-laday	. 37
Potocan	64	Salomagui	. 41
Pototan	44	Samac	. 78
Pugahan	88	Samat	. 72
' Puntas puntas	63	Sambac	
Puriquet		Sambon	
Pusopuso		Sambong gala	
Quiapo		Samilin	
Quibal		Sampac	
		1	
Quilamo	47	Sampaca	. 10

	Pa	ge.		Page.
Sampaga		58	Suma	17
Sampaga del monte		56	Sumag	55
Sampagang pougso		58	Sunting	41
Sampalagui		41	Suranga	27
Sampaloc		41	Susocoyoli	27
Sampang		33	Susoi	87
Sincamas		40	Susong damolag	16
Sandana	22.	- 1	Susosusoyan	37
Sandoc sandoc	,	50	Suspiros	70
San Francisco	67,	1	Sursur	91
Sangdigquit	٠.,	56	Suya	29
Sangilo		72	Taal	41
Sangue		40	Tabaco	64
-		17	Tabatabacohan	55
Sansao				
Sansaosansauan		17	Tabayag	49
Santa Ana		54	Tabigui	31
Santa Maria		56	Tabing	23
Santan		54	Tablon	78
Saog		72	Tabobog	49
Sapang		41	Tabog	29
Sapinit 23,			Tabogo	50
Sapinsapin	66,	67	Tabtabocol	62
Sapnit		41	Tabtabocol ti nuang	70
Sapolongan		22	Tac-an	70
Sasa		88	Tacatac	81
Sausanli		34	Taclang onac	20
Sayican	19,	75	Taclitan	30
Siac		78	Tacolin	90
Sibog		42	Tocor languit	95
Sibucao		41	Tacpo	54
Sigang dagat	55,	62	Tacpong dalaga	57
Silasila		40	Tacpus	54
Silipao		32	Tacurangan	23
Silisilihan	56.	67	Tadiang	32
Sinampaga		53	Tadiang anunang	53
Sincamas aso		39	Tagabas	94
Sintonis		29	Tagabili	55
Sipit olang		87	Tagactagac	67
Sirihuelas		36	Tagalhi	86
Sisiohan		75	Taganagtagua	17
Sitao	39.		Tagapi	36
Soal	00,	16	Tagarao	45
Sobsob		55	Tagaray	62
Solanga		27	Tagbac	84
Solasi		69		84
Solasolasihan		70	Tagbactagbac	66
		56	Taghilao	16
Solinao			Taghivalas	
Songai		83	Tagima	51 24
Sonting		41	Taglinao	
Soob cabayao		69	Taglocot	62
Sorogsorog		75	Tagolinao	56
Sorosoro		75	Taguilima	51
Sucao		17	Taguima	51
Suganda		69	Taguipan	
Suha		29	Tagung	
Sulbang		39	Tahid labuyo	
Subsuban		71	Tala	
Suliac daga		53	Talacatac	
Sulipa		53	Talactac	18

	Page.		Page.
Talailo	17	Tartarao	45
Talamponai	64	Tarumpalit	50
Talamponai na itim	64	Tatlong	32
Talanai	33	Tatumbulo	64
Talancao	56	Taval	82
Talang	57, 58	Tavatava	76
Talangtalang	73	Tavatavangsina	78
Talaonor	86		75
Talatala	64		64
Taliantar	54	Tayom	37
Talicnono	61		
Talicoran	85		37
Talicot	79		61
Talinas	33		37
Talingharap	70		67
Talinouno	35		
Talisai	44		93
	88		
Talola			81
Talolong	80		
Talong	64		79
Talong gubat	64	•	79
Talosan	25	Ticas	84
Taloto	24		84
Tamahilan	84		67
Tamauian	20		44
Tambalisa	40		91
Tambo	93	Tighiman	41
Tambobon	86	Tililes	71
Tamo	. 84	Timbagan	72
Tamocansi	84	Timbangantimban	72
Tamohilan	84	Timsim	91
Tampin banal	90	Tinaingui	35
Tampohin	83	Tinalong	83
Tampoi	45	Tinatinaan	76
Tanag	24	Tindalo	41
Tancapan	27	Tingang baguis	57
Tancong	62	Tingatinga	53
Tangain daga	27		66
Tangal	43	Tinisas	56
Tangantangan	64, 78		91
Tangi	92	Tintatinta	55
Tangili	22		76
Tangisang bayauac	79		83
Tanglad	92		80
Tanglay maloto	68		90
Tangolon	45	- ·	
Taointaoin	72	1 -	90
Tapiasin	62	· ·	
Tapit	78		59
	23		58
Tapolanga Taquing baca	23		63
			87
Taquipan	88	-	
Taquip asin	78		83
Taquip cohol	50		86
Taquip suso	50		43
Tarambolo	64		
Taramhampan	64	,	87
Taratara		1 -	21
Tarocanga	23	Torrecilla	66

	Page.		Page.
Toston	50	Ungalina mapula	36
Tres puntas	78	Varas hari	87
Trigo	91	Varas ni José	87
Tuba 17,	76, 77	Vilos	78
Tubang dalag	67	Vocabul	23
Tubli	87	Yabag	40
Tub6	92	Yacal	22
Tue	66	Yambo	45
Tugabang	41	Yampoi	45
Tuglima	51	Yantoc	88
Tugnang	61	Yate	67
Tulibas	28, 48	Yates	16
Tumahila	84	Yayo	27
Tumbong aso	54	Yerba buena	70
Tunas	17	Yerva mora	92
Tuttullacac	64	Yovas	67
Ualualisan	23	Zabache	39
Ubi	86	Zapote negro	58
Ubi ubihan	87	Zarzaparrilla	87
Timohona	41		

INDEX TO LATIN NAMES.1

Name.	This		co's Flo Filipinas		Novis- sima
Maine,	work.	First edition.	Second edition.		Appen- dix.
Abroma slata Blanco	25		422	2:404	29
communis Blanco	25 25		422	2:404	
A have a recetoring Tipp		606			29
Abrus precatorius Linn	38	565	394	2:361	62
Acalypha angatensis Blanco	77	750	516	8: 150	193
caroliniana Blanco	77	748	515	3:149	194
glandulosa Blanco	77	740	516	3:149	193
tomentosa Blanco	77	750		3:151	193
Acanthaceæ	66				152
Acanthus doloariu Blanco	66	487			154
ilicifolius Linn	66		341	2:267	154
Achras lucuma Blanco	57	237	166	1:299	124
sapota Linn	57	236	165	1:298	124
Achrostichum lagolo Blanco	95	826	570	3:240	343
simplex Blanco	95	826	571		
tripinnatum Blanco	95	826	571	3:241	343
Achyranthes aspera Blanco	71	188	133	1:239	170
villosa Blanco	71	189	134	1:240	170
Adelia acidoton Blanco	77	815	562	3:226	193
barbata Blanco	78	010	561	3:223	195
bernardia Blanco		814	301	0.220	195
glandulosa Blanco	77	814			194
monoico Planco	78	014		0.004	
monoica Blanco			561	3:224	194
papillaris Blanco	78		562	3:225	195
resinosa Blanco	77		562	3:225	193
Adenanthera gogo Blanco	42	353			78
Adiantum falcatum Blanco	94	833	576	3:251	330
lancea Blanco	94	833	576	3:250	330
lyratum Blanco	94	832	575	3:250	330
philippinense Blanco	94	831	575	3:249	330
tenuifolium Blanco	94	832	575	3:250	330
Aegiceras corniculatum	57	79	59	1:112	123
Aegilops fluviatilis Blanco	91	47	32	1:59	314
Aeginetia indica Roxb	65	l	342	2:270	149
Aeschynomene arborea Blanco	38	581	406	2:381	61
Agaricus	95	845	583	3:265	-
Agathis loranthifolia Blanco	82	,	528	3:170	211
Agave americana Blanco	85	258	180	1:322	253
Ageratium calomala Blanco	26		306	2:204	31
quadriflorum Blanco	55	624	300	2.204	114
Ailantus pongelion Blanco	29	380	268	0.194	349
Ajovea malabonga Blanco	74	233	200	2:134	
Alamaium astonatalum Dlamas	14	200		0.010	181
Alangium octopetalum BlancoAleurites lanceolata Blanco			310	2:210	108
Aleurites lanceolata Bianco	76	757	521	3:158	191
lobata Blanco	76	756			191
saponaria Blanco	76		517	8:156	191
triloba Forst	76		520	3: 157	191
trisperma Blanco	76	755			191
Allamanda cathartica Linn	58		64	1:120	128
Allantodia pinnata Blanco	94		571	3:242	334
Allium tricoccum Blanco	87	239	167	1:301	267
Aloe humilis Blanco	87	256	179	1:321	266
Alstonia batino Blanco	59	1	589	1:168	130

¹This index includes only the names used by Blanco, and family names according to the Bentham and Hooker system of classification. The numbers in the first column refer to the present paper; in the second, to the first edition of Blanco's Flora de Fllipinas; in the third, to the second edition; in the fourth, to the third edition; and in the fifth, to the Novissima Appendix.

111

Name. Ithaea rosea Cav sinensis Blanco marantaceæ marantus cruentus L	This work.	First	ı		
sinensis Blanco marantaceæ		edition.	Second edition.	Third edition.	sima Appen dix.
sinensis Blanco marantaceæ	23		386	2:345	2
	23	552			2
maranthiis critontiis I.	71		400	0.114	10
maraninus tructions II	71	711	492 492	3:114 3:114	10
mangostanus Blanco	71 71	711 710	492	3:114	1
spinosus Linn maryllidaceæ	85	/10	491	5.115	1 2
maryllis atamasco Blanco	85	254	178	1:319	2
merimnum mimosella Blanco	40	563	393	2:358	4
mmania debilis Ait	47	000	163	1:86	
monoflora Blanco	47	64			
ramosior Blanco	47		46	1:86	1
mmi glaucifolium Blanco	50	213			1
momum echinatum Blanco	84		3	1:4	1 2
zerumbet Linn	83	2	2	1:3	2
zingiber Linn	83	2	2	1:3	2
morpha glandulosa Blanco pedalis Blanco	37	555		0.040	
pedalis Blanco	19	553	387	2:348	1
mpelidaceæ	33				
nacardiacee	35 59	110	81	1:149	١.,
nasser laniti Blanco		112	. 61	1.149	1
navinga fuliginosa Blanco ndromeda japonica Blanco	48 28	372 371	261	2: 120	
ndropogon aciculatus Retz	92	9/1	26	1:49	۱ .
contortus Linn	91	38	26	1:49	8
nardus Blanco	92	39	27	1:51	8
ramosus Blanco	91	37	25	1:47	8
schoenanthus Blanco	92	38	27	1:50	8
nethum foeniculum Linn	50	214	150	1:270	ì
nonaceæ	15				1 1
nona muricata Linn	17		326	2:242	1
reticulata Linn	16	470	327	2:241	1
squamosa Linn	16	269	327	2:241	1
squamosa Linn nthemis cotula Blanco	55	638	l		. 1
nthistiria gigantea Cav ntholyza meriana Blanco	92	49	33	1:62	9
ntholyza meriana Blanco	85	24	18	1:33	2
ntidesma alexiteria Blanco	76		547	3:202	1
spicata Blanco	76	794		2:287	. 1
ntirrhinum comintanum Blanco	66	502	352	2:287	1
molle Blancopocynaceæ	66 58	503	353	2:288]]
pocynum mucronatum Blanco	60	852	143	1:259	1
poretica gemella Blanco	34	002	203	2:15	1
nenicillata Blanco	34	291	203	2:15	
penicillata Blanco ternata Blanco quilaria pentrandra Blanco	34	290	200	2.10	
quilaria pentrandra Blanco	19		373		
quilicia sambucina Blanco	33	178	0.0		
raceæ	89				
raceærachis hypogaea Linn	38	567	396	2:363	
raliaceæralia octophylla Blanco	51				. 1
ralia octophylla Blanco	51		158	1:284	
pendula Blanco	51	223	157	1:283	1 1
pipinnata Blancotripinnata Blanco	51	222	157	1:282	
tripinnata Blanco	51	223	401	0.100	.]]
reca catechu Linn	88	714	494	3:120	
catechu humilis Blanco	88	716	495	3:120	1 2
rethusa glutinosa Blanco	84 17	641 454	446	3:38 2:220	2
rgemone mexicana Linnrgophyllum pinnatum Blanco	31	186	316 131	1:235	
ristolochiaceæ	72	100	101	1.200	
ristolochia indica Blanco	72	282	197	1 349	1 5
sericea Blanco	72	282	198	1:350	1 1
subsagittata Blanco	$7\overline{2}$	283	197	1:350	1 1
rtemesia viridis Blanco	55		436	3:14	1 5
vulgaris Linn	56	625	435	3:14	1 1
rtocarpus camansi Blanco	80	740	467	3:76	1 3
in size T f	80	668	465	3:75	1 :
integrifolia L. f	80		466	3:76	2
integrifolia L. f	80	667	465	3:74	2
maxima bianco	80	669			
	80 80	671	467	3:78	2
odoratissima Blancoovata Blanco		666	464	3: 74 3: 77	

Nove		Blan	Nov:		
Name.	This work.	First edition.	Second edition.	Third edition.	Appe
Arum decurrens Blanco	90	656	457	3:62	
divaricatum L	90	657	458	3:62	
grandifolium Blanco	90		458	3:63	
Arundo tecta Blanco	93	48	33	1:60	:
Asclepiadaceæ	59				
Asclepias carnosa Blancodaemia Blanco	61 60	208 208	147	1:263	
gigantea Willd	60	208	146 146	1:263 1:262	
peregrina Blanco	60	207	146	1:262	
syriaca Blanco	60	204	144	1:261	
Atragene zeylanica Blanco	15	461	321	2:231	
Averrhoa bilimbi Linn	27	391	273	2:144	
carambola Linn	26	391	274	2:145	
pentandra Blanco	27	392	294	2:145	
Avicennia nitida Blanco	69	504			
tomentosa R. Br	69		353	2:289	
Azaola betis Blanco	57		281	2:157	
Azima nova Blanco	58	68	49	1:91	
Baccharis indica Linn ivæfolia Blanco	55 55	627	438	3:17	
Balingayum decumbens Blanco	48	627 187	437 132	3:16 1:237	
Bambus arundo Blanco	93	107	188	1:335	
diffusa Blanco	93	269	187	1:334	
levis Blanco	93	272	189	1:337	
lima Blanco	93	271	189	1:336	1
lumampao Blanco	93	272	189	1:338	
mitis Blanco	93	271	188	1:336	
monogyna Blanco	93	. 268	187	1:333	
pungens Blanco	93	270			
textoria Blanco	93	270	188	1:335	
Banara brevifolia Blanco	18	426			
racemosa Blanco	18	425			
Barreliera cristata Blanco prionitis Blanco	66	492	344	2:273	
Barringtonia racemosa Blume	66 46	491	343 373	2:272 2:325	
speciosa Forst	46	533	373	2:325	1
stravidium Blanco	46	533	313	2.320	1
Basella lucida Linn	71	216	151	1:273	
rubra Linn	71	215	151	1:272	1
Bassovia sylvatica Blanco	56		95	1:174	
Batis hermaphrodita Blanco	89	791	544	3:197	
Bauhinia binata Blanco	42	331	231	2:66	
castrata Blanco	42	331			
grandiflora Blanco	42	332	231	2:67	١.
purpurea Blanco	42		231	2:66	
scandens Blancotomentosa Blanco	42 42	332 330	232 230	2:68	
Begoniaceæ	50	330	250	2:65	
Begonia capensis Blanco	50 50	724	501	3:127	
Bergera compressa Blanco	28	360	254	2:107	
inodora Blanco	28	361		2.10.	
koenigii Blanco	28		254	2:108	}
ternata Blanco	28	360	254	2:108	
Bergia serrata Blanco	19	387			
Bidens bipinnata Linn	56	623	425	3:12	
Bignoniaceæ	66				
Bignonia quadripinnata Blanco	66	499	349	2:283	
spathacea Linn	66	499			
Biophytum sensitivum LBixaceæ	26 18		272	2:142	
Bixa orellana Linn	18	456	317	3:221	
Bladhia japonica Blanco	57	126	90	1:164	
Blechnum colubrinum Blanco	94	834	576	3: 252	1
Boerhaavia diffusa Linn	70	9	6	1:11	
Bombax ceiba Linn	24	L	372	2:372	
pentandrum Linn	$\frac{24}{24}$	531	371	2:324	1
Boraginaceæ	61				
Borago africana Blanco	62		60	1:114	
indica Linn	62	81	60	1:114	
Boswellia integra Blanco	30		242	2:86	
obliqua Blanco	30	1	243	2:87	1

Name		Blan	Novis- sima		
		First edition.	Second edition.	Third edition.	Appen- dix.
Brabejum caliculatum Blanco	75		39	1:72	355
concatenatum Blanco	58		40	1:73	129
lucidum Blanco	58		40	1:74	129
pinnatum Blanco	58		40	1:74	129
Brassica orientalis Blanco	17	• 519	361	2:303	10
Bromeliaceæ	86				256
Bromelia ananas Linn	86	230	162	1:291	256
Broussonetia luzonensis Blanco	78		488	3:107	198
tinctoria Blanco	80		488	3:108	202
Bruguiera nemorosa Blanco	44 43		275	2: 147 2: 47	79 78
Bryophyllum germinans Blanco			220		
serratum Blanco	43		220	2:48 2:48	78 78
triangulare Blanco	43	856	221	2:48	
Bucida comintana Blanco	44		265		80
Buddleia virgata Blanco	61	57	38	1:70 2:36	136
Buginvillea racemosa Blanco	70 29	307	214	2.30	167
Burseraceæ Bursaria inermis Blanco	29 18		124	1:222	40
Cacalia sarracenica Blanco	18 56	010	124	1.222	13
		618			119
sonchifolia	56 50	618			120 99
		414		9.171	
Cactus opuntia Blanco	50 50	414	288 289	$2:171 \\ 2:172$	99
pitajaya Blanco		000		0.70	
Caesalpinia ignota Blancosappan Linn	41	336	235	2:72 2:72	70
tarquata Planas	41 41	335 336	234	2.14	69 70
torquata Blanco	40	550	410	2:396	
Cajanus bicolor DC	38		416	2. 390	66
quinquepetalus Blanco	40		417	2:397 2:398	61 66
volubilis Blanco			417		
Caladium digitatum Blanco	90 90		459	3: 64 3: 64	$\frac{294}{292}$
esculentum Blanco	90 88	267	459 186	1:332	292
Calamus gracilis Blanco	88	266	185	1:331	276
maximus Blanco	88 88			1:329	
mollis Blanco	88 88	264 265	184 185	1:330	$275 \\ 275$
usitatus BlancoCalius lactescens Blanco	78	698	543	3:103	198
Calla badian Blanco	90	658-659	459	3:64	298
gaby Blanco	90	659	409	3.04	292
maxima Blanco	90	658			294
polyphylla Blanco	90	659			294
Callicarpa americana Blanco	67	517	360	2:300	158
Calophyllum apetallum Blanco	20	517	429	2:415	17
inonhyllum Linn	20	612	428	2:415	17
inophyllum Linn Calyptranthes jambolana Willd	46	418	420	2.110	86
makul Blanco	46	419			85
ramiflora Blanco	46	420			86
zuzugium Blanco	46	120	293	2:179	85
Calytriplex oboyata R. et P	65		361	2:203	147
Calytriplex obovata R. et PCambogia binucao Blanco	20	434	302	2:196	16
crassifolia Blanco	20		304	2:198	16
venulosa Blanco	20	435	303	2:197	1 16
Camellia sasangua Blanco	21	530	371	2:322	19
Campanulaceæ	56	000	0.1	2.022	1
Canarium album Blanco	30	793	546	3:201	40
commune Blanco	30	791	310	3.232	40
pimela Blanco	30		545	3:201	40
Canavalia ensiformis Blanco	39		404	2:578	64
gladiata DC	39		403	2:376	64
Canna indica Linn	84	6	5	1:8	229
Cansiera grossularioides Blanco	76	73	53	1:99	190
pentandra Blanco	76	73	53	1:98	190
rheedii Blanco	76	73	52	1:98	190
Canthium horridum Blume	53		J16	1:208	110
monoflorum Blanco	53	166			110
pauciflorum Blanco	53	165			110
pedunculare Cav	53	1	116	1:208	110
	17				10
Capparidaceæ	18	438			l îi
CapparidaceeCapparis baducca Blanco			1	1	
Capparis baducca Blanco			305	2:200	1 1 1
Capparis baducca Blanco	18	438	305 305	2:200 2:200	
Capparis baducca Blanco				2:200 2:200 2:199	11 11 11

Name.	This	Blan	co's Flor Filipinas	ra de	Novis- sima
Name.	work.	First edition.	Second edition.	Third edition.	Appen- dix.
Capparis odorata Blanco	18 51	439	305	2:201	11
CaprifoliaceæCapsicum minimum Roxb	64	133	95	1:174	103 145
Capura pinnata Blanco	35	264			52
purpurata BlancoCardamine glandulosa Blanco	$\frac{35}{17}$	521	184	1:328	52 9
impatiens Blanco	17		363	2:306	9
Cardiospermum halicacabum Linn	. 34	312	218	2:44	50
Carex glomerata Blancotuberosa Blanco	91 91	35	24 24	1:45 1:45	308 306
Carica hermaphrodita Blanco	49	803	554	3:212	95
papayā LCarmonea heterophylla Cav	49 62	803 209	553 147	3:212 1:265	95 138
Carthamus tinctorius Linn	56	616	431	3:6	120
Caryophyllaceæ	19	1			14
Caryota onusta Blanco	88 88	741	511 513	3: 143 3: 145	280 280
palindan Blancotremula Blanco	88	744	512	3: 145	280 280
urens Blanco	88	740	510	3:142	279
Casearia fuliginosa Blanco	48	990	$\frac{262}{237}$	2:123	98
Cassia alata Linn fistula Linn	41 41	339 339	$\frac{237}{237}$	2: 77 2: 76	70
longisiliqua Blanco	41	338	201	2.70	70
mimosoides Linn	41	340	237	2:78	7:
occidentalis Linnsulcata Blanco	41 41	338	$\frac{236}{236}$	2:75 2:76	70
tora Linn	41	337	235	2:74	70
Casuarina equisetifolia Forst	81	661	460	3:67	20'
Cassuvium reniforme Blanco	36	322	227	2:20	5-
Cassytha filiformis LCasuarinaceæ	74 81	321	226	2:58	182 200
Cedrela odorata Blauco	31	184	130	2:130	4
taratara Blanco	31		131	1:234	45
Cedrota guinanensis BlancoCelastraceæ	70 32		213	2:33	16'
Celosia argentea Linu	71	192	135	1:243	16
baccata Retz	71	193			167
bicolor Blancococcinea Blanco	71 71	191	134	1:241	168 16'
cristata Linn	71	191	194	1.241	16
glauca Blanco	71		135	1:242	16
lanata Blanco	71 47	192	134	1:241	169
nana BlancoCeltis lima Blanco	78	192	139	1:250	19
philippinensis Blanco	78	197	139	1:249	19
Cenchrus hexaflorus BlancoCephaelis expaleacea Blanco	91 52	36	24	1:46 1:189	31 10
Cerbera manghas Blume	58 58	125	103	1:189	10
thevetia Linn	58	125	89	1:162	12
Chailletiace:	31				4.
ChenopodiaceæChenopodium ambrosioides Linn	71 71	200	140	1:253	17 17
Chironia capsularis Blanco	63	102	140	1.200	14
lanosanthera Blanco	63		71	1:132	14
ChloranthaceæChloranthus inconspicuus Blanco	72 72		54	1:101	17'
Cieca acidissima Blanco	76	700	486	3:105	18
decandra Blanco	76	701	487	3:106	18
pentandra Blanco	76 17	701	486 563	3:105 3:227	18
Cissampelos pareira LinnCissus acida Roxb	33	815 69	50	1:94	4
alata Blanco	33		. 51	1:95	4
arborea Blanco	27		. 51	1:95	3
frutescens Blancolatifolia Vahl	27 33	70	52	1:96	3
pedata Blanco		71	52	1:96	5
pedata Blanco quadraugularis Linn	33	72	52	1:97	4
rubescens Blancosimplex Blanco	33	71 72			4
vesicatoria Blanco	33	12	50	1:94	4
Citrus aurantium	. 28	609	425	2:408	3
decumana Linn	28	608	424	2:406) 3

notissima Blanco papillaris Blanco reticulata Blanco reticulata Blanco reticulata Blanco eome alliacea Blanco alliodora Blanco gigantea Blanco pentaphylla Blanco retodendron capsulare Blanco fortunatum Blanco ethra alnifolia Blanco ethra alnifolia Blanco ethra atripularis Blanco diffusa Blanco diffusa Blanco glabra Blanco glabra Blanco yolubilis Blanco rifolia Blanco polyphylla Blanco routifusa Blanco polyphylla Blanco polyphylla Blanco routifera Linn damba blanco dichotoma Blanco nucifera Linn ffea arabica Linn yolubilis Blanco yolubilis Blanco enopteris quadripinnata Blanco motifera Linn dichotoma Blanco pumilaris Blanco yolubilis Blanco sucifera Linn dichotoma Blanco pumilus Blanco yolubilis Blanco	This		co's Flo Filipinas		Novis- sima	
Name.	work.	First edition.	Second edition.	Third edition.	Appen- dix.	
Citrus mitis Blanco	29	610	426	2:409	37	
notissima Blanco	28	607	424	2:406	37	
papiliaris Blanco	28	610	425	2:409	37	
torosa Blanco	28 28	610 609	425 425	2:408 2:407	37	
Cleome alliacea Blanco	18	522	420	2.401	10	
alliodora Blanco	18		363	2:307	10	
gigantea Blanco	18		364	2:307	10	
	18	523 522			10	
Clerodendron cansulare Blanco	17 68	509	355	2:292	160	
fortunatum Blanco	68	508	345	2:291	16	
Clethra alnifolia Blanco	56		259	2:117	121	
Clitorea ternatea L	40	590	412	2:391	6	
Chostis comiculate Planco	75 37	818 386	564 270	3:229 2:138	168	
diffusa Rianco	36	386	210	2.130	56	
erecta Blanco	37	287			5'	
glabra Blanco	37	387	271	2:138	5	
polyphylla Bianco	36		270	2:137	5	
trifolia Blanco	36		270	2:136	50	
VOIUDIIIS BIANCOCOCOS mammilaris Planco	36 88	385 722	499	3:123	28	
	88	716	495	3:123	28	
Codamba blancoi Azaola	61	710	591	2: 293	13	
dichotoma Blanco	61	510	355	2:293	13	
Coenopteris quadripinnata Blanco	94	833	576	3:252	33	
Coffea arabica Linn	54	156	110	1:198	11	
VOIUDIIIS Blanco	54 91	157 688	111 478	1:200 3:92	11 31	
Coldenia procumbens Linn	62	74	56	1.105	13	
Coleus grandifolius Blanco	69	482	336	2:258	16	
pumilus Blanco	69	482	336	2:257	163	
suganda Blanco	69	483	337	2:259	16	
Columbia anilao Blanco	26 26	654	426 427	2:412 2:412	30	
	44		721	2.412	80	
Combretum distillatorium Blanco	44	295			8	
laxum Blanco	44		206	2:206	8:	
Commelinaceæ	87		10	1.04	28	
Commercania cobinete Blanco	87 25	25	18 160	1:34 1:287	26 2	
Comocladia serrata Blanco	32	30	100	1.201	4	
Composite	54				11	
Coniferæ	82				21	
Conferva littoralis Blanco	95	843	582	3:262		
setosa BlancoConnaraceæ	95 36	844	583	3: 264	5	
Connarus foetens Blanco	28	525			3	
santaloides Blanco	28		366	2:314	3	
Convolvulaceæ					13	
Convolvulus batatas Linn	62	93	68	1:129	14	
boerhaavioides Blanco	63 63	94	67	1:125	14 14	
catharticus Blancocolubrinus Blanco	63	94	66	1:125	14	
dentatus Blanco	63	89	66	1:123	14	
distillatorius Blanco	63	95	70	1:130	14	
hederaceus Blanco	63	90	66	1:124	14	
longiflorus Spreng	63		69	1:130	14	
maximus Blanco muricatus Blanco	63 62	91 92	67 68	1:127 $1:127$	14 13	
nil Linn	62	92	68	1:127	14	
paniculatus Linn	63	96	70	1:131	14	
pes-caprae Linn	. 62	88	65	1:123	14	
pes-tigridis Linn	63	97	71	1:133	14	
reniformis Roxb	63	91	67	1:126	14	
repens Willd	62	92 90	68	1:128	14	
valerianoides BlancoConyza balsamifera Linn	63 55	628	438	3:18	14	
COMPAN TON	55	020	438	3:18	11	
Cadda Haii						
cappa Hamdentata Blancoerosa Blanco	55 55	629	439	3:19	. 11 11	

Name.	This	Blan 1	Novis sima		
name.	work.	First edition.	Second edition.	Third edition.	Apper dix.
Conyza gouana Blanco	55	629			1
Cookia anisodora Blanco	28	i	253	2:314	1
anisum-olens Blanco	28	359			
wampi Blanco	28	358	253	2:105	}
Corchorus acutangulus Linnaestuans Blanco	26		308	2:206	
aestuans Blanco	26	441		2:206	
capsularis Linncatharticus Blanco	$\frac{26}{26}$	442 442	308	2:206	
olitorius Linn	26 26	442	308	2:207	
Cordia banalo Blanco	61	124	300	2.201	1
dichotoma Blanco	62	123	88	1:159	ĺ
ignota Blanco	61		88	1:160	l ī
olitoria Blanco	70	123	88	1:159	1
sebestina Blanco	62	121	87	1:158	1
Coreopsis gracilis Blanco	56		591	3:25	1
Jornaceæ	51				1
Coronilla emerus Blanco	38	582		1.000	
Corypha minor Blanco	88 88	229 228	161	1:290 1:290	2
umbraculifera LiunCostus luteus Blanco	88 84	228	160	1:290	2
nigricans Blanco	84 84	3	3 3	1:5	
Cotula quinqueloba Blanco	56	626	436	3:15	ĺ
Cotyledon lanceolatum Blanco	43	382	100	0.10	
paniculata Blanco	43	381			
paniculata Blancoserrata Blanco	43	382			
Crassulaceæ	43				
Crataeva octandra Blanco	18	400	280	2:280	
religiosa Forst	18	399	279	2:154	
Crescentia trifolia Blanco	66	489	343	2:271	1
Crinum asiaticum Blanco	85	251	175	1:314	1 3
giganteum Blanco	85 37	570	175	1: 315	2
Crotalaria linifolia Linn. fpallida Blanco	37	570			1
pumila Blanco	37	370	397	2:365	1
quinquifolia Linn	37	572	398	2:267	1
Croton drupsceum Blanco	77	0.2	519	3:155	1
Croton drupaceûm Blancograndifolium Blanco	78	753	518	3:153	1 1
glandulosum Blanco	77	754			. 1
lacciferum Blanco	78	752	517	3:153	1 1
muricatum Blanco	77		518	3:154	1
variegatus L	77	751	517	3:152]]
Cruciferæ	17				-
Crudia spicata Blanco	41	776	261	2: 121 3: 179	
Cucumis acutangulus Linn	49 50	861	534 534	3:179	
luzonicus Blancomelo Linn	50	775	534	3:179	
Cucurbitaceæ	49	110	994	3.179	
Cucurbita lagenaria-oblonga Blanco	49	772	531	3:175	-
lagenaria-villosa Blanco	49	772	532	3:175	1
pepo-aspera Blanco	49	773	532	3:176	
sulcata Blanco	50	773	532	3:177	
Cupania spinosa BlancoCupulifereCupulifere	32	184	204	2:17	
Cupuliferæ	81				. :
Curcuma longa Linn	83	5	4	1:6	
Cycadaceæ Cycas circinalis Linn Cylista piscatoria Blanco	82			0.140	-
Cycas circinalis Linn	82 37	745 589	513	3:146	:
Cynanchum hirtum BlancoCynanchum hirtum Blanco	60	203	143	1:258	1
tenellum Blanco	60	203	143	1:258	
viminale Blanco	60	203	143	1:257	1
Cynomorium philippinense Blanco	96	665	464	3:72	
Cyperaceæ	90				-1
Cyperus difformis	80	32	22	1:41	
paniculatus Blanco	93	32	22	1:42	i
rotundus Blanco	91	. 31	26	1:40	
Cypripedium bulbosum Blanco	84	637	444	3:33	
Cyrtandra aristata Blanco	65		. 13	1:24	
glaberrima Blanco personata Blanco	65		- 12	1:23	
personata Blanco	65		13 269	1:25 2:135	
quinquestila Blanco	72		209	4;150	
Cytinaceæ	40		-		-1

Name. Sytisus quinquepetalus Blanco volubilis Blanco Dais laurifolia Blanco Dalea alopecuroides Blanco Dapne aquilaria Blanco indica Linu Datura fastuosa Linn metel Blanco Dalecus anisodorus Blanco Decringia celosioides R. Br Delima aspera Blanco Dendrobium teres Blanco teretifolium Blanco Dentella repens Forst Desmodium diversifolium Blanco gangeticum Blanco parvifolium Blanco parvifolium Blanco parvifolium Blanco parvifolium Blanco	work.	-			Novis-	
volubilis Blanco Dais laurifolia Blanco Dalea alopecuroides Blanco Dapne aquilaria Blanco foetida Blanco indica Linu Datura fastuosa Linn metel Blanco Deceringia celosioides R. Br Delima aspera Blanco Dendrobium teres Blanco Dentella repens Forst Desmodium diversifolium Blanco Dentella repens Forst Desmodium Blanco Dendrobium diversifolium Blanco	work.	First edition.	Second edition	Third edition.	Appen dix.	
Dais laurifolia Blanco Dalea alopecuroides Blanco Dapne aquilaria Blanco foetida Blanco indica Linu Datura fastuosa Linn metel Blanco Daucus anisodorus Blanco Deeringia celosioides R. Br Delima aspera Blanco Dendrobium teres Blanco Dendrobium teres Blanco Dentella repens Forst Desmodium diversifolium Blanco Dentella repens Forst Desmodium Blanco Dentella regens Forst Desmodium Blanco	38	598			6	
palea alopecuroides Blanco Dapne aquilaria Blanco foetida Blanco indica Linn Datura fastuosa Linn metel Blanco leucus anisodorus Blanco leeringia celosioides R. Br Delima aspera Blanco Dendrobium teres Blanco Dendrobium teres Blanco Dentella repens Forst Desmodium diversifolium Blanco gangeticum Blanco gangeticum Blanco gangeticum Blanco	$\frac{40}{74}$	599 375	263	2:125	18	
Dapne aquilaria Blanco foetida Blanco indica Linn Datura fastuosa Linn metel Blanco Daucus anisodorus Blanco Deeringia celosioides R. Br Delima aspera Blanco Dendrobium teres Blanco Dentella repens Forst Desmodium diversifolium Blanco Dentella regangeticum Blanco Dentella regens Forst Desmodium diversifolium Blanco Dentella regens Forst Desmodium diversifolium Blanco Dentella regens Forst	37	310	389	2; 351	5	
indica Linu Datura fastuosa Linn metel Blanco Daucus anisodorus Blanco Deeringia celosioides R. Br Delima aspera Blanco Dendrobium teres Blanco teretifolium Blanco Dentella repens Forst Desmodium diversifolium Blanco gangeticum Blanco gangeticum Blanco	74	310	216	2:39	18	
indica Linu Datura fastuosa Linn metel Blanco Daucus anisodorus Blanco Deeringia celosioides R. Br Delima aspera Blanco Dendrobium teres Blanco teretifolium Blanco Dentella repens Forst Desmodium diversifolium Blanco gangeticum Blanco gangeticum Blanco	74	308	214	2:37	18	
metel Blanco	74	309	215	2:38	18	
Daucus anisodorus Blanco Deeringia celosioides R. Br Delima aspera Blanco Dendrobium teres Blanco teretifolium Blanco Dentella repens Forst Desmodium diversifolium Blanco gangeticum Blanco gangeticum Blanco	64	100	73	1:136	14	
peeringia celosioides R. Br pelima aspera Blanco pendrobium teres Blanco teretifolium Blanco pentella repens Forst pesmodium diversifolium Blanco gangeticum Blanco gangeticum Blanco	64 50	98	72 150	1:136 1:269	14	
Delima aspera Blanco Lendrobium teres Blanco teretifolium Blanco Lentella repens Forst Lesmodium diversifolium Blanco gangeticum Blanco	71		135	1:244	10	
Pendrobium teres Blanco teretifolium Blanco Pentella repens Forst Pesmodium diversifolium Blanco gangeticum Blanco gangeticum Blanco	15	429	299	2:191		
Dentella repens Forst	84	638			24	
Desmodium diversifolium Blanco	84	7.40	444	3:34	24	
gangeticum Blanco	52 38	146	103 408	1:190 2:384	10	
parvifolium Blanco	38		408	2:384		
parvironum Dianco	38		408	2:386	(
spirale DC	38		408	2:385		
Dianthera americana Blanco	67	16	12	1:22	1	
ciliata Blanco	67	16	12	1:22	1	
subserrata Blanco Dicerma pulchellum DC	67 38	10	11 407	1:22 2:383	1	
Diceros stoloniferus Blanco	64		349	2: 282	1	
oilleniaceæ	15				1	
oillenia indica Blanco	15	427				
speciosa Blanco	15		329	2:244		
Dioscorea eae	86 86	799	550	3:207	$\frac{2}{2}$	
bolojonica Blanco	39	800	551	3:208		
divaricata Blanco	86	797	550	3:206	2	
papillaris Blanco	86	801	552	3:210	2	
pentaphylla Linn	86	802	552	3:210	2	
sativa Blanco	87		551	3:209	20	
triphylla Blancotugui Blanco	86 87	799 800	551	3:208	20	
Diosma serrata Blanco	32	168	119	1:213	1 -	
Diospyros biflora Blanco	57	303	210	2:28	1:	
embryopteris Blanco	58		209	2:28	1	
kaki Blanco	57, 58	302	211	2:29	126, 1	
lotus Blanco multiflora Blanco	57 57	210 303		2:29	1	
nigra Blanco	58	303	211	2:30	1	
pilosanthera Blanco	57	304	211	2:31	l î	
Dipterocarpaceae	21			·		
Dipterocarpus grandiflorus Blanco	21		314	2:218		
guiso Blanco	$\frac{22}{22}$		313 312	2:215 2:214		
malaanonan Blanco mangachapoi Blanco	22		313	2:214		
mayanis Blanco	$\frac{21}{21}$		313	2:215		
palosapis Blanco plagatus Blanco polyspermus Blanco	22		312	2:214		
plagatus Blanco	-22		311	2:212		
polyspermus Blanco	$\frac{22}{22}$		312	2:213 2:212		
thurifer Blancovernicifluus Blanco	21		310 314	2:217		
Oodonaea angustifolia Linn	35	312	217	2:40		
Polichos acinaciformis Blanco	39	578			.]	
echinulatus Blanco	40		401	2:373		
ensiformis Linn	39	577	402	2:373		
repens Blancosesquipedalis Blanco	39 40	577	402	2:375		
sinensis Linn	40	575	102	2.010		
tetragonolobus Linn	40	576	402	2:402	1	
trilobus Blanco	39		403	2:375		
Oombeya biserrata Blanco	25		244	2:90		
decandra Blanco	25	349	481	3:98	2	
Porstenia pubescens Blanco Pracaena terminalis Linn	80 87	692 263	183	1:328	2	
Proseraceæ	43	200	100	1.020		
Prosera hexagynia Blanco	43	226	159	1:286		
Chites caudata Blanco	57 59	106	77	 	1	

		Blan	Nov sin		
Name.	This work.	First edition.	Second edition.	Third edition.	App
Cchites procumbens Blanco	59		78	1:145	
repens Blanco	59	109			
scholaris Linn	59	106	77	1:144	1
trifida Blanco	59	109	79	1:146	
spiralis Blanco	59	110	79	1:146	
Ehretia beurreria Blanco	62	127	91	1:166	
virgata Blanco	62	127	90	1:165	
Elaeagnaceæ	74				
Elaeagnus angustifolia Blanco	$\frac{74}{32}$	74	53 306	1:100	ĺ
Elaeocarpus integrifolius Blancosylvestris Blanco	26		306	2: 202 2: 203	
Elatineace	19		300	2.203	
Elcana seminuda Blanco	58		584	3: 267	
Elephantopus dubius Blanco	55		442	3: 28	1
scaber Linn	54	634	441	3:27	
serratus Blanco	55	635	442	3:28	
Elytraria amara Blanco	66	000	8	1:15	
Emelia sonchifolia DC	56	432	432	3:8	1
Enrila dichotoma Blanco	32	709	102	0.0	
Entada pursaetha DC	42	100	247	2:96	1
Eperua decandra Blanco	41	368	259	2:118	İ
falcata Blanco	41	369	1 200	2.110	1
rhomboidea Blanco	41	000	261	2:119	1
Epidendrum equitans Blanco	84	645	449	3:44	
lineare Blanco	84	644	449	3:44	
vanilla Blanco	85	643	110	0.11	1
Ericaceæ	56	0.10			1
Erotium lanigerum Blanco	96	440	307	2:205	
Erythrina caffra Blanco	39	110	394	2:360	1
carnea Blanco	39	564	393	2:359	
picta Blanco	39	565			
pieta BlancoEugenia bauanguica Blanco	46	418	290	2:174	1
cauliflora Blanco	46		291	2:177	
glandulosa Blanco	46	417	291	2:176	1
jambos Blanco	45	416	290	2:175	
lobas Blanco	46	857		.	-1
malaccensis Blanco	45	415	290	2:173	
mananquil Blanco	45		_ 290	2:174	1
montana Blanco	45	418	291	2:175	l
Eupatorium ayapana Vent	55	619	432	3:9	
Euphorbiaceæ	75		-} -	-	-
Euphorbia capitata Lam	75	411		-	-
dulcis Blanco	75	412	287	2:167	
hirta L	75		286	2:166	
parannaquensis Blanco	. 75		286	2:165	1
pentagona Blanco	75	413	287	2:168	1
tirucalli L	75	412	287	2:167	
Euphoria cubili Blanco	35	287	200	2:10	1
didyma Blanco	35	288	201	2:10	
litchi Blanco	35	285	199	2:8	1
nephelium Blanco	22		200	2:9	
annularis Blanco	34	285	199	2:7	1
malaanonan Blanco	22	289		1.00	-
Evodia bintoco Blanco	27 27		- 50 - 50	1:93 1:91	
triphylla DC Evolvulus linifolius Linn	63	221	- 50 156		1
Exacum albens Blanco	61	58			
Excaecaria agallocha Blanco	78	786	541	3:193	
laevis Blanco	78	788	543		
sicca Blanco		787	542		
Exostemma philippicum Blanco	52	1 101	113	1:203	
Exostemma philippicum Blanco Fagara decandra Blanco	36	66		1:89	
octandra Blanco	27	67	48		
piperita Blanco	27	64	47	1:87	
pterota Blanco	27	66		1:88	
Fagraea scholaris Blanco	61		_ 93	1:171	1
Fagus philippensis Blanco	81		503	3:132	
Feronia ternata Blanco	28		252		
Ficoideæ					
Ficus argentea Blanco	79	681	473	3:84	
aspera Bianco	79				
aspera nota Blanco	79			1	1

Namo		Blan	Novis-		
Name.	This work.	First edition.	Second edition.	Third edition.	Apper dix.
Ficus aspera volubilis Blanco	79	676	472	3:82	2
dicarpa Blancoglomerata Blanco	79	682		0.05	20
hauili Blanco	79 79	682 684	475	3:87	$\frac{2}{2}$
heterophylla Blanco	79	685	476	3:89	2
hispida Blanco	79	685	476	3:88	2
hispida hastata Blanco	79 79	685 685	476 476	3:89 3:88	$\frac{2}{2}$
hispida linearis Blancohispida odorata Blanco	79	686	476	3:89	2
indica Linn	78	681	473	3:85	1
laccifera Blanco	80	673	468	3:80	2
laevigata Blancolaurifolia Blanco	79 79	682	474 475	3: 86 3: 87	$\frac{2}{2}$
nepalensis Blanco	79		474	3:85	2
payapa Blancopseudopalma Blanco	79	683	475	3:86	1
pseudopalma Blanco	80	680	473	3:84	2
rostratâ Blancoscabra Blanco	79 79	679	472 471	3:83 3:81	$\frac{2}{2}$
Filices	94		111	0.01	3
Fissilia psittacorum Blanco	32	- 28	20	1:38	
Flacourtia corollata Blanco	18		559	3:220	
parvifolia BlancoFlagellaria indica Linn	18 87		560 196	3:220 1:347	2
Fucus denticulatus Blanco	95	839	130	1.047	4
edulis Blanco	95		580	3:259	
gulaman Blanco	95	839			
natans Blancoprolifer Blanco	95 95	838	579 579	3:259 3:259	
Fusanus parasitus Blanco	75	000	53	1:100	1
alactia terminiflora Blanco	37		411	2:390	
Faledupa frutescens Blanco	40	559	391	2:354	
maculata Blancopungam Blanco	40 38	559 558	390 390	2:353 2:352	
Gentianacee.	61	000		2.002	1
deraniaceæ	26				
Geruma subtrilobata Blanco	25 65	182	130	1:232	1
desneriaceæ dethyllis acaulis Blanco	85	260	181	1:324	2
Simbernatia calamansanay Blanco	44		266	2:129	1
linus leptoides Linn	50	413	288	2:169	1
Huta orgyalis Blanco	75	578	451	3:49] 1
Hycine lucida Blanco Helina asiatica Blanco	40 68	492	344	2:274	1
inermis Blanco	68	493	345	2:274	l i
Anaphalium dichotomum Blanco Anetaceæ	55		439	3:20	1
Inetacee Inetum gnemon Linn	82 82	747	514	3:147	2
Inidia oppositifolia Blanco	45	299	208	2:25	3
doodeniacee	56				
Somphrena globosa Linn	71	198	139	1:251	
volubilis Blanco Fordonia polysperma Blanco	71 48	199 549	140 384	1:252 2:342	-
Sossypium herbaceum Linn	24	534	374	2:329	
paniculatum Blanco	24	539	378	2:331	
perenne Blanco	24	537	376	2: 330	
religiosum Linn Fouania domingensis Blanco	24 33	196	376 138	2:329 1:248	
Framineæ	91	150	100	1,240	
ratiola hyssipioides Blanco	65	11	8	1:16]
Frewia mallococca Blanco	26		310	2:209	
multiflora Blanco	26 26	444	309	2:209	
serrata Blanco	45	187	132	1:236	
luaiacum abilo Blanco	30	364			
łuettarda jazminiflora Blanco	51	722			
polyandra Blanco	58 51		500 499	3:126 3:124	
speciosa Blancovermicularis Blanco	51 21	723	500	3:124	1
Guilandina bonducella Linn	41	343	239	2:81	1
nuga Linn	41	344	240	2:81	
Auttiferæ	20			.	. I

Name		Blanco's Flora de Filipinas.				
Name.	work.	First edition.	Second edition.		sima Appe dix	
Tyrocarpus pendulos Blanco	81		55	1:104	2	
Haemanthus pubescens Blanco	85	253			2	
Halesia ternata Blanco	45	399	279	2:153	1	
Hedysarum pulchellum Linn	38	581				
vespertilionis LinnHelianthemum triflorum Blanco	38	581	407	2:382	1	
Ielianthemum trifforum Blanco	$\frac{21}{24}$	383	309	2:208		
lelicteres apetala Blanco pinnata Blanco	36	384				
leliotropium parviflorum Blanco	62	80	59	1:113	١.	
Ielminthostachys dulcis Kaulf	95		596	3:254]	
Iemionitis incisa Blanco	94	829	574	3:246		
Jernandia sonora Blanco	74	689	478	3:93	ì	
Ieritiera tinctoria Blanco	24	653	456	3:59		
Hibiscus abelmoschus Linn	23	545	380	2:335		
batacensis Blanco	$\frac{23}{23}$	544	380	2:334		
bifurcatus Blanco	23 23	545 546	380	2:334 2:336		
mutabilis Linnpopulneus Linn	23	544	381	2;350		
rosa-sinensis Linn	23	543	379	2:333		
tiliaceus Linn	23	541	379	2:332		
Iillia longiflora Blanco	75	235	165	1:297)]	
Hippocratea volubilis Blanco	32	27	20	1:27		
Jippoerenis comosa Blanco	38	584				
humilis Blanco	38	585				
humilis Blanco multisiliquosa Blanco rhomboidea Blanco	38	584				
rhomboidea Blanco	38	585			-	
Hiraea reclinata Blanco	26 92	378 47	32	1:58	١.	
Holeus saccharatus Linn	60	47		1: 257	1	
Hoya carnosa BlancoHoya carnosa BlancoHydnocarpus polyandra Blanco	18		142 545	3:200		
Hydrocharideæ	82		940	3.200		
Avdrocotyle asiatica Linn	50	212	149	1:268	1.	
monopetala Blanco	65	213				
Ivdrolea aravatensis Blanco	61	211				
zeylanica Vahl Hydrophyllaceæ	61		148	1:266	1 :	
Hydrophyllaceæ	61					
1vpericaceæ	19			0. 410		
Hypericum aegyptium Blanco	20 19	615 613	430 429	2:418 2:416		
olympicum Blanco	25	614	429	2:417		
eica abilo Blanco	30	014	256	2:113		
cica abilo Blanco gnatia amara Linn. f	61	82		2.110		
llecebrum lanatum L	71	190				
Impatiens triflora Blanco	27	636	443	3:32		
ndigofera angustifolia Blanco	37	596	415	2:394		
argentea Blanco	37		415	2:394		
hirsuta Blanco	37	591		0.000	-	
senegalensis Blanco	37 37	591	415 413	2:392 2:393		
tinetoria Blanco Inga lanceolata Blanco	43	991	370	2:322		
Ipomoea hepaticifolia Blanco	63		72	1:134		
quamoclit Linn	62	97	72	1:134		
ridaceæ	85					
[xora arborea Blanco	54	61	42	1:78		
coccinea Linn	54	59	41	1:76		
glandulosa Blanco	53	61	42	1:78		
manila Blanco	53	60	42	1:77		
Jatropha eureas L janipha Blanco, non L	76	759	522	3:160		
manihot Linn	76 77	758 760	521 522	3:159 3:160		
Juglandace	81	700	922	3.100	1	
Jussiaea erecta Blanco	48	365	257	2:114		
inclinata Blanco	48	366			.i	
fluviatilis Blanco	48		257	2:114	1	
Justicia dalaora Blanco	67	14	10	1:10		
ecbolium Blanco	67	13	10	1:18	1	
gendarussa Blanco	67	14	10	1:19	1	
nasuta Linn	67	14	10	1:19		
picta Linn	67	12	9	1:18		
viridis Blanco Kadsura blancoi Azaola	67 15	15	11 594	1:21 3:118		
			. 094		1	

Name.	This	Blan	Novis- sima		
Name.	work.	First edition.	Second edition.	Third edition.	Appen- dix.
Kirganelia alba Blanco	75	713	494	3:117	18
nigrescens Blanco	76	712	493	3:115	18
pumila Blanco	76	712	493	3:117	18
triandra Blanco	75	711	492	3:115	18
villosa Blanco	75	712	493	3:115	18
Kleinhovia hospita Linn	24	652	455		18
correte Plance				3:58	2
serrata Blanco	96	653	456	3:58	35
Knautia sagittata Blanco	55	54	36	1:67	11
Koelreuteria arborea Blanco	34		202	2:13	5
edulis Kyllingia triceps Blanco	35	34	202	2:13	5
	90		23	1:44	30
Kyrtandra aristata Blanco	65	18			15
capsularis Blanco	65	17			14
personata Blanco	65	18	[14
serrata Blanco	65	18	- -		14
Labiatæ	69				16
ablab cultratus DC	40		405	2:379	6
Lagerstroemia indica L	47	454	316	2:219	9
Lansium domesticum Rumph	31		228	2:62	4
antana viburnoides Blanco	67		345	2:275	15
auraceæ	73				17
Laurus cassia Blanco	73	317	223	2:53	17
cinnamomum Blanco	73	319	225	2:57	17
culilaban Blanco	73	316	222	2:51	17
hexandra Blanco	74		222	2:52	18
lanosa Blanco	$7\overline{4}$	318	224	2:54	18
martinicensis Blanco	73	317	223	2:53	17
persea L	73	01.	224	2:56	18
serrata Blanco	48	319	224	2:55	100
Lawsonia inermis Roxb		294	206	2:21	
Lechea minor Blanco	47 87	52	35	1:65	9
Leea aculeata Blanco		32		1:227	26
sambucina Blanco	34		127	1: 227	5
Sambucina Bianco	33		126	1:226	5
Legazpia triptera Blanco Legnotis lanceolata Blanco	65		339	2:264	14
Legnous lanceolata Blanco	96	445			
Leguminosæ	37				5
Lemnaceæ	90				29
emna gibba Blanco	90	672	468	3:78	29
igustrum quadriloculare Blanco	69	10	7	1:14	16
iliaceæ	87		l		26
Limonia corymb^sa Blanco	28		251	2:102	3
disticha Blanco	28	356		2:102	3
glutinosa Blanco	28	358			3
laureola Blanco	19		251	2:101	1
linearis Blanco	28	357			3
monophylla Linn	28		252	2:103	3
trifoliata Linn	28	357	252	2:103	3
Liparia badocana Blanco	37	597	l	l	5
Litsea luzonica Blanco	69		284	2:162	16
Llanosia toquian Blanco	21		319	2:225	î
Oraniacese	61				13
Lonicera symphoricarpos Blanco	74	161			18
oranthaceæ	74				18
Loranthus pauciflorus Blanco	74	235			18
philippinensis Cham and Schlecht	74	200	164	1:296	18
tomentosus Blanco	74		164	1:296	18
umanaja fluviatilis Blanco	78	821	568	3:236	19
unasia amara Blanco	27	783	500	3: 191	1 2
uninus angustifolius Rlango				9. 191	
upinus angustifolius Blanco ycopodiaceæ	38	566			- 6
ycopodiaceeycopodium dichotomum Blanco	93				32
	93		576	3:240	32
gnidioides Blanco	83	824	569	3:239	32
ythraceæ	47				
Iacanea arborea Blanco	16	431			
Iaesa membranacea Blanco	57		590	1:164	12
Magnoliaceæ	15				
Magnolia angatensis Blanco	15	859	328	2:243	
falaisia tortuosa Blanco	78	789	543	3:196	19
Malesherbia globosa Blanco	43	l	454	3:54	35
Mallococca parva Blanco	26	443			:
Malphigiaceæ	26				ì
dalvaceæ	23		L		3
	20	,			

Name.		Blan	Novis sims		
Name.	work.	First	Second		Apper dix.
		edition.	edition.	edition.	uix.
[alva coromandeliana Blanco	23	551			
luzonica Blanco	23		385	2:385	1 :
moschata Blanco	26	550	235	2:344	:
lamboga capitata Blanco	52	140			1
Iangifera altissima Blancoanisodota Blanco	35	181	129	1:230	
anisodota Blanco	35		129	1:229	
indica Linnpinnata Blanco	35 36	179 129	127	1:229 1:231	
rostrata Blanco	35	123	129	1:231 $1:231$	
Ianungala pendula Blanco	29	306	123	1.201	
aranta arundinacea Blanco	84	7	5	1:9	2
[archantia	95	835	577	3:256	
farrubium indicum Blanco	69	477	332	2:250	1
Iarsdena parasita Blanco	61	120	86	1:156	1
akkar Blanco	60	118	85	1:155	1
tagudina Blanco Iarsileaceæ	60 93	121	86	1:157	3
Iarsilea minuta Blanco	93	834	577	3:253	8
trifolia Blanco	93	835	577	3: 254	8
[atricaria chamomilla Blanco	56	631	440	3: 22	ľ
[elaleuca grandiflora Blanco	24	615			1
[elastomataceæ	46				1
Ielastoma aspera Blanco	46	368			Ì
dodecandra Blanco	47		258	2:116	j
malabarica Blanco	47	367	258	2:115	
obvoluta Blancotamonea Blanco	46 47	367	259	2:117	
[eliaceæ	30	307			ł
felia azedarach Blanco	30	345			1
composita Blanco	30	010	241	2:84	1
iloilo Blanco	31		241	2:85	1
Ielicocca triptera Blanco	35		203	2:16	1
felicope conferta Blanco	28		205	2:19	1
tetandra Blanco	27	293			1
fallococca crenata Blanco felochia arborea Blanco	26	442 524	365	2:311	1
supina Linn	25 25	524	856	2:311	
femecylon lanceolatum Blanco	47	301	209	2:26	1
parviflorum Blanco	47	300			
tinetorium Blanco	47		208	2:26	1
Ienais mollis Blanco	62	139	99	1:182]
[enispermaceæ	17				
Ienispermum cocculus Linn	17	809	557	3:216	ł
rimosum Blanco fentha auricularia Blanco	17 70	810	558 332	3:217 2:250	:
cablin Blanco	70	472	332	2.200	
crispa Blanco	70	474	330	2: 246	
Ienyanthes indica Bory	61	87	63	1:118	
fetrosideros pictapetala Blanco	45		295	2:183	1
lezoneurum procumbens Blanco	41		235	2:73	
Iichelia champaca Linn	15	462	322	2:232	1
fillingtonia pinnata Blanco	66	501	351	2:285	1 :
quadripinnata Blanco Iimosa acle Blanco	66	501	351	2:286	:
asperata Blanco	42	738 732	509 505	3:140	
carisquis Blanco	42	734	507	3:134 3:137	1
coriaria Blanco	42	734	506	3:136	i
coriaria Blancoscutifera Blanco	43	735	507	3:138	1 .
farnesiana Linn	42	729	504	3:155	1
lebbek Blanco	42	733	506	3:135	1
membranulacea Blanco	41	739			-{
peregrina Blanco	42	737	509	3:139	
punctata Blanco	42		508	3:139	1
quadrivalvis Blancotenuifolia Blanco	42	732	506	3:135	
tenuifolia Bianco	42 43	739 731	510	3:141	
unguis-cati Blanco	43	731			1
virgata Blanco fimulus violaceus Blanco	65	101	357	2:290	
fimusops talosan Blanco	25	284	198	2:6	1 '
firabilis longiflora Blanco firoxylum dicline Blanco focanera grandiflora Blanco	70	77 813	57	1:109	

Name.	This	Blan	Novis- sima		
rome.	work.	First edition.	Second edition.	Third edition.	Appe dix
Mocanera guiso Blanco		449			
malaanonan Blanco	22	858			
mangachapoi Blanco	$\frac{22}{22}$	450			
mayapis Blanco	22 21	449			
plagata Blanco	21 22	447			
polysperma Blanco	22	448			
thurifera Blanco	22	446			
vernicifluua Blanco	21	450			
Iodecca coccinea Blanco	49		453	3:53	I
parviflora Blanco	48		453	3: 52	
saponaria Blanco	49		453	3:53	
tríloba Blanco	49		452	3:52	
logorium aculeatum Blanco	58	9	7	1:13	1
Iolinaea arborea Blanco	34	292	204	2:18	
Iollugo stricta Linn	50	52	35	1:64	1
subserrata Blanco	50	51	34	1:63	ĺ
Iomordica balsamina Linn	49	768	529	3:271	1
cylindrica Blanco	49	769	530	3:271	
operculata Blanco	49	770	530	3:173	
sphaeroidea Blanco	49	771	531	3:174	1
Ionodora myristica Blanco Iorinda citrifolia Blanco	16		300	2: 193	1
lorinda citrifolia Blanco		149	300	2.155	
ligulata Blanco	54	143	105	1:196	1 1
litoralis Blanco	54		109	1:190	
royoc Blanco	. 54	148	105]]
umbellata Blanco	54	140		1:193	
oringacee	54		110	1:197	-
loringa oleifera Lam	36	341		0.00	
lorus alba Linn	36	705	238	2:80	١.
luzonica Planco	78		489	3:109	
luzonica Blanco	78	703]
tinctoria Blanco	80	704			2
Ioschosma tenuiflorum Blanco	69	481	335	2:255	1
Iunchausia speciosa Linn	47	611	427	2:413	
Iurraya cerassiformis Blanco	28	363			l
exotica Blanco	28		255	2:110	1.
lobata Blanco	28	363	256	2:111	j
odorata Blanco	28		256	2:111	Ι΄.
Iusa paradisiaca Linn	82	239	168	1:303	1 :
cinerea Blanco	83	250	175	1:313	:
compressa Blanco	83	240	168	1:304	1 :
glaberrina Blanco	83	245	171	1:308	1 2
glauca Blanco	83	250	175	1:312	1 :
lacatan Blanco	83	243	170	1:305	
longa Blanco	83	245	171	1:308	! :
magna Blanco	83	244	171	1:307	
maxima Blanco	83	245	171	1:308	1 :
pumila Blanco	83	244	171	1:306	
subrubea Blanco	83	242	171	1:307	
suaveolens Blanco	83	244	171	1:306	
ternatensis Blanco	83	243	171 170	1:305	
tombak Blanco	83	246	171	1:309	
ulnaris Blanco	83	246	172	1:309	
violacea Blanco	83	245	171	1:307	
troglodytarum Linn	83	246	172	1:310	3
dolioliformis Blanco	83	855	174	1:312	
errans Blanco	83	247	174 172	1:310	
textoria Blanco	83	247	173	1:310	
ussaenda frondosa Blanco		167	118	1:311 $1:211$	
yriotheca arborescens Blanco	53				
vristicacea	94	831	575	3: 248	
yristicaceæyristica luzonica Blanco	73		462	3:69	
	73	664	402	5.09	
[yrsinaceæ	56				1 .
Tyrtaceæ	45	100		0.100	1
lyrtus communis Blanco	45	422	295	2:182	
mananquil Blanco	45	421			
subrubens Blanco	46		294	2:182	1
tripinnata Blanco	46	421		i	1
aiadaceæ	90				2
aias lobata Blanco	90		459	3:65	1
obvoluta Blanco	94		460	3:66	1
palustris Blanco	90	660			1 2
ama jamaicensis Blanco	96	211	148	1:266	

Name.		Blan	Novis sima		
Name.	work.	First edition.	Second edition.	Third edition.	Appendix.
Vapaea latifolia Blanco	23		387	2: 346	2
scabra Blanco	23 52	553	386 102	2:846 1:188	2
Vauclea adina Blanco	52 52		102	1:188	10 10
calycina Blancodigitata Blanco	51		102	1:188	10
glaberrima Blanco	52		100	1:141	10
glabra Blanco	52		101	1:185	liŏ
glandulosa Blanco	52	143			10
lanceolata Blanco	52	144		<u></u>	10
latifolia Blanco	52	144			10
lutea Blanco	52	141			10
luzonensis Blanco	52 52		102 101	1:187	10
obtusa Blanco	39	588	410	1:187 2:388	10
pruriens Blanco	38	900	411	2:389	1 8
urens Blanco	38	586	409	2:387	6
Velumbium speciosum Willd	17	1 000	318	2:223	١,
turbinatum Blanco	17	458			
Vepenthaceæ	72				17
Vepenthes alata Blanco	72	805	555	3:214	17
ventricosa Blanco	72	807	556	3:215	17
Verium oleander Lour	59	104	75	1:140	18
Vicotiana frutescens Blanco	64	101			14
fruticosa Blanco	64		74	1:138	14
pusilla Blanco	64	100	74	1:137	14
tabacum Linn	64	101	74	1:138	14
Viota globosa Blanco	77 29		214	2:35	19
tetrapetala Lam Vipa fructicans Wurmb	29 88		213	2:35	28
littoralis Blanco	88	662	461	3:68	28
Vyctaginaceæ	70	002			16
Vyctanthes sambac Linn	58	10	6	1:12	12
lymphaeaceæ	17	10		1.12	14
lymphaeaceæ lymphaea lotus Linn	17	456	317	2:222	8
Ochnaceæ	29		01.		ì
Ochna fascicularis Blanco	29		245	2:92	
Ocymum album Blanco	69	479			10
americanum Blanco	69	480	335	2:254	16
citrodorum Blanco	69		591	2:256	10
flexuosum Blanco	69	481	335	2:255	10
sanctum Linn	69	480	334	2:254	10
virgatum Blanco	69		334	2:253	10
DlacaceæDlax baticulin Blanco	32			1.90	13
Oldenlandia capensis Blanco	74 52	62	589 45	1:38 1:82	10
affinis Blanco	52 52	02	44	1:83	1
paniculata Linn	52	61	44	1:82	10
)leaceæ	58	01	**	1.02	1
mphalobium pictum Blanco	37		271	2:139	
)nagraceæ	48				1 9
Ophiorrhiza oblongifolia DC	53		65	1:121	10
triandrum Bianco	65		65	1:122	1
ophrys cernua Blanco	85	639	445	3:35	2.
Orchidaceæ	84				2:
Orchis lanigera Blanco	84	641	446	3:37	2
Prixa ternata Blanco	27	62	45	1:84	1
Ormosia calavensis Blanco	41		230	2:64	! !
Ornitrophe triandra Blanco	35	291			
OrobanchaceæOryza sativa aristata Blanco	65 92	274	190	1:339	3
hinambang Blanco	92 82	274	189	1:339	3
binambang Blanco glutinosa Blanco	92 92	273	190	1:339	3
lamuyo Blanco	92	273	190	1:339	3
pilosa Blanco	92	275	191	1:341	3
praecox Blanco	92	274	190	1:340	3
quinanda Blanco	92	274	191	1:340	3
rubra Blanco	92	285	191	1:341	3
violacea Blanco	92	275	191	1:343	3
			205	2:20	
Sbeckia chinensis L	46				
	46 46 82	293 461	321	2:230	2

Name		Blar	Novis sims		
Name.	This work.	First edition	Second edition.	Third edition.	Appe
Oxalis acetosella Blanco	27	388	272	2:141	
sensitiva Blanco	27	389			-
Pachyrhizus angulatus Rich	40		405	2:380	ł
jicamas Blanco montanus Blanco	40 39	579	406	2:381	
teres Blanco	39	580			
Paederia foetida Blanco	54	159	112	1:201]
tacpo Blanco Palaquium lanceolatum Blanco	54 57	160 403	113 282	1:202 2:159	
latifolium Blanco	57	405	282	2:159	1
oleiferum Blanco	57	405	283	2:160]]
Paliurus dao Blancodubius Blanco	36 29	174 175	122 123	1:219 1:121	
edulis Blanco	36	173			l
lamio Blanco	$\frac{36}{29}$	174	122	1:218 1:220	
perforatus Blanco	88 88	1/4	122	1:220	2
anax fruticosa Blanco	51		156	1:281	j
Pancratium amboinense Linn	- 86	252	177	1:317	1 3
illyricum Blanco maritimun Blanco	86 86	251 252	176 177	1:315 1:316	
zeylanicum Linn	86	253	177	1:317	1 :
Pandanaceæ	89				1 1
Pandanus exaltatus Blanco gracilis Blanco	89 89	778 778	536 536	3: 183 3: 182	
inermis Blanco	89	110	537	3:184	:
malatensis Blanco	89		536	3:182	2
radicans Blancosabotan Blanco	89 89	780 779	537 537	3:184 3:183	
spiralis Blanco	89	777	535	3: 181	
Panicum miliaceum Blanco	91	39	28	1:52	
PapaveraceeParatropia crassa Blanco	17 51		158	1:285	
obtusa Blanco	51		159	1:285	:
Paspalum villosum Blanco	91	40	28	1:53	8
PassifloraceæPassiflora coccinea Blanco	48 49	650			
minima Blanco	48	647			
parviflora Blanco	48	649			
saponaria Blancoserrulata Jacq	49 48	650	452	3:50	
zucca Blanco	49	648	402	3.50	
Pavetta membranacea Blanco	54	59			
sumbueina Blanco Pedaliaceæ	54 66		41	1:75]
Pemphis setosa Blanco	47	410	285	2:164	
Pentapetes cebuana Blanco	25	529	369	2:319	
coccinea BlancoPerdicium tomentosum Blanco	25 55	529 630	439	3:21	. :
Pergularia glabra Blanco	60		141	1:254	
glandulosa Blanco	60	201	141	1:254	
procumbens Blanco Petaloma alba Blanco	60 44	201 344	240	2:82	
coccinea Blanco	44	345	240	2:83	
Phalaenopsis amabilis Blume	84		592	3:41	:
Phaseolus bulai Blancocaracalla Linn	37 39	572 575	401	2:372	
ilocanus Blanco	39	572	401	2.012	
inamoenus Linn	39	571	399	2:368	
lunatus Linn	39 3 9	573 573	400 400	2:370 2:370	1.5
mungo Linn tunkinensis Lour	39	913	399	2:369	
tunkinensis Lour vexillatus Blanco	39	574			
vulgaris Blanco	39	AP7 4	401 330	2: 371	:
Phlomis alba Blanco zeylanica Blanco	70 70	474 475	330	2:247 2:248	
Phyllanthus carolinianus Blanco	76	691			
kirganelia Blanconiruri Blanco	76		480	3:96	
niruri Blanco tetrander Blanco	76 76	690	480	3: 117	
Physalis angulata Linn	64	138	99	1:181	
peruviana Linn	64	138	98	1:180	1 :

Name.	This	Blan	Blanco's Flora de Filipinas.		
Trume.	work.	First edition.	Second edition.	Third edition.	Appen- dix.
Physalis pubescens Linn	64	138	98	1:181	144
PHOCATOUS amara Blanco	27	100	540	1.101	35
Pinus taeda Blanco	82	767	528	3:169	212
Piperaceæ	72				174
Piper anisodorum Blanco	72		16	1:31	176
anisumolens Blancobetel Blanco	$\frac{72}{72}$	23 22	16	1:30	176 175
nigrum Linn	72	21	16	1:29	175
nigrum Linnobliquum Blanco	72	22	16	1:30	175
parvifolium Blanco	72	23	17	1:32	175
Pisonia aculeata Linn	70	195	137	1:247	167
Pistia stratiotes Linn Pittosporaceæ	90 18	651	454	3:55	296
Pittosporaceæ Plagianthus humilis Blanco	30	526	366	2:315	13 41
Plantaginaceæ	70	320	300	2.010	166
Plantago crenata Blanco	70	56			166
media Blanco	70		38	1:70	166
Plinia paniculata Blanco	21	423	296	2:184	17
Plumbaginaceæ	56			1,111	122
Plumbago rosea Linnviscosa Blanco	56 56	78 78	58	1:111	122 122
Plumieria alba Blanco	59	111	58 80	1:111 1:148	129
Plumieria alba BlancoPoinciana pulcherrima Linn	41	333	232	2:69	69
Polanisia viscosa DC	17		364	2:308	10
Polemonium obscurum Blanco	63	103	75	1:139	142
Polyanthes tuberosa Linn	85	259	181	1:323	254
Polycarpon pholyphyllum Blanco	19	53	36	1:66	15
Polygalacee Polygala monspeliaca Blanco	19 19	557	388	2:350	14 14
Polygonaceæ	71	997	300	2:350	172
Polygonum bellardi Blanco	71	314	219	2:45	172
stoloniferum Blanco	71	314	219	2:45	172
Polyozus bipinnatus Blanco	51		43	1:79	103
Polypodium adiantiforme Blanco	94		573	3:245	338
phymatodes Linn	94	827 827	572	3:242	340
quereifolium Linnserratum Blanco	94 94	827	572 572	3:244 3:243	340 337
signatum Blanco	95	021	572	3:243	340
Polyscias digitata Blanco	51	224			102
disperma Blanco	51	226			102
obtusa Blanco	51	226			102
odorata Blanco	51	225			102
Pongatium spongiosum BlancoPontederiaceæ	56 87	86			121 267
Pontederia vaginalis Blanco	87	255	178	1:320	267
Porana volubilis Burm	63	88	64	1:119	143
Portesia rimosa Blanco	31	297			43
Portulacaceæ	19				15
Portulaca axilliflora Blanco	50 19		285	2:163	100
meridiana Linn oleracea Linn	19	406 407	285 284	2:163 2:162	15 15
toston Blanco	50	408	204	2.102	100
Pothos pinnata Blanco	90	646	450	3:49	289
Poupartia pinnata Blanco	36	392	275	2:146	55
Premna cordata Blanco	68	489			159
integrifolia Blanco	68		342	2:268	159
nauseosa Blancoodorata Blanco	68 68	489 488	341	2:268	159 159
serratifolia Blanco	68	400	342	2:269	159
tomentosa Blanco	68		342	2:269	159
Prockia albicans Blanco	77	430	299	2:192	194
Procris erecta Blanco	80	707	490	3:111	203
violacea Blanco	80	706	490	3:110	203
Prunus armeniaca Blanco	43		296	2:185	76
Psidium aromaticum Blanco pyriferum Blanco	45 45	714	292	2:178	83 83
Psoralea badocana Blanco	37		416	2:178	58 58
Ptelea arborea Blanco	35	63	710	2,000	53
Pteris grandifolia Blanco	94	829	574	3:246	331
piloselloides Blanco	94	830	574	3:248	337
sinuata Blanco	94	830			338
signata Blanco	95	830	l	l	340

Name.	This	Blan]	Novis- sima		
	work.	First edition.	Second edition.	Third edition.	Appen dix.
Pteris trichomanoides Blanco	94	830	574	3:247	38
Pterocarpus diadelphus Blanco	40	563	393	2:357	3
frutescens Blanco	40	562	392	2:356	ì
pallidus Blanco	40	560	391	2:355	è
santalinus Blanco	. 40	561	392	2:356	. (
Pterospermum hastatum Blanco	25	528	367	2:317	:
obliquum Blanco	25	529			:
semisagittatum Blanco	25		368	2:318	
Punica granatum Linn	47	422	295	2:184	9
Pycnanthemum decurrens Blanco	69		333	2:251	16
elongatum Blanco	69		333	2: 252	10
subulatum Blanco	69 34		333 247	2: 251 2: 94	16
Quassia simaruba Blanco	34 34	351	247	2:94	
tricarpa Blanco tricarpa Blanco	34	351	246	2:94	3
Quercus cerris Blanco	81	727	503	3:130	20
concentrica Blanco	81	121	503	3:129	20
cooperta Blanco	81		503	3:130	20
glabra Blanco	81	727		0.100	20
molucca Blanco	81	726			20
ovalis Blanco	81		502	3:129	20
Quilamum luteum Blanco	47	851	136	1:245	1 -3
Quilesia sericea Blanco	32	177	125	1:224	
Quirosia anceps Blanco	37		398	2:367	1 8
secunda Blanco	37		398	2;366	8
Quisqualis indica Linn	45	361			8
spinosa Blanco	44		254	2:109	1 8
Rafflesia lagascæ Blancophilippinensis Blanco	72		595	2: 231 3: 280	17
philippinensis Blanco	72		365		1'
Randia aculeata Blanco	53	141	99	1:183	10
Ranunculaceæ	15		233		١.
Reichardia pentapetala Blanco	41		233	2:71	1 .
Reichelia palustris Blanco	56	220	155	1:227	15
Remijia angatensis Blancoobscura Blanco	53 53		115 116	1:206 1:207	10
odorata Blanco	53		115	1:205	1 10
Renealmia gracilis Blanco	84	1	110	1.200	25
exaltata Blanco	84	1 *	1	1:2	25
Rhamnaceæ	32		•	1	- 4
Rhamnus carolinianus Blanco	33	169	119	1:214	1
dalanta Blanco	33	170			. 4
jujuba Linn	32	172			. 4
talanai Blanco	33	170			. 4
zizyphus Blanco	33	171			. 4
zonulatus Blanco	33	172			. 4
Rhaphis trivalvis Lour	92	45			3
Rheum muricatum Blanco	72	327			1'
Rhizophoraceæ	43				
Rhizophora candel Blanco	44	396	277	2:150	1 :
gymnorrhiza Linnlongissima Blanco	44	555	276	2:149	
nongissima Bianco	43	398	278	2:151	
mangle Linn	43	397	278	2:151	
plicata Blanco	44 44	398 396	279 277	2:152 2:277	
polyandra Blanco			211	2.211	
tinetoria Blanco Riana tricapsularis Blanco	44 32	394 850	126	1:225	
Ricinus communis Linn	78	761	523	3:161	1
Ronabea arborea Blanco	53	101	114	1:204	1
bipinnata Blanco	53	162			1
Rondeletia asiatica Blanco	52	146			1
Rosaceæ	43				.]
Rosmarinus officinalis Linn	70	20	15	1:28	1
Rubiaceæ	52		l _		1
Rubus comintanus Blanco	43		298	2:190	1
idaeus Blanco	43	428	298	2:190	1
jamaicensis Blanco	43	427			
moluccanus Linu	43	428	298	2:190	
Ruellia contorta Blanco	67	496	347	2:278	1
repens Blanco	66	493	345	2:276	1
secunda Blanco	67	495	348	2:278	1
uliginosa Blanco	66	494	346	2:277	1
Rumex muricatus Blanco	72	1	195	1:346	1

	This	Blan	Novis sima			
Name.		First edition.	Second edition.	Third edition.	Appe dix.	
Rutaceæ	27					
Saccharum koenigii Blanco	92	44	30	1:56		
officinarum Linn	92	42	29	1:55	1	
Salacia sinensis Blanco	32	26	19	1:36	`	
Salceda montana Blanco	$\tilde{21}$		374	2:327		
Salgada lauriflora Blanco	$\overline{73}$		221	2:50	1 1	
Salicaceæ	81				1 2	
Salix azaolana Blanco	81		539	3:188	1 2	
Salvadoraceæ	58		000	0.100	;	
Salvia violacea Blanco	70		14	1:27	1 1	
Sambucus javanica Blume	51		151	1:271	1 3	
Samydaceæ	48		101	1.11	1 -	
Samyda pubescens Blanco	48		263	2:124		
serrulata Blanco	48	374				
trialvis Blanco	48	374	263	2:124	1	
Sandoricum indicum Cav	31	0.1	242	2:85	1	
ternatum Blanco	31	346		2.00	1	
Sapindaceæ	34	0.0				
Sapindus baccatus Blanco	35	290				
edulis Blanco	34	230	201	2:12		
guisian Blanco	34		201	2:11		
koelreuteria Blanco	34	289	201	2.11	1	
saponaria Blanco	34	288			İ	
Sapotacee	57	200			1 :	
Sapota nigra Blanco	58	409				
Saxafragaceæ	43	103	1		1 .	
Scaevola lobelia Blanco	56	147	104	1:193		
Schmidelia conferta Blanco	34	141	217	2:41		
triandra Blanco	35		218	2:42	-	
Schotia speciosa Blanco	41	356	251	2:100		
Scitimineæ		350	201	2:100		
Scirpus niloticus Blanco	82 91	33	23	1:43	1 :	
Sclerotium subterraneum Blanco	96				١ '	
Zoonorio dulois Linn		845	584	3: 266 1: 69		
Scoparia dulcis Linn	65	55	37	1:09		
Scrophulariaceæ	64					
Sebifera glutinosa Lour	73	819	566	3: 233		
balongai Blanco	74	820	567	3: 235		
Securidaca complicata Blanco	19	556	388	2: 349		
volubilis Blanco	19		150	1.075		
Semecarpus anacardium Blanco	36	217	152	1:275	1	
cuneiformis Blanco	36	220	155	1:276	1 .	
Senecio caliaster Blanco	56		441	3: 24		
Seringia lanceolata Blanco	35		45	1:85	1 .	
Serissa myrtifolia Blanco	53	164				
pinnata Blanco	53	163				
Serratula multiflora Blanco	54	617	431	3:7		
Sesamum indicum Linn	66	507	353	2: 290		
Sesbania cannabina Blanco	38		418	2:400		
grandinora Pers	38	599	418	2:399		
sesuvium portulacastrum Linn	50	426	297	2:187		
sida irutescens Bianco	23		384	2:341		
indica Linn	23	547	283	2:339	1	
lanceolata Roxb	23	548			.	
philippica DC	23		283	2:340		
semicrenata Link	23		. 348	2:341	1	
truncatula Blanco	23	548				
Sideroxylon balitbitan Blanco	57	130	92	1:169	1	
duclitan Blanco	57	129	92	1:168		
Simarubaceæ	29				4	
Sinapis brassicata Linn	17	219	362	2:305	1	
Junea Linn	17		362	2:304		
Sillensis Riengo	17	520			.1	
Smilax divaricata Blanco	86	795	548	3:203		
	87	796	549	3: 205		
latifolia Blanco	87		548	3:204	İ	
latifolia Blanco pseudochina Blanco smithia higeminata Blanco	87	795	548	3: 204		
	38		395	2: 362		
Soala litoralis Blanco	16	437	304	2: 199		
	63	101				
Solanum coagulans Blanco	64	135	97	1:177	1	
lycopersicum Linn	63	134	96	1:176		
mauritianum Blanco	64	134	96	1:176		
21996	1 04	Total	. 50			

Name.	This	Blan	Novis-		
	work.	First edition	Second edition.		Apper dix.
Solanum melongena Linn	64	135	96	1:177	14
nigrum Linn	64	134	96	1:175	ī
nigrum Linnserratum Blanco	64		97	1:179	1
sinense Blancotuberosum Linn	64 64	137	97	1:179	1
zeylanicum Blanco	64	136	97	1:178	1
Sonneratia pagatnat Blanco	47	424	296	2:186	1
pathodea luzonica Blanco	66		350	2:284	1
pathodea luzonica Blanco pergula serrata Blanco permacoce muriculata Blanco	19 54		271 44	2:140 1:81	1
mutilata Blanco	54		43	1:80	î
Sphaeranthus alatus Blanco	55	635		l	î
elongatus Blanco	55	636	443	3:30	1
indicus Linn	55 56		442 62	3:29 1:117	1
Sphenoclea zeylanica Gaertn Sophora heptaphylla DC	40	329	229	2:63	1
tomentosa Linn	40	328	229	2:63	
Spilanthes acmella Blanco	55	620	433	3:10	1
lobata Blanco	55	622	434	3:11 3:11	1
peregrina Blanco Spinifex squarrosus Linn	55 91	622 46	434 31	1:57	1 3
Spondias dulcis Blanco	36	390	273	2:143	
Spondias dulcis Blanco Stachys artemesia Lour	70	476	331	2:249	1
Stalagmites cambogioides Blanco	20		301	2:195	
Stapelia meliflua Blancoquadrangula Blanco	60 60	202 202	142 142	1:256 1:255	1
Stemodia ruderalis Blanco	64	498	346	2: 348	ĺ
Sterculiaceæ	24		!	!	i
sterculia alata Blanco	24		525	3:165	
Sterculia balanghas Blancocordifolia Blanco	$\frac{24}{24}$	765 764	525 525	3:165 3:163	
cymbiformis Blanco	24	704	526	3:167	
decandra Blanco	24, 73	766	526	3:166	27, 1
foetida Linn glandulosa Blanco	24	763	524	3:162	
glandulosa Blanco glomerata Blanco	24, 73	764	525	3: 164 3: 164	27, 1
Stigmarota edulis Blanco	$24,73 \\ 18$	764	560	3: 221	21,1
Stilago bunius L	76	782	539	3:189	1
Stipa spinifex Linn	91	41	29	1:54	1 8
Strychnos philippensis Blanco	61 53		61	1:116 1:209]]
Stygmanthus cymosus Blanco Styraceæ	58		117	1.203	. i
Sulipa globosa Blanco	46		348	2:280	
pseudopsidium Blanco	53		347	2:280	
Syzygium jambolanum DClatifolium Blanco	46 46		293 294	2:180 2:181	
Tabernaemontana cirrhosa Blanco	59	115	83	1:152	
elliptica	59	115	83	1:152	
globosa Blanco	59 59	116	83 82	1:153 1:150	
laurifolia Blancopolygama Blanco	59	114	82	1:150	1 :
Taccaceæ	86			1.101	
Tacea gaogao Blanco	86	262, 856			.) :
palmata Blume	86		182	1:325	
pinnatifida Forstvesicaria Blanco	86 86	261	182	1:327	
Tagetes patula Linn	56	632	440	3:23	
Tala odorata Blanco	64	485	338	2:262	
Taligalea umbellata Blanco	54 41	29	337	2:261 1:39	
Tamarindus indica LinnTapogomea rubra Blanco	52		20	1:59	
Tayotum nigrescens Blanco	61	105	76	1:141	1
Tayotum nigrescens Blanco Tectona grandis Linn. f Terminalia angustifolia Blanco	67	130	92		
Terminalia angustifolia Blanco	. 44	377		0.105	-{
edulis Blancolatifolia Blanco	. 44		_ 265	2:127	
mauritiana Blanco	44		264	2:126	-
Ternstroemiaceæ	. 21				-1
Tetracera monocarpa Blanco	. 15	459		-	-
sarmentosa Blanco Tetragonolobus simplicifolius Blanco	15		320	2: 227 2: 364	

Name.	This work.	Blar	Novis-			
Areally.		First edition.	Second Thir edition.		d Appen-	
Theobroma cacao Linn	25	601	410	(). 100	20	
Thespesia banalo Blanco	23	601	419 382	2:403 2:338	29 25	
populnea Corr	23		381	2:337	25	
sublobata Blanco	23		382	2:338	25	
Thoa edulis Blanco	82		514	3:146	211	
pendula Blanco	82	746	02.	0.110	211	
Thunbergia fragrans Blanco	66		360	2:301	152	
subsagittata Blanco	66	518			152	
Thymelaeaceæ	74				182	
Thymus biserratus Blanco	69	478			164	
virginicus Blanco	• 69	478			164	
Ticorea aculeata Blanco Tiliaceæ	34	86			349	
	25				29	
Tillaea rubella BlancoTillandsia pseudo-ananas Blanco	19 89	75 853	56	1:106	15	
Torenia paniculata Blanco	65	486	162 339	1:292	286	
soriana Blanco	65	400	266	2:265	148	
Tournefortia arborea Blanco	62	129	91	1:167	148 138	
hirsutissima Blanco	62	128	91	1: 167		
Tovomita pentapetala Blanco	20	432	301	2: 194	138 17	
Tradescantia cristata Jacq	87	231	163	1:293	272	
discolor Smith	87	232	163	1: 294	278	
discolor Smith geniculata Blanco	87	232	100	1:294	272	
Tragia bracteata Blanco	78		481	3:94	196	
innocua Blanco	77		479	3:94	194	
Tremella	95	837	578	3:258		
Tribulus lanuginosus Blanco	26	350	245	2:91	32	
Trichilia pentandra Blanco	31	355	249	2:97	42	
rimosa Blanco tripetala Blanco	31		250	2:99	48	
tripetala Blanco	31	354	248	2:97	49	
volubilis Blanco	31		249	2:98	44	
Trichosanthes amara Blanco	49	774	533	3:178	96	
Triopteris jamaicensis Blanco polyandra Blanco	$\frac{26}{25}$	379 380	267 268	2: 133 2: 133	32	
Tristillateia malintana Blanco	26	380			30	
Tristillateia malintana Blanco Triumfetta semitriloba Linn	26 26	406	$\frac{267}{283}$	2: 132 2: 161	32 31	
Turraea decandra Blanco	30	347	200	2. 101	42	
octandra Blanco	31	389	244	2:89	42	
virens Blanco	30	000	243	2:88	42	
Typhaceæ	90				297	
Typha angustifolia Linn	90	687	477	3:91	297	
Ugena alba Blanco	95	823	569	3:238	345	
semihastata Blanco	95	822	568	3: 237		
Ulva compressa Blanco intestinalis Blanco	95	842	581	3:261		
intestinalis Blanco	95	842	5.2	3:262		
reticulata Blanco	95	842	582	3:262	-	
umbilicalis Blanco Umbelliferæ	95	842	581	3:261		
Umbelliferæ Unona cabog Blanco	50 16	466			100	
camphorata Blanco	16	468	326	2: 239	5	
cauliflora Blanco	16	408	326	2:239	1	
corniculata Blanco	16	469	326	2:240	ļ ;	
dehiscens Blanco	16	466	325	2:240		
latifolia Blanco	16	400	323	2:236	5 6 5 5	
odorata Blanco	16		325	2:237	}	
odoratissima Blanco	16	467	325	2:239	Ì	
ossea Blanco	16	467			Ì	
setigera Blanco tripetala Blanco	15	468			4	
tripetala Blanco	16	i	324	2:236	7	
uncinata Blanco	16		324	2:237		
Urena multifida Blanco	23	540	378	2:332	28	
Urticaceæ	78				19	
Urtica arborescens Blanco	81		483	3:100	205	
baccifera Blanco	81	695			205	
capitata Blanco	80.		483	3:101	208	
ferox Blanco japonica Blanco	80		484	3:102	204	
nivea Linn	81	694	482	3:99	205	
sessiliflora Blanco	81 80	697 696	484	3:102	205	
umbellata Blanco	80	696			203	
			483	3:99		
villosa Blanco	81	695			205	

Name.	This work.	Blar	Novis-		
		First edition	Second edition.		Apper dix.
Uvaria lanotan Blanco	16	464			
ossea Blanco	16		322	2:233	
setigera Blanco	15		323	2:234	
sinensis Blanco	16	465		2.201	
tripetala Blanco	16	645			
Vallea cafomala Blanco	26	439			3
Vallisneria sphaerocarpa Blanco	82	780	538	3:186	21
spiralis Linn	82	781	538	3:187	21
Vandellia diffusa Blanco	65	505	000	0.107	14
multiflora Blanco	65	505			14
soriana Blanco	65	506		2 266	
Vangueria stellata Blanco	53	167	117	1:210	1- 11
Vanilla majaijensis Blanco	85	107	593	3:43	$\frac{11}{24}$
ovalis Blanco	85		448	3:42	
Vatica apteranthera Blanco	22		281	2:156	24
mangachapoy Blanco	22	401	201	2:100	2
sinensis Blanco	$\frac{22}{22}$	401	280	2:156	2
Ventilago monoica Blanco	32				2
Verbenaceæ	67		124	:223	4
Verbena capitata Forsk	67	19		1.00	15
Vinca rosea Linn	58	116	14	1:26	15
Vitex altissima Blanco	68	516	80	1:154	12
geniculata Blanco	68	514	359	2:299 2:299	16
latifolia Blanco	68	514	358		16
leucoxylon Blanco		514	358	2:298	16
ropons Planco	68		359	2:300	10
repens Blanco trifolia Linn. f	68	513	358	2:297	15
Volkameria casopanguil Blanco	68	513	358	2:297	16
inormic Planco	69		356	2:294	16
inermis Blanco grandiflora Blanco	69	511	5		10
Waltheria americana Linn	68	512	357	2:295	16
Wendlandia exserta Blanco	25	523	364	2:309	1
Williagh air deserta Bianco	52		104	1:191	10
Willughbeia drupacea Blanco	57	132	94	1:173	12
multilocularis Blanco	59	131	94	1:172	13
multilocularis Blanco	20	131	94	1:172	34
Xeranthemum staehelina Blanco	55	629			11
Xylocarpus granatum Koenig	31	298	207	2:24	4
Zea mays Linn	91	686	476	3:90	31
Zizyphus dalanta Blanco	33		121	1:217	. 4
jujuba Lam	32		120	1:215	4
latifolia Blanco	33		121	1:217	4
lotus Blanco	33			1:216	4
zonulatus Blanco	33		120	1:215	4
Zygophyllaceæ	26		1	l	;

PREVIOUS PUBLICATIONS OF THE BUREAU OF GOVERNMENT LABORATORIES.

(Continued from second page of cover.)

- No. 19, 1904, Biological Laboratory.—Some Observations on the Biology of the Cholera Spirillum. By W. B. Wherry, M. D.
- No. 20, 1904, Biological Laboratory.—I. Does Latent or Dormant Plague Exist Where the Disease is Endemic? By Maximilian Herzog, M. D., and Charles B. Hare, Serum Laboratory: II. Broneho-Pneumonia of Cattle: Its Association with B. Bovisepticus. By Paul G. Woolley, M. D., and Walter Sorrell, D. V. S. III. Pinto (Paño Blanco). By Paul G. Woolley, M. D. Chemical Laboratory: IV. Notes on Analysis of the Water from the Manila Water Supply. By Charles L. Bliss. Serum Laboratory: V. Frambeesa: Its Occurrence in Natives in the Philippine Islands. By Paul G. Woolley, M. D.
- No. 21, 1904, Biological Laboratory,—Some Questions Relating to the Virulence of Micro-Organisms, with Particular Reference to Their Immunizing Powers. By Richard P. Strong, M. D.
- No. 22, 1904, Bureau of Government Laboratories.—I. A Description of the New Buildings of the Bureau of Government Laboratories. By Paul C. Freer, M. D., Ph. D. II. A Catalogue of the Library of the Bureau of Government Laboratories. By Mary Polk, Librarian.
- No. 23, 1904, Biological Laboratory.—Plague: Bacteriology, Morbid Anatomy, and Histo-pathology (including a consideration of insects as plague carriers). By Maximilian Herzog, M. D.
- No. 24, 1904, Biological Laboratory.—Glanders: Its Diagnosis and Prevention, together with a Report on Two Cases of Human Glanders Occurring in Manila, and some Notes on the Bacteriology and Pleomorphism of Bacterium Mallei. By William B. Wherry, M. D.
- * No. 25, 1904.—Birds from the Islands of Romblon, Sibuyan, and Cresta de Gallo. By Richard C. McGregor.
- No. 26, 1994, Biological Laboratory.—The Clinical and Pathological Significance of Balantidium Coli. By Richard P. Strong, M. D.

Publications of the Bureau are given out in serial number pertaining to the entire Bureau, Publications which may be desired can be obtained by applying to the Librarian of the Bureau of Government Laboratories, Manila, P. I., or to the Superintendent of Government Laboratories, Manila, P. I.

^{*}The first four bulletins in the ornithological series were published by the Ethnological Survey under the title "Bulletins of the Philippine Museum." Future ornithological publications of the Government will appear as publications of the Bureau of Government Laboratories.

